

Teorema de Fubini

Teorema 1 (de Fubini). Sean (X_1, Σ_1, μ) y (X_2, Σ_2, λ) espacios de medida σ -finitos.

Sea $f: X_1 \times X_2 \longrightarrow \mathbb{R}$ una función $\Sigma_1 \otimes \Sigma_2$ -medible. Si definimos

$$\begin{aligned} \varphi: X_1 &\longrightarrow \mathbb{R} & \psi: X_2 &\longrightarrow \mathbb{R} \\ x &\longmapsto \int_{X_2} f(x, y) \, d\lambda(y) & y &\longmapsto \int_{X_1} f(x, y) \, d\mu(x), \end{aligned}$$

$$\implies \begin{cases} (1) & f \geq 0 \implies \varphi \text{ es } \Sigma_1\text{-medible y } \psi \text{ es } \Sigma_2\text{-medible.} \\ (2) & f \in \mathcal{L}^1(\mu \otimes \lambda) \implies \varphi \in \mathcal{L}^1(\mu) \text{ y } \psi \in \mathcal{L}^1(\lambda). \end{cases}$$

Además, en cualquiera de estos dos casos, se cumple que

$$\int_{X_1 \times X_2} f(x, y) \, d(\mu \otimes \lambda) = \int_{X_1} \varphi(x) \, d\mu(x) = \int_{X_2} \psi(y) \, d\lambda(y).$$

Referenciado en

- `lem-transformada-fourier-conmutativa-integral`
- `teo-inversion-transformada-fourier`
- `lem-transformada-fourier-convolucion`
- `desigualdad-young-convolucion`
- `teo-aproximacion-identidad-convolucion`
- `ley-fuerte-grandes-numeros`
- `lem-convolucion`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.