

Fórmula para el laplaciano en función de las derivadas de Wirtinger

Lema 1. Sea $f \in \mathcal{C}^2(\mathbb{D})$, entonces $\forall z \in \mathbb{D} : \Delta f(z) = 4 \frac{\partial^2 f}{\partial z \partial \bar{z}}(z) = 4 \frac{\partial^2 f}{\partial \bar{z} \partial z}(z)$.

Demostración: Desarrollando las derivadas de Wirtinger en el lado derecho, tenemos:

$$\begin{aligned} \frac{\partial^2 f}{\partial z \partial \bar{z}} &= \frac{\partial}{\partial z} \left(\frac{\partial f}{\partial \bar{z}} \right) = \frac{\partial}{\partial z} \left(\frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right) \right) \\ &= \frac{1}{2} \left[\frac{1}{2} \left(\frac{\partial^2 f}{\partial x^2} - i \frac{\partial^2 f}{\partial y \partial x} \right) + \frac{1}{2} i \left(\frac{\partial^2 f}{\partial x \partial y} - i \frac{\partial^2 f}{\partial y^2} \right) \right] \\ &= \frac{1}{4} \left(\frac{\partial^2 f}{\partial x^2} - i \frac{\partial^2 f}{\partial y \partial x} + i \frac{\partial^2 f}{\partial x \partial y} + \frac{\partial^2 f}{\partial y^2} \right) = \frac{1}{4} \left(\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} \right) = \frac{1}{4} \Delta f. \end{aligned}$$

Por tanto, $\Delta = 4 \frac{\partial^2}{\partial z \partial \bar{z}}$ (también se puede ver que $\Delta = 4 \frac{\partial^2}{\partial \bar{z} \partial z}$ de forma análoga). ■

Referenciado en

- lem-laplaciano-cambio-variable

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.