

Prop schwartz imp lp

Proposición 1. *La clase de Schwartz $\mathcal{S}(\mathbb{R})$ es un espacio vectorial sobre $\mathbb{C}$ contenido en $\mathcal{L}^p(\mathbb{R})$ para todo $p \in [1, \infty]$.*

Demostración: Que $\mathcal{S}(\mathbb{R})$ es un espacio vectorial es fácil de comprobar. Sea $f \in \mathcal{S}(\mathbb{R})$, entonces $\rho_{0,0} = \sup_{x \in \mathbb{R}} |f(x)| < \infty$, luego $f \in \mathcal{L}^\infty(\mathbb{R})$. Por otro lado, para $p \in [1, \infty)$,

$$\|f\|_p^p = \int_{\mathbb{R}} |f(x)|^p dx = \int_{|x| < 1} |f(x)|^p dx + \int_{|x| \geq 1} |f(x)|^p dx.$$

Como $f \in \mathcal{L}^\infty(\mathbb{R})$, la primera integral está acotada por $2\|f\|_\infty^p$. Para la segunda integral,

$$\int_{|x| \geq 1} |f(x)|^p dx = \int_{|x| \geq 1} \frac{|x|^{2p} |f(x)|^p}{|x|^{2p}} dx \leq \rho_{2,0}^p \int_{|x| \geq 1} \frac{1}{|x|^{2p}} dx = \rho_{2,0}^p \cdot 2 \int_1^\infty \frac{1}{x^{2p}} dx < \infty.$$

Por tanto, $\|f\|_p < \infty$ y $f \in \mathcal{L}^p(\mathbb{R})$. ■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.