

Teorema de diferenciación de Lebesgue para funciones continuas

Teorema 1 (de diferenciación de Lebesgue para continuas). Sea $f \in \mathcal{C}(\mathbb{R})$

$$\implies \forall x \in \mathbb{R} : \lim_{h \rightarrow 0} \frac{1}{2h} \int_{x-h}^{x+h} f(t) dt = f(x).$$

Demostración: Sea $x \in \mathbb{R}$. Entonces, dado $\varepsilon > 0$, $\exists \delta > 0$ tal que si $|x - y| < \delta$ se tiene que $|f(x) - f(y)| < \varepsilon$. Por tanto, si $0 < h < \delta$,

$$\left| f(x) - \frac{1}{2h} \int_{x-h}^{x+h} f(t) dt \right| \leq \frac{1}{2h} \int_{x-h}^{x+h} |f(x) - f(t)| dt < \frac{1}{2h} \int_{x-h}^{x+h} \varepsilon dt = \varepsilon.$$

Entonces, $\forall x \in \mathbb{R} : \lim_{h \rightarrow 0} \frac{1}{2h} \int_{x-h}^{x+h} f(t) dt = f(x)$. ■

Referenciado en

- `lem-diferenciacion-lebesgue-l1-imp-casi-toda-parte`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.