

Teorema de derivación bajo el signo integral

Teorema 1. Sea (X, Σ, μ) un espacio de medida y $f: X \times [a, b] \rightarrow \mathbb{K} = \mathbb{R} \text{ o } \mathbb{C}$ tal que $-\infty \leq a < b \leq \infty$. Definimos $F: [a, b] \rightarrow \mathbb{K}$ mediante

$$\forall t \in [a, b] : F(t) := \int_X f(x, t) d\mu(x).$$

Supongamos que $\exists \frac{\partial f}{\partial t}(x, t)$ para todo $(x, t) \in X \times [a, b]$ y $|\frac{\partial f}{\partial t}(x, t)| \leq g(x)$ para $g \in \mathcal{L}^1(X, \mu)$.

$$\implies F \text{ es diferenciable en } [a, b] \text{ y } \forall t \in [a, b] : F'(t) = \int_X \frac{\partial f}{\partial t}(x, t) d\mu(x).$$

Demostración: Definimos $\forall h \neq 0 : F_h(x, t) = \frac{f(x, t+h) - f(x, t)}{h}$. Tenemos que

$$F'(t) = \lim_{h \rightarrow 0} \frac{F(t+h) - F(t)}{h} = \lim_{h \rightarrow 0} \int_X \frac{f(x, t+h) - f(x, t)}{h} d\mu(x) = \lim_{h \rightarrow 0} \int_X F_h(x, t) d\mu(x).$$

Como $\forall (x, t) \in X \times [a, b] : \exists \frac{\partial f}{\partial t}(x, t)$, se tiene que $\lim_{h \rightarrow 0} F_h(x, t) = \frac{\partial f}{\partial t}(x, t)$. Además, por el teorema del valor medio en la variable t , para cada $(x, t) \in X \times [a, b]$ y $h \neq 0$, existe ξ entre t y $t+h$ tal que

$$|F_h(x, t)| = \left| \frac{f(x, t+h) - f(x, t)}{h} \right| = \left| \frac{\partial f}{\partial t}(x, \xi) \right| \leq g(x).$$

Por tanto, podemos aplicar el teorema de la convergencia dominada:

$$\forall t \in [a, b] : F'(t) = \int_X \lim_{h \rightarrow 0} F_h(x, t) d\mu(x) = \int_X \frac{\partial f}{\partial t}(x, t) d\mu(x). \quad \blacksquare$$

Referenciado en

- cor-convolucion-regularidad

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.