

Teo fn continua soporte compacto denso lp

Teorema 1. $\mathcal{C}_c(\mathbb{R})$ es denso en $\mathcal{L}^p(\mathbb{R}, m)$ para todo $p \in [1, \infty)$.

Demostración: Sea $f \in \mathcal{L}^p(\mathbb{R}, m)$ y $\varepsilon > 0$. Por la `prop-fn-simples-denso-lp/??`, tenemos que $\exists s \in \mathcal{S} \cap \mathcal{L}^p$ tal que $\|f - s\|_p < \varepsilon/2$. Si $s = \sum_{i=1}^n a_i \mathbb{1}_{A_i}$, tomamos $M = \sum_{j=1}^n |a_j|$. Entonces, por el `lem-aprox-indicatriz-continua-norma-lp/Lema 1`, $\forall j \in \mathbb{N}_n : \exists g_j \in \mathcal{C}_c(\mathbb{R}) : \|\mathbb{1}_{A_j} - g_j\|_p < \varepsilon/2M$. Sea $g = \sum_{j=1}^n a_j g_j \in \mathcal{C}_c(\mathbb{R})$, entonces

$$\|s - g\|_p = \left\| \sum_{j=1}^n a_j (\mathbb{1}_{A_j} - g_j) \right\|_p \leq \sum_{j=1}^n |a_j| \cdot \|\mathbb{1}_{A_j} - g_j\|_p \leq \sum_{j=1}^n |a_j| \cdot \frac{\varepsilon}{2M} = \frac{\varepsilon}{2}.$$

Ahora, por Minkowski, $\|f - g\|_p \leq \|f - s\|_p + \|s - g\|_p \leq \varepsilon/2 + \varepsilon/2 = \varepsilon$. ■

Referenciado en

- `teo-fn-suave-soporte-compacto-denso-lp`
- `lem-diferenciacion-lebesgue-l1-imp-casi-toda-parte`
- `lem-convergencia-lp-traslacion`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.