

Teorema de aproximación de la identidad por convolución

Teorema 1. Sea $\phi \in \mathcal{L}^1(\mathbb{R}^n)$ tal que $\phi \geq 0$ y $\int_{\mathbb{R}^n} \phi(x) dx = 1$. Entonces,

- (1) $\forall f \in \mathcal{L}^p(\mathbb{R}^n)$ con $p \in [1, \infty) : \lim_{t \rightarrow 0^+} \|f * \phi_t - f\|_p = 0$.
- (2) $\forall f \in \mathcal{L}^\infty(\mathbb{R}^n)$ uniformemente continua : $\lim_{t \rightarrow 0^+} \|f * \phi_t - f\|_\infty = 0$.
- (3) $\forall f$ continua en $x_0 \in \mathbb{R}^n : \lim_{t \rightarrow 0^+} f * \phi_t(x_0) = f(x_0)$.

Demostración: Como $\int_{\mathbb{R}^n} \phi(x) dx = 1$, por la obs-propiedades-dilatacion-isotropica/??, sabemos que $\forall t > 0 : \int_{\mathbb{R}^n} \phi_t(x) dx = 1$. Entonces, $\forall x \in \mathbb{R}^n :$

$$\begin{aligned} |f * \phi_t(x) - f(x)| &= \left| \int_{\mathbb{R}^n} f(x-y) \phi_t(y) dy - \int_{\mathbb{R}^n} f(x) \phi_t(y) dy \right| \\ &= \left| \int_{\mathbb{R}^n} [f(x-y) - f(x)] \phi_t(y) dy \right| \leq \int_{\mathbb{R}^n} |T_y f(x) - f(x)| \phi_t(y) dy. \end{aligned} \quad (1)$$

(1) Sea q el exponente conjugado de p (i.e. $1/p + 1/q = 1$). Entonces, por Hölder,

$$\begin{aligned} |f * \phi_t(x) - f(x)| &\stackrel{(\text{??})}{\leq} \int_{\mathbb{R}^n} |T_y f(x) - f(x)| \phi_t^{1/p}(y) \phi_t^{1/q}(y) dy \\ &\stackrel{\text{Hölder}}{\leq} \left(\int_{\mathbb{R}^n} |T_y f(x) - f(x)|^p \phi_t(y) dy \right)^{1/p} \cdot \underbrace{\left(\int_{\mathbb{R}^n} \phi_t(y) dy \right)^{1/q}}_1. \end{aligned} \quad (2)$$

$$\begin{aligned} \|f * \phi_t - f\|_p^p &= \int_{\mathbb{R}^n} |f * \phi_t(x) - f(x)|^p dx \stackrel{(\text{??})}{\leq} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |T_y f(x) - f(x)|^p \phi_t(y) dy dx = \\ &\stackrel{\text{Fubini}}{\leq} \int_{\mathbb{R}^n} \phi_t(y) \left(\int_{\mathbb{R}^n} |T_y f(x) - f(x)|^p dx \right) dy = \int_{\mathbb{R}^n} \phi_t(y) \|T_y f - f\|_p^p dy. \end{aligned} \quad (3)$$

Por el lem-convergencia-lp-traslacion/??, dado $\varepsilon > 0$, existe $\eta > 0$ tal que $\forall y \in \mathbb{R}^n : \|y\| < \eta : \|T_y f - f\|_p < \varepsilon$. Entonces, sea $\eta > 0$ tal que $\|y\| < \eta \implies \|T_y f - f\|_p \leq \frac{\varepsilon}{2^{1/p}}$. Por la obs-propiedades-dilatacion-isotropica/??, $\exists \delta > 0 : \forall t \in (0, \delta) :$

$\int_{\|y\| \geq \eta} \phi_t(y) \, dy < \frac{\varepsilon^p}{2^{p+1} \|f\|_p^p}$. Entonces, para $t \in (0, \delta)$,

$$\begin{aligned} \|f * \phi_t - f\|_p^p &\stackrel{(\text{??})}{\leq} \int_{\|y\| < \eta} \|T_y f - f\|_p^p \phi_t(y) \, dy + \int_{\|y\| \geq \eta} \|T_y f - f\|_p^p \phi_t(y) \, dy \\ &\leq \frac{\varepsilon^p}{2} \int_{\|y\| \leq \eta} \phi_t(y) \, dy + 2^p \|f\|_p^p \frac{\varepsilon^p}{2^{p+1} \|f\|_p^p} \leq \frac{\varepsilon^p}{2} + \frac{\varepsilon^p}{2} = \varepsilon^p. \end{aligned}$$

Por tanto, $\lim_{t \rightarrow 0^+} \|f * \phi_t - f\|_p = 0$.

(2) Como f es uniformemente continua, $\forall \varepsilon > 0 : \exists \eta > 0 : \forall x, y \in \mathbb{R}^n : \|y\| < \eta \implies |f(x - y) - f(x)| < \frac{\varepsilon}{2}$. Entonces, por (??), tenemos que

$$\begin{aligned} |f * \phi_t(x) - f(x)| &\stackrel{(\text{??})}{\leq} \int_{\|y\| < \eta} |T_y f(x) - f(x)| \phi_t(y) \, dy + \int_{\|y\| \geq \eta} |T_y f(x) - f(x)| \phi_t(y) \, dy \\ &\leq \frac{\varepsilon}{2} \int_{\mathbb{R}^n} \phi_t(y) \, dy + 2 \|f\|_\infty \int_{\|y\| \geq \eta} \phi_t(y) \, dy \leq \frac{\varepsilon}{2} + 2 \|f\|_\infty \frac{\varepsilon}{4 \|f\|_\infty} = \varepsilon. \end{aligned}$$

Por tanto, $\lim_{t \rightarrow 0^+} \|f * \phi_t - f\|_\infty = 0$.

(3) Como f es continua en x_0 , tenemos que

$$\forall \varepsilon > 0 : \exists \eta > 0 : \forall y \in \mathbb{R}^n : \|y\| < \eta \implies \int_{\|y\| < \eta} \phi_t(y) \, dy \leq \frac{\varepsilon}{4 \|f\|_\infty}.$$

Entonces, el resultado se sigue igual que en (2) reemplazando x por x_0 . ■

Referenciado en

- teo-inversion-transformada-fourier

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.