

Lema de Schwarz

Lema 1 (de Schwarz). Sea $f \in \mathcal{H}(\mathbb{D})$ tal que $f(0) = 0 \wedge \forall z \in \mathbb{D} : |f(z)| \leq 1$, entonces
 (1) $\forall z \in \mathbb{D} : |f(z)| \leq |z| \wedge$ (2) $|f'(0)| \leq 1$.

Además, si se cumple la igualdad en ?? para $z \neq 0$ o en ??, f es una rotación.

Demostración: Sabemos que $f'(0) = \lim_{z \rightarrow 0} \frac{f(z)}{z}$ y definimos la función g dada por

$$\forall z \in \mathbb{D} : g(z) = \begin{cases} f(z)/z & \text{si } z \neq 0 \\ f'(0) & \text{si } z = 0 \end{cases} \implies g \in \mathcal{H}(\mathbb{D}).$$

Sea $r \in (0, 1)$, entonces $g \in \mathcal{H}(D(0, r)) \cap \mathcal{C}(\overline{D(0, r)})$ y, por el principio del módulo máximo, $\forall z \in \overline{D(0, r)} : |g(z)| < \max_{|w|=r} |g(w)|$.

$$\implies \forall z \in \mathbb{D} : |g(z)| \leq \max_{|w|=r} \frac{|f(w)|}{r} \leq \frac{1}{r} \xrightarrow{r \rightarrow 1^-} 1 \implies |g(z)| \leq 1.$$

Por tanto, se verifica que $|g(z)| \leq |z|$ y $|f'(0)| = |g(0)| \leq 1$.

Por otra parte, si $\exists z_0 \in \mathbb{D}^* : |f(z_0)| = |z_0| \vee |f'(0)| = 1 \implies |g(z_0)| = 1 \vee |g(0)| = 1$. Entonces, $|g|$ alcanza su máximo en el $\mathbb{D}$ y, por el teorema del módulo máximo, $g \equiv C \in \mathbb{C}$ constante con $|C| = 1$ (luego $\exists \theta \in \mathbb{R} : C = e^{i\theta}$). Por tanto, $f(z) = Cz = e^{i\theta}z$. ■

Referenciado en

- teo-schwarz-pick
- teo-formula-aut-disco-unidad

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.