

Teo fn suave soporte compacto denso lp

Teorema 1. $\mathcal{C}_c^\infty(\mathbb{R})$ es denso en $\mathcal{L}^p(\mathbb{R}, m)$ para todo $p \in [1, \infty)$.

Demostración: Por las demostraciones del `lem-aprox-indicatriz-continua-norma-lp`/Lema 1 y del `teo-fn-continua-soporte-compacto-denso-lp/??`, basta con hallar $g \in \mathcal{C}_c^\infty(\mathbb{R})$ tal que $\mathbb{1}_{[a,b]}(x) \leq g(x) \leq \mathbb{1}_{[a-\varepsilon_1, b+\varepsilon_2]}(x)$ en función de $a, b, \varepsilon_1, \varepsilon_2$ dados. Para ello, basta con hallar $\varphi \in \mathcal{C}_c^\infty(\mathbb{R})$ tal que

$$\varphi(x) = \begin{cases} 0 & x \leq 0, \\ 1 & x \geq 1, \end{cases} \implies g(x) = \varphi\left(\frac{a-x}{\varepsilon_1}\right) \cdot \varphi\left(\frac{x-b}{\varepsilon_2}\right).$$

Para definir φ , usaremos la función ψ dada por

$$\psi(x) = \begin{cases} e^{-\frac{1}{x}} & x > 0, \\ 0 & x \leq 0. \end{cases} \implies \lim_{x \rightarrow 0^+} \forall n \in \mathbb{N} : \psi^{(n)}(x) = 0.$$

Sea $h(x) = \frac{1}{e} - \psi(x)$, entonces $\varphi(x) := e^e \cdot \psi(h(x))$ cumple lo pedido. ■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.