

Teorema de Hahn-Banach II

Teorema 1 (de Hahn-Banach II). Sea X un espacio vectorial normado, $M \subset X$ un subespacio y $T: M \rightarrow \mathbb{K}$ lineal y continua

$$\implies \exists F \in \mathcal{L}(X, \mathbb{K}) : \begin{cases} \forall x \in M : F(x) = T(x) \\ \|F\| = \|T\|. \end{cases}$$

Demostración: Como T es continua, por el `teo-carac-continuidad-apl-lineal`/Teorema 1, $\forall x \in M : |T(x)| \leq \|T\| \|x\|$. Tomando $p(x) = \|T\| \|x\|$, tenemos que p es una seminorma en X y $\forall x \in M : |T(x)| \leq p(x)$.

Por tanto, por el `teo-hahn-banach-i`/Teorema 1, existe $\exists F: X \rightarrow \mathbb{K}$ lineal tal que $\forall x \in X : |F(x)| \leq p(x)$ y $F|_M = T$. Entonces, $\|F\| = \sup_{\|x\|=1} |F(x)| \leq \sup_{\|x\|=1} p(x) = \|T\|$.

Por otro lado, $\|F\| = \sup_{\|x\|=1} |F(x)| \geq \sup_{\substack{x \in M \\ \|x\|=1}} |F(x)| = \sup_{\substack{x \in M \\ \|x\|=1}} |T(x)| = \|T\|$. ■

Referenciado en

- `teo-subespacio-reflexivo`
- `lem-separacion-punto-conjunto-convexo-abierto`
- `teo-esp-normado-separacion-punto-cero`
- `teo-esp-normado-separacion-punto-subespacio-cerrado`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.