

En un espacio de Hilbert, convergencia débil y en norma implica convergencia fuerte

Proposición 1. *Sea H un espacio de Hilbert real y $(x_n)_{n \in \mathbb{N}} \subset H$, entonces*

$$x_n \rightharpoonup x \wedge \|x_n\| \rightarrow \|x\| \implies x_n \rightarrow x \text{ en } H.$$

Demostración: Basta ver que $\|x_n - x\| \rightarrow 0$:

$$\begin{aligned} \|x_n - x\|^2 &= \langle x_n - x, x_n - x \rangle = \|x_n\|^2 - 2 \langle x_n, x \rangle + \|x\|^2 \\ &= \|x_n\|^2 - 2 (L_x(x_n)) + \|x\|^2. \end{aligned}$$

Al tomar límites cuando $n \rightarrow \infty$ en la expresión anterior, obtenemos que

$$\|x_n - x\|^2 \xrightarrow{n \rightarrow \infty} \|x\|^2 - 2 (L_x(x)) + \|x\|^2 = \|x\|^2 - 2 \|x\|^2 + \|x\|^2 = 0.$$

■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.