

Directamente conforme implica holomorfa

Lema 1. Sea $f: \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$ diferenciable en el sentido real y directamente conforme

$$\implies f \text{ es holomorfa en } \Omega.$$

Demostración: Sea $(x, y) \in \Omega$, como f es directamente conforme, $\exists \lambda > 0, Q \in SO(2)$ tales que $Df(x, y) = \lambda Q$ con $\det Q = 1$. Por tanto, Q actúa como una rotación en $\mathbb{R}^2$, es decir, $\exists \theta \in \mathbb{R}$ tal que

$$Df(x, y) = \begin{pmatrix} \frac{\partial u}{\partial x}(x, y) & \frac{\partial u}{\partial y}(x, y) \\ \frac{\partial v}{\partial x}(x, y) & \frac{\partial v}{\partial y}(x, y) \end{pmatrix} = \lambda Q = \lambda \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

donde $u(x, y) = \Re(f(x + iy))$ y $v(x, y) = \Im(f(x + iy))$.

Observamos que se satisfacen las ecuaciones de Cauchy-Riemann en (x, y) :

$$\frac{\partial u}{\partial x}(x, y) = \lambda \cos \theta = \frac{\partial v}{\partial y}(x, y) \quad \wedge \quad \frac{\partial u}{\partial y}(x, y) = -\lambda \sin \theta = -\frac{\partial v}{\partial x}(x, y).$$

Dado que f es $\mathbb{R}$ -diferenciable, concluimos entonces que f es $\mathbb{C}$ -diferenciable en (x, y) y, como este punto era arbitrario, tenemos que $f \in \mathcal{H}(\Omega)$. ■

Referenciado en

- ??
- ??
- ??
- ??
- ??
- ??

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.