

Corolario del principio del módulo máximo

Corolario 1. Sea $\Omega \subset \mathbb{C}$ un dominio acotado y $f \in \mathcal{H}(\Omega) \cap \mathcal{C}(\overline{\Omega})$

$$\implies \max_{z \in \overline{\Omega}} |f(z)| = \max_{z \in \partial\Omega} |f(z)|.$$

Demostración: Como Ω es acotado, $\overline{\Omega}$ es compacto y, por tanto, como $f \in \mathcal{C}(\overline{\Omega})$, $|f| \in \mathcal{C}(\overline{\Omega})$ alcanza su máximo en $\overline{\Omega}$ (Teorema de Weierstrass). Veamos dos casos:

- I. Si $\exists z_0 \in \Omega$ tal que $|f(z_0)| = \max_{z \in \overline{\Omega}} |f(z)|$, por el principio del módulo máximo, f es constante en Ω y, por tanto, $\max_{z \in \partial\Omega} |f(z)| = |f(z_0)| = \max_{z \in \overline{\Omega}} |f(z)|$.
- II. Si $\forall z \in \Omega : |f(z)| < \max_{z \in \overline{\Omega}} |f(z)|$, entonces, necesariamente, $\exists z_1 \in \partial\Omega$ tal que $|f(z_1)| = \max_{z \in \overline{\Omega}} |f(z)|$. Por tanto, $\max_{z \in \partial\Omega} |f(z)| = |f(z_1)| = \max_{z \in \overline{\Omega}} |f(z)|$. ■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.