

Obs Involucion Disco Unidad

Diametro

Observación 1. Sea $w \in \mathbb{D} \setminus \{0\}$, entonces $\Gamma_w := \{t w / |w| : t \in (-1, 1)\} \subset \mathbb{D}$ es el diámetro del disco unidad en la dirección de w . Al aplicar τ_w , obtenemos que

$$\forall t \in (-1, 1) : \tau_w \left(t \frac{w}{|w|} \right) = \frac{w - t w / |w|}{1 - \bar{w} t w / |w|} = \frac{|w| - t}{1 - t |w|} \cdot \frac{w}{|w|} \in \Gamma_w$$

porque $\frac{|w|-t}{1-t|w|} \in \mathbb{R}$ y $\tau_w(\mathbb{D}) \subset \mathbb{D}$. Por tanto, $\tau_w(\Gamma_w) \subset \Gamma_w$ y, aplicando el apartado ?? del `lem-involucion-disco-unidad/lema 1`, $\Gamma_w = \tau_w \circ \tau_w(\Gamma_w) \subset \tau_w(\Gamma_w)$. En consecuencia, $\tau_w(\Gamma_w) = \Gamma_w$.

Referenciado en

- Prop bola pseudohiperbolica bola euclidea

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.