

# Lema de Schwarz desnormalizado

**Corolario 1.** Sea  $\Omega \subset \mathbb{C}$  un dominio y  $f \in \mathcal{H}(\Omega) : \exists a \in \Omega, M > 0 : f(\Omega) \subset D(f(a), M)$ . Entonces, si definimos  $R := \text{dist}(a, \partial\Omega)$ , se tiene que

$$(1) \quad \forall z \in D(a, R) : |f(z) - f(a)| \leq \frac{M}{R} |z - a|, \quad (2) \quad |f'(a)| \leq \frac{M}{R}.$$

**Demostración:** Consideramos la función  $g : \mathbb{D} \rightarrow \mathbb{C}$  dada por

$$\forall z \in \mathbb{D} : g(z) = \frac{f(a + Rz) - f(a)}{M}.$$

Entonces, si  $z \in \mathbb{D}$ ,  $a + Rz \in D(a, R) \subset \Omega$ , luego  $f(a + Rz) \in D(f(a), M)$ , de donde se deduce que  $g(z) \in \mathbb{D}$ . Por tanto,  $g$  es una función holomorfa en  $\mathbb{D}$  con  $g(0) = 0$  y  $\forall z \in \mathbb{D} : |g(z)| \leq 1$ . Por el lema de Schwarz, tenemos que:

$$(1) \quad \forall z \in \mathbb{D} : |g(z)| \leq |z|, \text{ es decir, } |f(a + Rz) - f(a)| \leq M |z|, \text{ luego}$$

$$\forall z \in D(a, R) : |f(z) - f(a)| \leq \frac{M}{R} |z - a|.$$

$$(2) \quad |g'(0)| \leq 1 \text{ y como } g'(0) = \frac{R}{M} f'(a), \text{ entonces } |f'(a)| \leq \frac{M}{R}. \quad \blacksquare$$

## Referenciado en

- Teo Lem Schwarz Desnormalizado Derivadas
- Teorema de Liouville

### **Temporary page!**

L<sup>A</sup>T<sub>E</sub>X was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L<sup>A</sup>T<sub>E</sub>X now knows how many pages to expect for this document.