

Desigualdad de Young para convoluciones

Proposición 1 (Young). Sean $f \in \mathcal{L}^1(\mathbb{R}^n)$, $g \in \mathcal{L}^p(\mathbb{R}^n)$ con $p \in [1, \infty]$

$$\implies \|f * g\|_p \leq \|f\|_1 \|g\|_p.$$

En particular, $|f * g(x)| < \infty$ c.t.p. $x \in \mathbb{R}^n$.

Demostración: Veamoslo en tres casos mutuamente excluyentes:

Caso I: Si $p = \infty$, por la convolucion/Observación 2, tenemos que $\|f * g\|_\infty \leq \|f\|_1 \|g\|_\infty$.

Caso II: Si $p = 1$, por el Teorema de Fubini, tenemos que

$$\begin{aligned} \|f * g\|_1 &= \int_{\mathbb{R}^n} |f * g(x)| dx = \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} f(x-y) g(y) dy \right| dx \\ &\leq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f(x-y)| |g(y)| dy dx \stackrel{\text{Fubini}}{=} \int_{\mathbb{R}^n} |g(y)| \left(\int_{\mathbb{R}^n} |f(x-y)| dx \right) dy = \\ &\leq \int_{\mathbb{R}^n} |g(y)| \|f\|_1 dy = \|f\|_1 \|g\|_1. \end{aligned}$$

Caso III: Si $p \in (1, \infty)$, tenemos que

$$\|f * g\|_p^p = \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} f(x-y) g(y) dy \right|^p dx \leq \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} |f(x-y)| |g(y)| dy \right)^p dx. \quad (1)$$

Por otro lado, tomando $q \in (1, \infty)$ como el exponente conjugado de p , tenemos que

$$\int_{\mathbb{R}^n} |f(x-y)| |g(y)| dy = \int_{\mathbb{R}^n} |f(x-y)|^{1/q} |f(x-y)|^{1/p} |g(y)| dy.$$

Entonces, por la desigualdad de Hölder para $|f(x-\cdot)|^{1/q}$ y $|f(x-\cdot)|^{1/p} |g|$, tenemos

$$\begin{aligned} \int_{\mathbb{R}^n} |f(x-y)| |g(y)| dy &\leq \left(\int_{\mathbb{R}^n} |f(x-y)| dy \right)^{\frac{1}{q}} \left(\int_{\mathbb{R}^n} |f(x-y)| |g(y)|^p dy \right)^{\frac{1}{p}} = \\ &\leq \|f\|_1^{1/q} \left(\int_{\mathbb{R}^n} |f(x-y)| |g(y)|^p dy \right)^{\frac{1}{p}}. \end{aligned}$$

Sustituyendo con esta expresión en (??), tenemos que

$$\begin{aligned}
\|f * g\|_p^p &\leq \int_{\mathbb{R}^n} \left(\|f\|_1^{1/q} \left(\int_{\mathbb{R}^n} |f(x-y)| |g(y)|^p dy \right)^{\frac{1}{p}} \right)^p dx = \\
&\leq \|f\|_1^{\frac{p}{q}} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f(x-y)| |g(y)|^p dy dx \stackrel{\text{Fubini}}{=} \|f\|_1^{\frac{p}{q}} \int_{\mathbb{R}^n} |g(y)|^p \left(\int_{\mathbb{R}^n} |f(x-y)| dx \right) dy = \\
&= \|f\|_1^{\frac{p}{q}} \|f\|_1 \int_{\mathbb{R}^n} |g(y)|^p dy = \|f\|_1^p \|g\|_p^p.
\end{aligned}$$

Por tanto, elevando a $1/p$, llegamos a $\|f * g\|_p \leq \|f\|_1 \|g\|_p$. ■

Referenciado en

- `lem-transformada-fourier-convolucion`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.