

Desigualdad de Minkowski

Proposición 1 (Desigualdad de Minkowski). Sea $1 \leq p \leq \infty$ y $f, g \in \mathcal{L}^p(\mu)$

$$\implies \|f + g\|_p \leq \|f\|_p + \|g\|_p.$$

Demostración: Dividimos la demostración en tres casos excluyentes:

I. Si $p = 1$, entonces podemos aplicar directamente la desigualdad triangular de $|\cdot|$:

$$\begin{aligned} \|f + g\|_1 &= \int_{\Omega} |f(x) + g(x)| \, d\mu(x) \stackrel{\Delta}{\leq} \int_{\Omega} [|f| + |g|] \, d\mu(x) \\ &\stackrel{\text{lineal}}{=} \int_{\Omega} |f| \, d\mu(x) + \int_{\Omega} |g| \, d\mu(x) = \|f\|_1 + \|g\|_1. \end{aligned}$$

II. Si $1 < p < \infty$, entonces

$$\begin{aligned} |f(x) + g(x)|^p &= |f(x) + g(x)| |f(x) + g(x)|^{p-1} \\ &\leq |f(x)| |f(x) + g(x)|^{p-1} + |g(x)| |f(x) + g(x)|^{p-1}. \end{aligned} \tag{1}$$

Integrando el primer sumando del lado derecho, obtenemos

$$\begin{aligned} \int_{\Omega} |f(x)| |f(x) + g(x)|^{p-1} \, d\mu(x) &\stackrel{\text{Hölder}}{\leq} \|f\|_p \| |f + g|^{p-1} \|_q = \\ &\leq \|f\|_p \left[\int_{\Omega} |f + g|^{(p-1)q} \, d\mu \right]^{1/q} = \\ &\leq \|f\|_p \left[\int_{\Omega} |f + g|^p \, d\mu \right]^{\frac{1}{p} \cdot \frac{p}{q}} = \|f\|_p \left(\|f + g\|_p \right)^{p/q}. \end{aligned} \tag{2}$$

Análogamente, integrando el segundo sumando del lado derecho de (??) se obtiene

$$\int_{\Omega} |g(x)| |f(x) + g(x)|^{p-1} \, d\mu(x) \leq \|g\|_p \left(\|f + g\|_p \right)^{p/q}. \tag{3}$$

Entonces, aplicando las cotas de (??) y (??) en la integral de (??), se tiene que

$$\begin{aligned}
\|f + g\|_p^p &\stackrel{(\text{??})}{=} \int_{\Omega} |f(x)| |f(x) + g(x)|^{p-1} d\mu(x) + \int_{\Omega} |g(x)| |f(x) + g(x)|^{p-1} d\mu(x) \\
&\stackrel{(\text{??}) \text{ y } (\text{??})}{\downarrow} \leq \|f\|_p \|f + g\|_p^{p/q} + \|g\|_p \|f + g\|_p^{p/q} = \|f + g\|_p^{p/q} [\|f\|_p + \|g\|_p] . \\
&\iff \|f + g\|_p^{p-p/q} = \|f + g\|_p \leq \|f\|_p + \|g\|_p .
\end{aligned}$$

III. Si $p = \infty$, entonces tomamos

$$A_1 \subset \Omega : \|f\|_{\infty} = \sup_{x \in A_1^c} |f(x)| \quad \text{y} \quad A_2 \subset \Omega : \|g\|_{\infty} = \sup_{x \in A_2^c} |g(x)| .$$

Definimos $A := A_1 \cup A_2$, entonces

$$\begin{aligned}
\|f\|_{\infty} + \|g\|_{\infty} &= \sup_{x \in A_1^c} |f(x)| + \sup_{x \in A_2^c} |g(x)| \geq \sup_{x \in A^c} |f(x)| + \sup_{x \in A^c} |g(x)| \\
&\geq \sup_{x \in A^c} [|f(x)| + |g(x)|] \geq \sup_{x \in A^c} |f(x) + g(x)| \\
&\quad \uparrow \Delta \\
&\geq \inf_{\substack{B \in \Sigma \\ \mu(B)=0}} \sup_{x \in B^c} |f(x) + g(x)| = \|f + g\|_{\infty} .
\end{aligned}$$

Como se ha demostrado en los tres casos, se tiene que $\|f + g\|_p \leq \|f\|_p + \|g\|_p$. ■

Referenciado en

- teo-fn-continua-soporte-compacto-denso-lp
- lem-esp-lp-normado
- teo-esp-lp-banach
- lem-esp-lp-vectorial
- prop-convergencia-lp-imp-subsucesion-ctp
- lem-convergencia-lp-traslacion
- ley-debil-grandes-numeros

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.