

Anillo polinomios

Proposición 1 (Anillo de polinomios). *Sea A un anillo (conmutativo con unidad)*

$$\implies A[x] := \{a_n x^n + \cdots + a_1 x + a_0 : a_i \in A \wedge n \in \mathbb{N}\}$$

es un anillo (conmutativo con unidad) con las siguientes operaciones:

- *Suma:* $(a_n x^n + \cdots + a_0) + (b_m x^m + \cdots + b_0) = (a_n + b_n) x^n + \cdots + (a_0 + b_0).$
- *Producto:* $(a_n x^n + \cdots + a_0) \cdot (b_m x^m + \cdots + b_0) = \sum_{k=0}^{n+m} \left(\sum_{i=0}^k a_i b_{k-i} \right) x^k.$

Lo llamamos el anillo de polinomios en la variable x con coeficientes en A .

Referenciado en

- alg-cerrado
- ralgebra-finitamente-generada
- ejer-variedad-algebraica-ideal-radical
- variedad-algebraica-afin
- prop-variedad-algebraica-afin-ideal
- teo-cuerpo-imp-polinomios-dip
- prop-grado-polinomio
- lem-anillo-polinomios-variables-cuerpo-ideal-maximal-extension-algebraica-grado-fin
- con-ceros-polinomios-esp-afin
- teo-base-hilbert
- teo-ceros-hilbert
- prop-di-imp-anillo-polinomios-di
- elemento-entero-sobre-anillo

- teo-ideal-maximal-anillo-polinomios-cuerpo-alg-cerrado
- funcion-regular-variedad-algebraica-afin
- ideal-anulacion
- lem-con-ceros-ideal-generado
- elemento-algebraico-sobre-anillo
- teo-con-ceros-polinomios-esp-afin-ideal-propio-cuerpo-alg-cerrado-no-vacio
- obs-con-ceros-ideal-generado-contenido-trivial
- polinomio-monico-variable
- independencia-algebraica
- cor-base-hilbert-n-var
- grado-polinomio
- anillo-coordenadas-variedad-algebraica-afin

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.