

Propiedades de la convolución

Lema 1. Sean $f, g, h: \mathbb{R}^n \rightarrow \mathbb{K} = \mathbb{R} \text{ o } \mathbb{C}$ dos funciones medibles, entonces

- (1) *Conmutatividad:* $f * g = g * f$ c.t.p.
- (2) *Asociatividad:* $f * (g * h) = (f * g) * h$ c.t.p.
- (3) $T_z(f * g) = (T_z f) * g = f * (T_z g)$ c.t.p.
- (4) $\text{supp}(f * g) \subset \text{supp}(f) + \text{supp}(g)$.

Demostración:

$$(1) \quad f * g(x) = \int_{\mathbb{R}^n} f(x - y) g(y) dy \stackrel{z = x - y}{=} \int_{\mathbb{R}^n} f(z) g(x - z) dz = g * f(x) \text{ c.t.p. } x \in \mathbb{R}^n.$$

(2) Calculamos y aplicamos el Teorema de Fubini:

$$\begin{aligned} (f * g) * h(x) &= \int_{\mathbb{R}^n} (f * g)(x - y) h(y) dy = \int_{\mathbb{R}^n} g * f(x - y) h(y) dy \\ &= \int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} g(x - y - z) f(z) dz \right) h(y) dy \\ &\stackrel{\text{Fubini}}{=} \int_{\mathbb{R}^n} f(z) \left(\int_{\mathbb{R}^n} g(x - y - z) h(y) dy \right) dz = \int_{\mathbb{R}^n} g * h(x - z) f(z) dz \\ &= f * (g * h)(x) \end{aligned}$$

(3) Se demuestra la primera igualdad, la segunda es análoga. Tenemos que

$$\begin{aligned} T_z(f * g)(x) &= f * g(x - z) = \int_{\mathbb{R}^n} f(x - z - y) g(y) dy \\ &= \int_{\mathbb{R}^n} T_z f(x - y) g(y) dy = (T_z f) * g(x). \end{aligned}$$

(4) Basta probarlo para el soporte sin cerrar. Es decir, si $s(f) := \{x \in \mathbb{R}^n : f(x) \neq 0\}$ basta ver que $s(f * g) \subset s(f) + s(g)$. Sea $x \in \mathbb{R}^n \setminus (s(f) + s(g))$, entonces $\forall y \in s(g) : f(x - y) = 0$ porque $f(x - y) \neq 0 \implies x - y \in s(f) \implies x = (x - y) + y \in s(f) + s(g)$.

Como $f(x - y) = 0$, $f * g(x) = \int_{\mathbb{R}^n} f(x - y) g(y) dy = 0$, luego $x \in \mathbb{R}^n \setminus s(f * g)$. ■

Referenciado en

- cor-convolucion-regularidad

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.