

Teorema de diferenciación de Lebesgue en $\mathcal{L}^1$ para casi todo punto

Lema 1. Sea $f \in \mathcal{L}^1(\mathbb{R})$ y $\lambda > 0$. Entonces, el conjunto

$$A_\lambda := \left\{ x \in \mathbb{R} : \limsup_{r \rightarrow 0} \left| \frac{1}{2r} \int_{x-r}^{x+r} f(y) dy - f(x) \right| > \lambda \right\} \text{ tiene medida de Lebesgue cero.}$$

Demostración: Sea $\varepsilon > 0$. Como $\mathcal{C}_c(\mathbb{R})$ es denso en $L^1(\mathbb{R})$ (teo-fn-continua-soporte-compacto-dens) $\exists g \in \mathcal{C}_c(\mathbb{R}) : \|f - g\|_1 \leq \varepsilon\lambda/8$. Como g es continua, por el teo-diferenciacion-lebesgue-continuas/?? sabemos que

$$\forall x \in \mathbb{R} : \lim_{r \rightarrow 0^+} \frac{1}{2r} \int_{x-r}^{x+r} g(y) dy = g(x).$$

Se tiene así que $\left| \frac{1}{2r} \int_{x-r}^{x+r} f(y) dy - f(x) \right| =$

$$\begin{aligned} &= \left| \frac{1}{2r} \int_{x-r}^{x+r} f(y) dy - \frac{1}{2r} \int_{x-r}^{x+r} g(y) dy + \frac{1}{2r} \int_{x-r}^{x+r} g(y) dy - g(x) + g(x) - f(x) \right| \\ &\leq \frac{1}{2r} \int_{x-r}^{x+r} |f(y) - g(y)| dy + \left| \frac{1}{2r} \int_{x-r}^{x+r} g(y) dy - g(x) \right| + |g(x) - f(x)|. \end{aligned}$$

Como $\lim_{r \rightarrow 0^+} \frac{1}{2r} \int_{x-r}^{x+r} g(y) dy = g(x)$, tomando el $\limsup$ cuando $r \rightarrow 0^+$, se tiene que

$$\begin{aligned} \limsup_{r \rightarrow 0} \left| \frac{1}{2r} \int_{x-r}^{x+r} f(y) dy - f(x) \right| &\leq \limsup_{r \rightarrow 0} \frac{1}{2r} \int_{x-r}^{x+r} |f(y) - g(y)| dy + 0 + |g(x) - f(x)| \\ &\leq M(f - g)(x) + |g(x) - f(x)|. \end{aligned} \tag{1}$$

Definimos ahora $B_\lambda := \{x \in \mathbb{R} : M(f - g)(x) > \frac{\lambda}{2}\}$ y $C_\lambda := \{x \in \mathbb{R} : |g(x) - f(x)| > \frac{\lambda}{2}\}$.

Entonces, $A_\lambda \subset B_\lambda \cup C_\lambda$ porque $x \in (B_\lambda \cup C_\lambda)^c = B_\lambda^c \cap C_\lambda^c \implies x \in A_\lambda^c$ ya que

$$\begin{cases} x \in B_\lambda^c \implies M(f - g)(x) \leq \lambda/2 \\ x \in C_\lambda^c \implies |g(x) - f(x)| \leq \lambda/2 \end{cases} \implies \limsup_{r \rightarrow 0^+} \left| \frac{1}{2r} \int_{x-r}^{x+r} f(y) dy - f(x) \right| \leq \lambda$$

por la desigualdad (??).

Por tanto, por subaditividad de la medida de Lebesgue, $m(A_\lambda) \leq m(B_\lambda) + m(C_\lambda)$. Ahora

bien, por la desigualdad de Hardy-Littlewood, $m(B_\lambda) \leq \frac{3}{\lambda/2} \|f - g\|_1$. Entonces,

$$\begin{aligned} m(A_\lambda) &\leq \frac{3}{\lambda/2} \|f - g\|_1 + \int_{C_\lambda} 1 \, dy \\ &\leq \frac{6}{\lambda} \|f - g\|_1 + \frac{2}{\lambda} \int_{C_\lambda} |g(x) - f(x)| \, dx \leq \frac{8}{\lambda} \|f - g\|_1 \leq \varepsilon. \end{aligned}$$

Como $\varepsilon > 0$ era arbitrario, concluimos que $m(A_\lambda) = 0$. ■

Referenciado en

- teo-diferenciacion-lebesgue-l1

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.