

(Anti)holomorfía en términos de las derivadas de Wirtinger

Proposición 1. Sea $f \in \mathcal{C}^1(\Omega)$ con $\Omega \subset \mathbb{C}$ abierto, entonces

$$(1) \ f \text{ es holomorfa en } \Omega \iff \forall z \in \Omega : \frac{\partial f}{\partial \bar{z}}(z) = 0.$$

$$(2) \ f \text{ es antiholomorfa en } \Omega \iff \forall z \in \Omega : \frac{\partial f}{\partial z}(z) = 0.$$

Demostración: En ambos casos, procedemos por definición de las derivadas de Wirtinger:

(1) Escribimos $z = x + iy$ con $x, y \in \mathbb{R}$ y $f = u + iv$ con $u = \Re(f), v = \Im(f)$. Entonces,

$$\begin{aligned} \forall z \in \Omega : \frac{\partial f}{\partial \bar{z}}(z) = 0 &\iff \frac{1}{2} \left(\frac{\partial f}{\partial x}(z) + i \frac{\partial f}{\partial y}(z) \right) = 0 \iff \frac{\partial f}{\partial x}(z) + i \frac{\partial f}{\partial y}(z) = 0 \\ &\iff \frac{\partial u}{\partial x}(z) + i \frac{\partial v}{\partial x}(z) + i \frac{\partial u}{\partial y}(z) - \frac{\partial v}{\partial y}(z) = 0. \end{aligned}$$

Tomando ahora partes real e imaginaria tenemos que esto es equivalente a

$$\frac{\partial u}{\partial x}(z) - \frac{\partial v}{\partial y}(z) = 0 \quad \wedge \quad \frac{\partial v}{\partial x}(z) + \frac{\partial u}{\partial y}(z) = 0,$$

que son las Ecuaciones de Cauchy-Riemann, luego esto es equivalente a que f sea holomorfa en Ω . Es decir, f es holomorfa en $\Omega \iff \forall z \in \Omega : \frac{\partial f}{\partial \bar{z}}(z) = 0$.

(2) De forma análoga, tenemos que

$$\begin{aligned} \forall z \in \Omega : \frac{\partial f}{\partial z}(z) = 0 &\iff \frac{\partial u}{\partial x}(z) + i \frac{\partial v}{\partial x}(z) - i \frac{\partial u}{\partial y}(z) + \frac{\partial v}{\partial y}(z) = 0 \\ &\iff \frac{\partial u}{\partial x}(z) + \frac{\partial v}{\partial y}(z) = 0 \quad \wedge \quad \frac{\partial v}{\partial x}(z) - \frac{\partial u}{\partial y}(z) = 0. \end{aligned}$$

Por otro lado, $\bar{f} = u - iv$ y, por tanto, $\frac{\partial \bar{f}}{\partial x} = \frac{\partial u}{\partial x} - i \frac{\partial v}{\partial x}$ y $\frac{\partial \bar{f}}{\partial y} = \frac{\partial u}{\partial y} - i \frac{\partial v}{\partial y}$. Por tanto, por el Teorema de Cauchy-Riemann, $\bar{f}$ es holomorfa en Ω

$$\iff \frac{\partial u}{\partial x}(z) + \frac{\partial v}{\partial y}(z) = 0 \quad \wedge \quad -\frac{\partial v}{\partial x}(z) + \frac{\partial u}{\partial y}(z) = 0,$$

que son precisamente las ecuaciones anteriores. Por tanto, f es antiholomorfa en $\Omega \iff \forall z \in \Omega : \frac{\partial f}{\partial z}(z) = 0$. ■

Referenciado en

- `lem-laplaciano-cambio-variable`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.