

Convergencia puntual dominada implica convergencia en $\mathcal{L}^p$

Proposición 1. Sean $(f_n)_{n \in \mathbb{N}}$ funciones medibles $f_n: X \rightarrow \mathbb{K}$ en (X, Σ, μ) dominadas por $g \in \mathcal{L}^p(X)$ tales que $f_n(x) \xrightarrow[n \rightarrow \infty]{} f(x)$ c.t.p. $x \in X$

$$\implies f_n \xrightarrow[n \rightarrow \infty]{\mathcal{L}^p} f.$$

Demostración: Tenemos que $\forall n \in \mathbb{N} : |f_n(x)| \leq g(x)$ c.t.p. $x \in X$. Por tanto,

$$\forall n \in \mathbb{N} : |f_n(x) - f(x)|^p \leq (|f_n(x)| + |f(x)|)^p \leq (2g(x))^p \quad \text{c.t.p. } x \in X.$$

Luego, por el Teorema de la Convergencia Dominada, tenemos que

$$\lim_{n \rightarrow \infty} \|f_n - f\|_p^p = \lim_{n \rightarrow \infty} \int_X |f_n(x) - f(x)|^p d\mu(x) \stackrel{\text{TCD}}{\downarrow} \int_X \lim_{n \rightarrow \infty} |f_n(x) - f(x)|^p d\mu(x).$$

Como $\lim_n |f_n(x) - f(x)| = 0$ c.t.p. $x \in X$, se tiene que $\lim_n \|f_n - f\|_p^p = 0$. ■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.