

Teo cauchy goursat convexo

Teorema 1 (de Cauchy-Goursat para convexos).

Sea $\Omega \subset \mathbb{C}$ un conjunto abierto y convexo y $f \in \mathcal{H}(\Omega)$, entonces

- (1) *Existencia de primitiva: $\exists F \in \mathcal{H}(\Omega)$ tal que $F' = f$ en Ω .*
- (2) *Teorema de Cauchy-Goursat: $\forall \gamma$ camino cerrado en $\Omega : \int_{\gamma} f(z) dz = 0$.*

Observación 1. Para la existencia de primitiva podemos rebajar la hipótesis sobre f y que se siga garantizando la existencia de dicha primitiva: f continua en Ω y $\int_{\partial R} f = 0$ para todo rectángulo $R \subset \Omega \implies \exists F \in \mathcal{H}(\Omega)$ tal que $F' = f$ en Ω .

Referenciado en

- teo-formula-integral-cauchy-disco
- teo-morera-rectangulo

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.