

Lem aprox indicatriz compacto abierto continua

Lema 1. Sea $V \subset \mathbb{R}$ un abierto y $K \subset \mathbb{R}$ un compacto tales que $K \subset V$

$$\implies \exists g \in \mathcal{C}_c(\mathbb{R}) : \forall x \in \mathbb{R} : \mathbb{1}_K(x) \leq g(x) \leq \mathbb{1}_V(x).$$

Demostración: Como V es abierto, $\exists \{I_j\}_{j \in \mathbb{N}}$ intervalos abiertos tales que $V = \bigsqcup_{j \in \mathbb{N}} I_j$. Como K es compacto y $K \subset V$ (luego $\{I_j\}_{j \in \mathbb{N}}$ es un cubrimiento por abiertos de K), $\exists \{I_{j_\ell}\}_{\ell=1}^m$ tal que $K \subset \bigsqcup_{\ell=1}^m I_{j_\ell}$.

Definimos $\forall \ell \in \mathbb{N}_m : K_\ell := K \cap \overline{I_{j_\ell}}$, entonces K_ℓ es un intervalo compacto y $K_\ell \subset I_{j_\ell}$. Por el `lem-aprox-indicatriz-continua/??`, tenemos que

$$\forall \ell \in \mathbb{N}_m : \exists g_\ell \in \mathcal{C}_c(\mathbb{R}) : \forall x \in \mathbb{R} : \mathbb{1}_{K_\ell}(x) \leq g_\ell(x) \leq \mathbb{1}_{I_{j_\ell}}(x).$$

Ahora definimos $g = \sum_{\ell=1}^m g_\ell \in \mathcal{C}_c(\mathbb{R})$, entonces

$$\forall x \in \mathbb{R} : \mathbb{1}_K(x) \leq \sum_{\ell=1}^m \mathbb{1}_{K_\ell}(x) \leq \sum_{\ell=1}^m g_\ell(x) = g(x) \leq \sum_{\ell=1}^m \mathbb{1}_{I_{j_\ell}}(x) \leq \mathbb{1}_V(x). \quad \blacksquare$$

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.