

Regla de la cadena para derivadas de Wirtinger

Proposición 1 (Regla de la cadena). Sean $f \in \mathcal{C}^1(\Omega_1)$ y $g \in \mathcal{C}^1(\Omega_2)$ con $f(\Omega_1) \subset \Omega_2$, entonces se satisfacen

$$(1) \quad \frac{\partial}{\partial z}(g \circ f)(z) = \frac{\partial g}{\partial z}(f(z)) \frac{\partial f}{\partial z}(z) + \frac{\partial g}{\partial \bar{z}}(f(z)) \frac{\partial \bar{f}}{\partial z}(z).$$

$$(2) \quad \frac{\partial}{\partial \bar{z}}(g \circ f)(z) = \frac{\partial g}{\partial z}(f(z)) \frac{\partial f}{\partial \bar{z}}(z) + \frac{\partial g}{\partial \bar{z}}(f(z)) \frac{\partial \bar{f}}{\partial \bar{z}}(z).$$

Demostración: Veremos únicamente la primera identidad, la segunda se demuestra de forma análoga. Escribimos $f(z) = u(x, y) + i v(x, y)$. Entonces, por definición de derivada de Wirtinger y la regla de la cadena para derivadas parciales, se tiene que

$$2 \frac{\partial}{\partial z}(g \circ f)(z) = \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (g \circ f)(z) = \frac{\partial g}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial g}{\partial v} \frac{\partial v}{\partial x} - i \frac{\partial g}{\partial u} \frac{\partial u}{\partial y} - i \frac{\partial g}{\partial v} \frac{\partial v}{\partial y} = (*).$$

Aplicando ahora el `lem-derivadas-parciales-wirtinger`/Lema 1, tenemos que

$$\begin{aligned} (*) &= \left(\frac{\partial g}{\partial z} + \frac{\partial g}{\partial \bar{z}} \right) \frac{\partial u}{\partial x} + i \left(\frac{\partial g}{\partial z} - \frac{\partial g}{\partial \bar{z}} \right) \frac{\partial v}{\partial x} - i \left(\frac{\partial g}{\partial z} + \frac{\partial g}{\partial \bar{z}} \right) \frac{\partial u}{\partial y} + \left(\frac{\partial g}{\partial z} - \frac{\partial g}{\partial \bar{z}} \right) \frac{\partial v}{\partial y} \\ &= \frac{\partial g}{\partial z} \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} - i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \right) + \frac{\partial g}{\partial \bar{z}} \left(\frac{\partial u}{\partial x} - i \frac{\partial v}{\partial x} - i \frac{\partial u}{\partial y} - \frac{\partial v}{\partial y} \right) \\ &= \frac{\partial g}{\partial z} \left(2 \frac{\partial u}{\partial z} + 2i \frac{\partial v}{\partial z} \right) + \frac{\partial g}{\partial \bar{z}} \left(2 \frac{\partial u}{\partial \bar{z}} + 2i \frac{\partial v}{\partial \bar{z}} \right) = 2 \frac{\partial g}{\partial z}(f(z)) \frac{\partial f}{\partial z}(z) + 2 \frac{\partial g}{\partial \bar{z}}(f(z)) \frac{\partial \bar{f}}{\partial \bar{z}}(z). \end{aligned}$$

Luego el resultado se sigue dividiendo entre 2. ■

Referenciado en

- `lem-laplaciano-cambio-variable`

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.