

Lema de aproximación por funciones simples

Lema 1. Sea (X, Σ) un espacio medible y $f: X \rightarrow [0, \infty]$ medible

$$\implies \exists (s_n)_{n \in \mathbb{N}} \subset \mathcal{S} : \forall x \in X : 0 \leq s_n(x) \leq s_{n+1}(x) \leq f(x) \wedge \lim_{n \rightarrow \infty} s_n(x) = f(x).$$

Demostración: Descomponiendo “en filas” la imagen de f , tenemos que la siguiente sucesión que cumple las condiciones del enunciado:

$$\forall n \in \mathbb{N} : s_n := \sum_{m=0}^{n2^n} \frac{m}{2^n} \mathbb{1}_{E_{n,m}} \text{ con } E_{n,m} := \left\{ x \in X : \min(f(x), n) \in \left[\frac{m}{2^n}, \frac{m+1}{2^n} \right) \right\}$$

Se tiene que $\forall n \in \mathbb{N} : s_n$ es simple por ser suma finita de funciones características escaladas. Veamos que es creciente. Sea $x \in X$ y $y = f(x)$.

- $y = \infty \implies s_n = n \implies \forall n \in \mathbb{N} : s_n \leq s_{n+1}$.
- $y < \infty \implies s_n(x) = \frac{\lfloor n2^n \rfloor}{2^n}$ y queremos ver que $\frac{\lfloor (n+1)2^{n+1} \rfloor}{2^{n+1}} \geq \frac{\lfloor n2^n \rfloor}{2^n}$.

$$\frac{\lfloor (n+1)2^{n+1} \rfloor + 1}{2^{n+1}} > y \geq \frac{\lfloor n2^n \rfloor}{2^n} \implies \frac{\lfloor (n+1)2^{n+1} \rfloor + 1}{2^{n+1}} > \frac{\lfloor n2^n \rfloor}{2^n}$$

$$\implies \lfloor (n+1)2^{n+1} \rfloor + 1 > 2 \lfloor n2^n \rfloor \implies \lfloor (n+1)2^{n+1} \rfloor \geq 2 \lfloor n2^n \rfloor$$

$$\implies \frac{\lfloor (n+1)2^{n+1} \rfloor}{2^{n+1}} \geq \frac{2 \lfloor n2^n \rfloor}{2^{n+1}} = \frac{\lfloor n2^n \rfloor}{2^n} \implies s_{n+1}(x) \geq s_n(x) \implies s_{n+1} \geq s_n.$$

■

Referenciado en

- prop-fn-simples-denso-lp

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.