

Teo base schauder imp separable

Teorema 1. *Sea X un espacio de Banach, entonces*

$$\exists (e_n)_{n \in \mathbb{N}} \subset X \text{ base de Schauder} \implies X \text{ es separable.}$$

Demostración: Suponemos que $\forall n \in \mathbb{N} : \|e_n\| = 1$ (si no, consideramos $\frac{e_n}{\|e_n\|}$). Sea $x \in X$, entonces $\exists! (a_n)_{n \in \mathbb{N}} \subset \mathbb{R} : \|x - \sum_{i=0}^n a_i e_i\| \xrightarrow{n \rightarrow \infty} 0$. Por tanto, dado $\varepsilon > 0$, $\exists N \in \mathbb{N} : \forall n \geq N : \|x - \sum_{i=0}^n a_i e_i\| < \frac{\varepsilon}{2}$.

Sea $(q_n)_{n \in \mathbb{N}} \subset \mathbb{K}$ tal que $\forall n \in \mathbb{N} : \Re(q_n), \Im(q_n) \in \mathbb{Q}$ y $|a_n - q_n| < \frac{\varepsilon}{2N}$. Entonces,

$$\begin{aligned} \left\| x - \sum_{i=0}^N q_i e_i \right\| &\leq \left\| x - \sum_{i=0}^N a_i e_i \right\| + \left\| \sum_{i=0}^N (a_i - q_i) e_i \right\| \\ &< \frac{\varepsilon}{2} + \sum_{i=0}^N |a_i - q_i| \|e_i\| < \frac{\varepsilon}{2} + N \cdot \frac{\varepsilon}{2N} = \varepsilon. \end{aligned}$$

Por tanto, el conjunto $\left\{ \sum_{i=0}^n q_i e_i : n \in \mathbb{N} \wedge \Re(q_i), \Im(q_i) \in \mathbb{Q} \right\}$ es denso en X y, como también es numerable, X es separable. ■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.