

Lem convergencia uniforme compactos imp convergencia sucesion

Lema 1. Sea $\Omega \subset \mathbb{C}$ un dominio, $(f_n)_{n \in \mathbb{N}} \subset \mathcal{C}(\Omega)$ y $f: \Omega \rightarrow \mathbb{C}$ tales que $f_n \xrightarrow[n \rightarrow \infty]{\Omega} f$. Sea $(z_n)_{n \in \mathbb{N}} \subset \Omega$ tal que $z_n \xrightarrow[n \rightarrow \infty]{} a \in \Omega$, entonces $f_n(z_n) \xrightarrow[n \rightarrow \infty]{} f(a)$.

Demostración: Como $z_n \xrightarrow[n \rightarrow \infty]{} a$, entonces $\exists N_0 \in \mathbb{N} : \forall n \geq N_0 : z_n \in D(a, 1)$. Por tanto, $K := \{z_n : n \geq N_0\} \cup \{a\}$ es un subconjunto compacto de Ω . Entonces, $f_n \xrightarrow[n \rightarrow \infty]{K} f$, es decir, dado $\varepsilon > 0$, $\exists N \in \mathbb{N} : \forall n \geq N : \forall z \in K : |f_n(z) - f(z)| < \varepsilon/2$.

En particular, $\forall n \geq \max\{N, N_0\} : |f_n(z_n) - f(z_n)| < \varepsilon/2$ y, como por el `lem-convergencia-uniforme-compactos` $f \in \mathcal{C}(\Omega)$, $|f(z_n) - f(a)| < \varepsilon/2$. Por tanto, para todo $n \geq \max\{N, N_0\}$,

$$|f_n(z_n) - f(a)| \leq |f_n(z_n) - f(z_n)| + |f(z_n) - f(a)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \quad \blacksquare$$

Referenciado en

- Lem Convergencia Uniforme Compactos Subsucesiones

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.