

Lem Precompacidad Imp Acotacion Uniforme Compactos

Lema 1. Sea $\Omega \subset \mathbb{C}$ un dominio y $\mathcal{F} \subset \mathcal{C}(\Omega)$ precompacto, entonces $\mathcal{F}$ está uniformemente acotado sobre compactos.

Demostración: Supongamos por contradicción que

$$\exists K \subseteq \Omega \text{ compacto} : \forall n \in \mathbb{N} : \exists f_n \in \mathcal{F}, z_n \in K : |f_n(z_n)| > n.$$

Como K es compacto, $\exists z \in K, (z_{n_k})_{k \in \mathbb{N}} \subset (z_n)_{n \in \mathbb{N}}$ tal que $z_{n_k} \xrightarrow[k \rightarrow \infty]{} z$. Por el lem-carac-precompacidad- como $\mathcal{F}$ es precompacto, $\exists (f_{n_{k_j}})_{j \in \mathbb{N}} \subset (f_{n_k})_{k \in \mathbb{N}} : \exists f \in \mathcal{C}(\Omega) : f_{n_{k_j}} \xrightarrow[j \rightarrow \infty]{\Omega} f$. Entonces, tenemos que $f_{n_{k_j}}(z) \xrightarrow[j \rightarrow \infty]{} f(z)$, pero

$$\left| f_{n_{k_j}}(z) \right| \geq \left| f_{n_{k_j}}(z_{n_{k_j}}) \right| - \left| f_{n_{k_j}}(z) - f_{n_{k_j}}(z_{n_{k_j}}) \right| > n_{k_j} - \left| f_{n_{k_j}}(z) - f_{n_{k_j}}(z_{n_{k_j}}) \right| \xrightarrow[j \rightarrow \infty]{} \infty,$$

lo que es una contradicción. ■

Temporary page!

L^AT_EX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because L^AT_EX now knows how many pages to expect for this document.