Numerical Analysis Homework 2

131044009 - Hasan MEN

7 March 2017

Apply Gaussian and Jaboci's method (50 points) 1

Program Usage:

\$./solver -i system.txt -m GESP/JCB

Use Gaussian Elimination with Scaled Partial Pivoting

Question 6.2-1c 1.1.1

$$2x_1 - 3x_2 + 2x_3 = 5 (1)$$

$$-4x_1 + 2x_2 - 6x_3 = 14 (2)$$

$$2x_1 + 2x_2 + 4x_3 = 8 (3)$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(1)} = \begin{bmatrix} 2 & -3 & 2 \mid 5 \\ -4 & 2 & -6 \mid 14 \\ 2 & 2 & 4 \mid 8 \end{bmatrix}$$

Find Max values for each lines:

$$S_i = \max_{i=1}^n ||a_{ij}|| \tag{4}$$

$$S_1 = \max\{\|2\|, \|-3\|, \|2\|\} = 3 \tag{5}$$

$$S_2 = \max\{\|-4\|, \|2\|, \|-6\|\} = 6 \tag{6}$$

$$S_3 = \max\{\|2\|, \|2\|, \|4\|\} = 3 \tag{7}$$

Now, find scaled partial pivots of first column to check if need interchanges

$$\max_{i=1}^{n} \{ \frac{\|a_{ij}\|}{S_i} \} \tag{8}$$

$$\frac{\|a_{11}\|}{S_1} = 0.666667\tag{9}$$

$$\frac{\|a_{21}\|}{S_2} = 0.666667 \tag{10}$$

$$\frac{\|a_{11}\|}{S_1} = 0.666667$$

$$\frac{\|a_{21}\|}{S_2} = 0.666667$$

$$\frac{\|a_{31}\|}{S_3} = 0.500000$$
(11)

Maximum pivot index=1. There is no row interchange Now calculate m_{21} and m_{31} to reduce rows

$$m_{21} = \frac{a_{21}}{a_{11}} = -2 (12)$$

$$m_{31} = \frac{a_{31}}{a_{11}} = 1 \tag{13}$$

Apply this operations on matrix:

$$E_2 - m_{21} * E_1 - > E_2 \tag{14}$$

$$E_3 - m_{31} * E_1 - > E_3 \tag{15}$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(2)} = \begin{bmatrix} 2 & -3 & 2 & 5 \\ 0 & -4 & -2 & 24 \\ 0 & 5 & 2 & 3 \end{bmatrix}$$

Now, find scaled partial pivots of second column to check if need interchanges

$$\frac{\|a_{22}\|}{S_2} = 0.666667$$

$$\frac{\|a_{32}\|}{S_3} = 1.250000$$
(16)

$$\frac{\|a_{32}\|}{S_2} = 1.250000\tag{17}$$

Maximum pivot index=3 and pivot=1.25. Change row 3 with 2.

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(2)} = \begin{bmatrix} 2 & -3 & 2 \mid 5 \\ 0 & 5 & 2 \mid 3 \\ 0 & -4 & -2 \mid 24 \end{bmatrix}$$

Now calculate m_{32} to reduce row

$$m_{32} = \frac{a_{32}}{a_{22}} = -2 \tag{18}$$

Apply this operations on matrix:

$$E_3 - m_{32} * E_2 - > E_3 \tag{19}$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(3)} = \begin{bmatrix} 2 & -3 & 2 & 5 \\ 0 & 5 & 2 & 3 \\ 0 & 0 & -0.4 & 26.4 \end{bmatrix}$$

Solve matrix with backward substitution.

Results:

$$\begin{bmatrix} X \end{bmatrix} = \begin{bmatrix} 109 & 27 & -66 \end{bmatrix}^t$$

1.1.2 Question 7.3.1-a

$$3x_1 - 1x_2 + 1x_3 = 1 (20)$$

$$3x_1 + 6x_2 + 2x_3 = 0 (21)$$

$$3x_1 + 3x_2 + 7x_3 = 4 (22)$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(1)} = \begin{bmatrix} 3 & -1 & 1 \mid 1 \\ 3 & 6 & 2 \mid 0 \\ 3 & 3 & 7 \mid 4 \end{bmatrix}$$

Find Max values for each lines:

$$S_i = \max_{i=1}^n ||a_{ij}|| \tag{23}$$

$$S_1 = \max\{\|3\|, \|-1\|, \|1\|\} = 3 \tag{24}$$

$$S_2 = \max\{\|3\|, \|6\|, \|2\|\} = 6 \tag{25}$$

$$S_3 = \max\{\|3\|, \|3\|, \|7\|\} = 7 \tag{26}$$

Now, find scaled partial pivots of first column to check if need interchanges

$$\max_{i=1}^{n} \{ \frac{\|a_{ij}\|}{S_i} \} \tag{27}$$

$$\frac{\|a_{11}\|}{S_1} = 1.000000 \tag{28}$$

$$\frac{\|a_{21}\|}{S_2} = 0.500000 \tag{29}$$

$$\frac{\|a_{31}\|}{S_3} = 0.428571\tag{30}$$

Maximum pivot index=1 and pivot=1.0. There is no row interchange Now calculate m_{21} and m_{31} to reduce rows

$$m_{21} = \frac{a_{21}}{a_{11}} = 1 \tag{31}$$

$$m_{21} = \frac{a_{21}}{a_{11}} = 1$$
 (31)
 $m_{31} = \frac{a_{31}}{a_{11}} = 1$ (32)

Apply this operations on matrix:

$$E_2 - m_{21} * E_1 - > E_2 \tag{33}$$

$$E_3 - m_{31} * E_1 - > E_3 \tag{34}$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(2)} = \begin{bmatrix} 3 & -1 & 1 \mid 1 \\ 0 & 7 & 1 \mid -1 \\ 0 & 4 & 6 \mid 3 \end{bmatrix}$$

Now, find scaled partial pivots of second column to check if need interchanges

$$\frac{\|a_{22}\|}{S_2} = 1.166667\tag{35}$$

$$\frac{\|a_{22}\|}{S_2} = 1.166667$$

$$\frac{\|a_{32}\|}{S_3} = 0.571429$$
(35)

Maximum pivot index=2 and pivot=1.166667. No Row Change Now calculate m_{32} to reduce row

$$m_{32} = \frac{a_{32}}{a_{22}} = 0.571429 \tag{37}$$

Apply this operations on matrix:

$$E_3 - m_{32} * E_2 - > E_3 \tag{38}$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(3)} = \begin{bmatrix} 3 & -1 & 1 & 1 \\ 0 & 7 & 1 & -1 \\ 0 & 0 & 5.429 & 3.571 \end{bmatrix}$$

Solve matrix with backward substitution.

Results:

$$[X] = \begin{bmatrix} 0..351 & -0.2368 & 0.6579 \end{bmatrix}^t$$

Question 7.3.1-b 1.1.3

$$10x_1 - x_2 + 0 = 9 (39)$$

$$-x_1 + 10x_2 - 2x_3 = 7 (40)$$

$$0 - 2x_2 + 10x_3 = 6 (41)$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(1)} = \begin{bmatrix} 10 & -1 & 0 \mid 9 \\ -1 & 10 & -2 \mid 7 \\ 0 & -2 & 10 \mid 6 \end{bmatrix}$$

Find Max values for each lines:

$$S_i = \max_{i=1}^n ||a_{ij}|| \tag{42}$$

$$S_1 = \max\{\|10\|, \|-1\|, \|0\|\} = 10 \tag{43}$$

$$S_2 = \max\{\|-1\|, \|10\|, \|-2\|\} = 10 \tag{44}$$

$$S_3 = \max\{\|0\|, \|-2\|, \|10\|\} = 10 \tag{45}$$

Now, find scaled partial pivots of first column to check if need interchanges

$$max_{i=1}^{n} \left\{ \frac{\|a_{ij}\|}{S_i} \right\} \tag{46}$$

$$\frac{\|a_{11}\|}{S_1} = 1.000000$$

$$\frac{\|a_{21}\|}{S_2} = 0.100000$$
(47)

$$\frac{\|a_{21}\|}{S_2} = 0.100000 \tag{48}$$

$$\frac{\|a_{31}\|}{S_3} = 0.000000 \tag{49}$$

Maximum pivot index=1 and pivot=1.0. There is no row interchange. Now calculate m_{21} and m_{31} to reduce rows

$$m_{21} = \frac{a_{21}}{a_{11}} = -0.1 \tag{50}$$

$$m_{31} = \frac{a_{31}}{a_{11}} = 0 (51)$$

Apply this operations on matrix:

$$E_2 - m_{21} * E_1 - > E_2 \tag{52}$$

$$E_3 - m_{31} * E_1 - > E_3 \tag{53}$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(2)} = \begin{bmatrix} 10 & -1 & 0 & 9 \\ 0 & 9.9 & -2 & 7.9 \\ 0 & -2 & 10 & 6 \end{bmatrix}$$

Now, find scaled partial pivots of second column to check if need interchanges

$$\frac{\|a_{22}\|}{S_0} = 0.990000 \tag{54}$$

$$\frac{\|a_{22}\|}{S_2} = 0.990000$$

$$\frac{\|a_{32}\|}{S_3} = 0.200000$$
(54)

Maximum pivot index=2 and pivot=0.990000. No Row Change Now calculate m_{32} to reduce row

$$m_{32} = \frac{a_{32}}{a_{22}} = 0.202020 \tag{56}$$

Apply this operations on matrix:

$$E_3 - m_{32} * E_2 - > E_3 \tag{57}$$

$$\begin{bmatrix} A \mid b \end{bmatrix}^{(3)} = \begin{bmatrix} 10 & -1 & 0 & 9 \\ 0 & 9.9 & -2 & 7.9 \\ 0 & 0 & 9.596 & 7.596 \end{bmatrix}$$

Solve matrix with backward substitution.

Results:

$$[X] = \begin{bmatrix} 0.9958 & 0.9579 & 0.7916 \end{bmatrix}^t$$