

Functions

Section 2.3


Section Summary

- Definition of a Function.
 - Domain, Codomain
 - Image, Preimage
- Injection, Surjection, Bijection
- Inverse Function
- Function Composition
- Graphing Functions
- Floor, Ceiling, Factorial

Functions


Definition: Let A and B be nonempty sets. A *function* f from A to B , denoted $f: A \rightarrow B$ is an assignment of each element of A to exactly one element of B . We write $f(a) = b$ if b is the unique element of B assigned by the function f to the element a of A .

- Functions are sometimes called *mappings* or *transformations*.



Given a function $f: A \rightarrow B$

- We say f maps A to B or f is a *mapping* from A to B .
- A is called the *domain* of f .
- B is called the *codomain* of f .
- If $f(a) = b$,
 - then b is called the *image* of a under f .
 - a is called the *preimage* of b .
- The range of f is the set of all images of points in A under f . We denote it by $f(A)$.
- Two functions are *equal* when they have the same domain, the same codomain and map each element of the domain to the same element of the codomain.



Representing Functions

- Functions may be specified in different ways:
 - An explicit statement of the assignment.
Students and grades example.
 - A formula.
$$f(x) = x + 1$$
 - A computer program.
 - A C++ program that when given an integer n , produces the n th Fibonacci Number (covered in the next section and also in Chapter 5).

Questions

$f(a) = ?$ **z**

The image of d is ? **z**


The domain of f is ? **A**

The codomain of f is ? **B**

The preimage of y is ? **b**

$f(A) = ?$ **{y,z}**

The preimage(s) of z is (are) ? **{a,c,d}**




Question on Functions and Sets

- If $f : A \rightarrow B$ and S is a subset of A , then

$$f(S) = \{f(s) | s \in S\}$$



$f\{a,b,c\}$ is ? $\{y,z\}$

$f\{c,d\}$ is ? $\{z\}$




Injectons

Definition: A function f is said to be *one-to-one*, or *injective*, if and only if $f(a) = f(b)$ implies that $a = b$ for all a and b in the domain of f . A function is said to be an *injection* if it is one-to-one.




Surjections

Definition: A function f from A to B is called *onto* or *surjective*, if and only if for every element $b \in B$ there is an element $a \in A$ with $f(a) = b$. A function f is called a *surjection* if it is onto.



Bijections

Definition: A function f is a *one-to-one correspondence*, or a *bijection*, if it is both one-to-one and onto (surjective and injective).



Showing that f is one-to-one or onto

Suppose that $f : A \rightarrow B$.

To show that f is injective Show that if $f(x) = f(y)$ for arbitrary $x, y \in A$ with $x \neq y$, then $x = y$.

To show that f is not injective Find particular elements $x, y \in A$ such that $x \neq y$ and $f(x) = f(y)$.

To show that f is surjective Consider an arbitrary element $y \in B$ and find an element $x \in A$ such that $f(x) = y$.

To show that f is not surjective Find a particular $y \in B$ such that $f(x) \neq y$ for all $x \in A$.

Showing that f is one-to-one or onto

Example 1: Let f be the function from $\{a, b, c, d\}$ to $\{1, 2, 3\}$ defined by $f(a) = 3$, $f(b) = 2$, $f(c) = 1$, and $f(d) = 3$. Is f an onto function?

Solution: Yes, f is onto since all three elements of the codomain are images of elements in the domain. If the codomain were changed to $\{1, 2, 3, 4\}$, f would not be onto.


Example 2: Is the function $f(x) = x^2$ from the set of integers onto?

Solution: No, f is not onto because there is no integer x with $x^2 = -1$, for example.



Inverse Functions

Definition: Let f be a bijection from A to B . Then the *inverse* of f , denoted f^{-1} , is the function from B to A defined as $f^{-1}(y) = x$ iff $f(x) = y$

No inverse exists unless f is a bijection. Why?



Inverse Functions



Questions

Example 1: Let f be the function from $\{a, b, c\}$ to $\{1, 2, 3\}$ such that $f(a) = 2$, $f(b) = 3$, and $f(c) = 1$. Is f invertible and if so what is its inverse?

Solution: The function f is invertible because it is a one-to-one correspondence. The inverse function f^{-1} reverses the correspondence given by f , so $f^{-1}(1) = c$, $f^{-1}(2) = a$, and $f^{-1}(3) = b$.

Questions

Example 2: Let $f: \mathbf{Z} \rightarrow \mathbf{Z}$ be such that $f(x) = x + 1$. Is f invertible, and if so, what is its inverse?

Solution: The function f is invertible because it is a one-to-one correspondence. The inverse function f^{-1} reverses the correspondence so $f^{-1}(y) = y - 1$.

Questions


Example 3: Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be such that $f(x) = x^2$. Is f invertible, and if so, what is its inverse?

Solution: The function f is not invertible because it is not one-to-one .



Composition

- **Definition:** Let $f: B \rightarrow C$, $g: A \rightarrow B$. The composition of f with g , denoted $f \circ g$ is the function from A to C defined by

$$(f \circ g)(a) = f(g(a))$$



Composition



Composition

Example 1: If $f(x) = x^2$ and $g(x) = 2x + 1$,
then

$$f(g(x)) = (2x + 1)^2$$

and

$$g(f(x)) = 2x^2 + 1$$

Composition Questions

Example 2: Let g be the function from the set $\{a,b,c\}$ to itself such that $g(a) = b$, $g(b) = c$, and $g(c) = a$. Let f be the function from the set $\{a,b,c\}$ to the set $\{1,2,3\}$ such that $f(a) = 3$, $f(b) = 2$, and $f(c) = 1$.

What is the composition of f and g , and what is the composition of g and f .

Solution: The composition $f \circ g$ is defined by

$$f \circ g (a) = f(g(a)) = f(b) = 2.$$

$$f \circ g (b) = f(g(b)) = f(c) = 1.$$

$$f \circ g (c) = f(g(c)) = f(a) = 3.$$

Note that $g \circ f$ is not defined, because the range of f is not a subset of the domain of g .

Composition Questions

Example 2: Let f and g be functions from the set of integers to the set of integers defined by $f(x) = 2x + 3$ and $g(x) = 3x + 2$.

What is the composition of f and g , and also the composition of g and f ?


Solution:

$$f \circ g(x) = f(g(x)) = f(3x + 2) = 2(3x + 2) + 3 = 6x + 7$$


$$g \circ f(x) = g(f(x)) = g(2x + 3) = 3(2x + 3) + 2 = 6x + 11$$

Graphs of Functions

- Let f be a function from the set A to the set B . The *graph* of the function f is the set of ordered pairs $\{(a, b) \mid a \in A \text{ and } f(a) = b\}$.



Graph of $f(n) = 2n + 1$
from \mathbb{Z} to \mathbb{Z}



Graph of $f(x) = x^2$
from \mathbb{Z} to \mathbb{Z}

Some Important Functions

- The *floor* function, denoted

$$f(x) = \lfloor x \rfloor$$

is the largest integer less than or equal to x .

- The *ceiling* function, denoted



$$f(x) = \lceil x \rceil$$

is the smallest integer greater than or equal to x

Example: $\lceil 3.5 \rceil = 4$ $\lfloor 3.5 \rfloor = 3$

$$\lceil -1.5 \rceil = -1 \quad \lfloor -1.5 \rfloor = -2$$

Floor and Ceiling Functions



Graph of (a) Floor and (b) Ceiling Functions

Floor and Ceiling Functions

TABLE 1 Useful Properties of the Floor and Ceiling Functions.

(n is an integer, x is a real number)

(1a) $\lfloor x \rfloor = n$ if and only if $n \leq x < n + 1$

(1b) $\lceil x \rceil = n$ if and only if $n - 1 < x \leq n$

(1c) $\lfloor x \rfloor = n$ if and only if $x - 1 < n \leq x$

(1d) $\lceil x \rceil = n$ if and only if $x \leq n < x + 1$

(2) $x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$

(3a) $\lfloor -x \rfloor = -\lceil x \rceil$

(3b) $\lceil -x \rceil = -\lfloor x \rfloor$

(4a) $\lfloor x + n \rfloor = \lfloor x \rfloor + n$

(4b) $\lceil x + n \rceil = \lceil x \rceil + n$

Proving Properties of Functions

Example: Prove that if x is a real number, then

$$\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$$

Solution: Let $x = n + \varepsilon$, where n is an integer and $0 \leq \varepsilon < 1$.

Case 1: $0 \leq \varepsilon < 1/2$

- $2x = 2n + 2\varepsilon$ and $\lfloor 2x \rfloor = 2n$, since $0 \leq 2\varepsilon < 1$.
- $\lfloor x + 1/2 \rfloor = n$, since $x + 1/2 = n + (1/2 + \varepsilon)$ and $0 \leq 1/2 + \varepsilon < 1$.
- Hence, $\lfloor 2x \rfloor = 2n$ and $\lfloor x \rfloor + \lfloor x + 1/2 \rfloor = n + n = 2n$.

Case 2: $1/2 \leq \varepsilon < 1$

- $2x = 2n + 2\varepsilon = (2n + 1) + (2\varepsilon - 1)$ and $\lfloor 2x \rfloor = 2n + 1$, since $0 \leq 2\varepsilon - 1 < 1$.
- $\lfloor x + 1/2 \rfloor = \lfloor n + (1/2 + \varepsilon) \rfloor = \lfloor n + 1 + (\varepsilon - 1/2) \rfloor = n + 1$ since $0 \leq \varepsilon - 1/2 < 1$.
- Hence, $\lfloor 2x \rfloor = 2n + 1$ and $\lfloor x \rfloor + \lfloor x + 1/2 \rfloor = n + (n + 1) = 2n + 1$. ◀

Factorial Function

Definition: $f: \mathbf{N} \rightarrow \mathbf{Z}^+$, denoted by $f(n) = n!$ is the product of the first n positive integers when n is a nonnegative integer.

$$f(n) = 1 \cdot 2 \cdots (n-1) \cdot n, \quad f(0) = 0! = 1$$

Examples:

$$f(1) = 1! = 1$$

$$f(2) = 2! = 1 \cdot 2 = 2$$

$$f(6) = 6! = 1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 = 720$$

$$f(20) = 2,432,902,008,176,640,000.$$

Stirling's Formula:

$$n! \sim \sqrt{2\pi n} (n/e)^n$$

$$f(n) \sim g(n) \doteq \lim_{n \rightarrow \infty} f(n)/g(n) = 1$$