


Hyperbolicity, injective hulls, and Helly graphs


Presenter: Heather M. Guarnera
Algorithmic Lab

Applications of Hyperbolicity

- Many real world networks have **small hyperbolicity** (biological, social, collaboration, communication, etc.)
- Smaller value means the network
 - is **metrically closer to a tree**
 - has **negative curvature**



A graph is **δ -hyperbolic** provided for any vertices x, y, u, v in it, the two larger of the three sums $d(u, v) + d(x, y)$, $d(u, x) + d(v, y)$, and $d(u, y) + d(v, x)$ differ by at most 2δ .





Small hyperbolicity implies that the shortest path between two points curves inward towards the core of the network.

How hyperbolicity relates to Injective Hulls

Every graph G can be isometrically embedded into the smallest **Helly** graph $H(G)$ [1,2]

- $H(G)$ is called the **injective hull** of G
- $H(G)$ **preserves hyperbolicity**
- If G is δ -hyperbolic, any vertex in $H(G)$ is **within 2δ** to a vertex in G [3]

- A set S of sets S_i has the **Helly property** if for every subset T of S the following hold: if the elements of T pairwise intersect, then the intersection of all elements of T is also non-empty.
- A graph is called **Helly** if its family of disks satisfies the Helly property.



We want to understand:

- **(Q1) what governs hyperbolicity in Helly graphs** in order to understand what governs hyperbolicity in regular graphs, and
- **(Q2) how does the injective hull grow** for various graph classes?

[1] J. Isbell. *Six theorems about injective metric spaces*, Comment. Math. Helv (1964).

[2] A. Dress. *Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups*, Adv. in Math (1984).


[3] U. Lang, *Injective hulls of certain discrete metric spaces and groups*, J. Topol. Anal. (2013)

(Q1) Interval thinness governs hyperbolicity in Helly graphs

- An interval $I(x,y)$ is the set of all vertices from shortest (x,y) -paths.


- A slice of an interval at distance k is defined as:

$$S_k(x, y) = \{z \in I(x, y) : d(z, x) = k\}$$



- An interval is τ -thin if for any natural number k and any two u, v vertices of $S_k(x, y)$ are at most τ apart.
- A graph is τ -thin if all of its intervals are at most τ -thin.

For general graphs $\tau(G) \leq 2hb(G)$,
but $\tau(G)$ and $hb(G)$ can be far apart.




example: odd cycle with $4k+1$ vertices

Our Contribution

Theorem [4]:

For Helly graphs, $\tau(G) \leq 2hb(G) \leq \tau(G) + 1$.




Example when $2hb(G) = \tau(G) + 1$

(Q1) Special subgraphs of a chess grid govern hyperbolicity in Helly graphs

Our Contribution


Theorem [4]: We show that for Helly graphs and any integer k ,

- $hb(G) \leq k$ if and only if G has no isometric H_2^k




$hb(G)$ is an integer

- $hb(G) \leq k + 1/2$ if and only if G has neither isometric H_1^{k+1} nor H_3^k




$hb(G)$ is a half-integer

Example: forbidden isometric subgraphs for 1-hyperbolic Helly graphs.




Example: forbidden isometric subgraphs for 1/2-hyperbolic Helly graphs.



(Q2) How big is $H(G)$ with respect to G ?


Every graph G can be isometrically embedded into the smallest **Helly** graph $H(G)$, called the **injective hull** of G .



Our Contribution


Theorem [5]: There are some graph classes for which $H(G)$... contains at most $2n$ vertices.

- Interval graphs
- Helly graphs
- Distance hereditary graphs




... can contain at least 2^n vertices.

- Chordal bipartite graphs
- Chordal graphs
- Cocomparability graphs



Open Questions and Future Work

- What other graph classes can be Hellified efficiently?
- What other graph classes require exponentially many Helly vertices?
- What kind of problems can use $H(G)$ to solve problems efficiently on G ?
 - Diameter
 - Radius
 - Center



idea: use center of $H(G)$
to find center of G