

Lattice Quantum Chromo Dynamics

Giovanni Pedervia, Mathias Vege

May 13, 2018

University of Oslo

Introduction

- Why Lattice QCD? Or QCD in general?
- Low-energy regime is non-perturbative.
- To understand nuclear forces, one needs a first principles approach.
- Constants achieved from LQCD can be used in Chiral effective field theory, which can then be used in CC methods.
- Can be used to investigate Dark Matter phenomena.

The full QCD Lagrangian,

$$\mathcal{L} = -\frac{1}{4} G_{\mu\nu}^a G_a^{\mu\nu} + \bar{\psi}_i (i\gamma^\mu D_{\mu ij} - m\delta_{ij}) \psi_j \quad (1)$$

a is color indices, i, j is flavour indices.

Challenges with LQCD

- Create a program that...
 - generates field configurations.
 - flows generated field configurations.
 - Samples the field for P, E, Q, Q_{t_e} .
- Take care of technicalities such as parallelization, $SU(3)$ exponentiation, ect.
- Perform analysis with bootstrap, jackknife, autocorrelation.
- Side projects: animating the field energy density and topological charge.

- Discretizing the Gluon field in spacetime by introducing *link variables*, $U_\mu(x)$.

Path integrals

For field theories we discretize the field into points in space time.
For a scalar field, this becomes

$$\int \mathcal{D}\phi \rightarrow \int \prod_{x_i \in \text{lattice}} d\phi(x_i)$$

From this we can extract observables $\Gamma[\phi]$ dependent on this field as

$$\langle \Gamma[\phi] \rangle = \frac{1}{Z} \int \mathcal{D}\phi \Gamma[\phi] e^{-S[\phi]}$$

where the normalization Z is given again by

$$Z = \int \mathcal{D}\phi e^{-S[\phi]}$$

Setup

- How do you share a face in 4D?
- 3 different approaches
 - **Face-sharing.** Edge cases, not possible in the Metropolis algorithm
 - **Single link-sharing.** Relatively simple to implement, but slow if network is slow.
 - **Shifts.** Large non-blocking data packets at a time.

Data is generated with the Metropolis algorithm, and following setup,

- Thermalize lattice. Roughly 10000 updates.
 1. Perform N_{corr} updates on the *whole* Lattice.
 2. For each N_{corr} update, we also perform N_{up} updates on every *single* link.
 3. Measure the change in ΔS , the action at each update. Accept config if $\exp(-\Delta S) > r$, where r is a number drawn from a uniform distribution.
 4. Flow the lattice and calculate observable at each step or save it to file.

A way of renormalizing the field.

Table 1: Every set is generated with 600 correlation updates between each configuration, and 30 updates per link, except for 6.45, which has 1600 correlation updates. The the side of the cube is then around $L = 2.2$ fermi.


β	Size	N_{cfs}	Size
6.0	24×48	1000	356 GB
6.1	28×56	500	330 GB
6.2	32×64	500	562 GB
6.45	48×96	250	1.4 TB

Observables

Plaquette

The *plaquette* is the simplest possible gauge invariant object on the lattice. It is a square made of 4 link variables:

$$\begin{aligned} P_{\mu\nu} &= U_\mu(n) U_\nu(n + \hat{\mu}) U_{-\mu}(n + \hat{\mu} + \hat{\nu}) U_{-\nu}(n + \hat{\nu}) \\ &= U_\mu(n) U_\nu(n + \hat{\mu}) U_\mu(n + \hat{\nu})^\dagger U_\nu(n)^\dagger \end{aligned}$$



Can be thought of as the curl of the gluon field. Analogous to

$$u \times v = \varepsilon^{ijk} u_j v_k$$

$$Q = - \sum_x \frac{1}{64 \cdot 32 \pi^2} \epsilon_{\mu\nu\rho\sigma} \text{Tr}\{ G_{\mu\nu}^{clov} G_{\rho\sigma}^{clov} \} \quad (2)$$

$$\chi^{1/4} = \frac{\hbar c}{a V^{1/4}} \langle Q^2 \rangle^{1/4} \quad (3)$$

N_f - checking the flavor

Using Witten-Veneziano formula to look at how different χ behave in the continuum limit and which that gives the best error estimate.

$$\frac{F_\pi^2 m_{\eta'}^2}{2N_f} = \chi$$

where F_π is the decay rate constant of pion given as $F_\pi = 130.0 \pm 5.0 \text{ MeV}$. The mass of the meson η' is given as $m_{\eta'} = 957.78 \pm 0.06 \text{ MeV}$. N_f is the number of flavors.

- G.P. Lepage, *Lattice QCD for Novices*, arXiv:hep-lat/0506036 (2005)
- C. Gattringer & C.P. Lang, *Quantum Chromodynamics on the Lattice*, Springer (2010)