

Solving $SU(3)$ Yang-Mills theory on the lattice: a calculation of selected gauge observables with gradient flow

Hans Mathias Mamen Vege
April 26, 2019

Supervisor: *Andrea Shindler*
Co-supervisor: *Morten Hjorth-Jensen*
University of Oslo

Introduction

- Quantum Chromodynamics(QCD).

- **QCD.** We will go through and explain what QCD as well as motivate its existence.

- Quantum Chromodynamics(QCD).
- Lattice QCD.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.
- **GLAC.** Will briefly present the code which we developed as well as some benchmarks. We will also present the Metropolis algorithm.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.
- Results.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.
- **GLAC.** Will briefly present the code which we developed as well as some benchmarks. We will also present the Metropolis algorithm.
- **Results.** We will present the results obtained from pure gauge calculations.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.
- Results.

Quantum Chromodynamics(QCD)

- The standard model: Six quarks and eight gluons

- The standard model.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom

- The standard model.
- Asymptotic freedom.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement


- The standard model.
- Asymptotic freedom.
- Confinement.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement
- Highly nonlinear due to gluon self-interactions


- The standard model.
- Asymptotic freedom.
- Confinement.
- Nonlinearity.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement
- Highly nonlinear due to gluon self-interactions

The Standard Model




Consists of the innermost square of the six quarks and the gluons.



- The coupling constant **decreases** as we **increase** the energy
- One of the experimental proofs of QCD along with triple γ decay and muon cross section ration R .

Confinement



5

If we try to pull apart **two mesons**, more and more energy is required until we have enough energy to spontaneously create a **quark-antiquark** pair, forming thus **two new mesons**.


The non-linearity of QCD

The QCD Lagrangian

$$\mathcal{L}_{\text{QCD}} = \sum_{f=1}^{N_f} \bar{\psi}^{(f)} \left(i \not{D} - m^{(f)} \right) \psi^{(f)} - \frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu},$$

with action

$$S = \int d^4x \mathcal{L}_{\text{QCD}}. \quad (1)$$



- Instantons

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.

- Instantons
- Topological charge, Q

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.
- Measuring **topological charge** is a measure of the *Winding number* of the gauge field.

- Instantons
- Topological charge, Q
- Winding number




Figure 1: The figure is taken from Forkel [1, p. 32].

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.
- Measuring **topological charge** is a measure of the *Winding number* of the gauge field.
- An illustration of how one can view the winding number given a function f that parametrizes a path around a circle S^1 . Given that it starts and ends at the same point, we have that the number of times it wraps around the circle gives us the winding number.

- Instantons
- Topological charge, Q
- Winding number




Figure 1: The figure is taken from Forkel [1, p. 32].

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.
- Measuring **topological charge** is a measure of the *Winding number* of the gauge field.
- An illustration of how one can view the winding number given a function f that parametrizes a path around a circle S^1 . Given that it starts and ends at the same point, we have that the number of times it wraps around the circle gives us the winding number.

A formula connecting pure gauge theory and full QCD.

$$m_{\eta'}^2 = \frac{2N_f}{f_\pi^2} \chi_{\text{top}} \quad (2)$$

- Pion decay constant $f_\pi = 0.130(5)/\sqrt{2}$ GeV.
- η' meson mass $m_{\eta'} = 0.95778(6)$ GeV.
- χ_{top} is the *topological susceptibility*.

8

- We use the experimental values for the pion decay constant and the η' mass.
- Allows us to estimate the number of flavors in our theory N_f .
- χ_{top} is the topological susceptibility, calculated from the expectation value of Q .

Lattice Quantum Chromodynamics(LQCD)

1. Divide spacetime into a cube of size $N^3 \times N_T$.


1. Divide spacetime into a cube of size $N^3 \times N_T$.
2. Fermions live on the each *point* in the cube.

1. Divide spacetime into a cube of size $N^3 \times N_T$.
2. Fermions live on the each *point* in the cube.
3. The gauge fields live on the sites *in between* the points, and is called links.

A link

$$U_\mu(n) = \exp [iaA_\mu(n)] ,$$

connects one lattice site to another and is a $SU(3)$ matrix.



where $U_{-\mu}(n) = U_\mu(n - \hat{\mu})^\dagger$.

- Defined from the gauge transporter.
- A link in the positive $\hat{\mu}$ direction is shown in the figure to the left.
- A link in the negative $\hat{\mu}$ direction is shown in the figure to the right.

Links gauge transform as

$$\begin{aligned}U_\mu(n) &\rightarrow U'_\mu(n) = \Omega(n) U_\mu(n) \Omega(n + \hat{\mu})^\dagger, \\U_{-\mu}(n) &\rightarrow U'_{-\mu}(n) = \Omega(n) U_\mu(n - \hat{\mu})^\dagger \Omega(n - \hat{\mu})^\dagger.\end{aligned}$$

Links gauge transform as

$$\begin{aligned}U_\mu(n) &\rightarrow U'_\mu(n) = \Omega(n) U_\mu(n) \Omega(n + \hat{\mu})^\dagger, \\U_{-\mu}(n) &\rightarrow U'_{-\mu}(n) = \Omega(n) U_\mu(n - \hat{\mu})^\dagger \Omega(n - \hat{\mu})^\dagger.\end{aligned}$$

Two main types of gauge invariant objects,

Links gauge transform as

$$\begin{aligned}U_\mu(n) &\rightarrow U'_\mu(n) = \Omega(n) U_\mu(n) \Omega(n + \hat{\mu})^\dagger, \\U_{-\mu}(n) &\rightarrow U'_{-\mu}(n) = \Omega(n) U_\mu(n - \hat{\mu})^\dagger \Omega(n - \hat{\mu})^\dagger.\end{aligned}$$


Two main types of gauge invariant objects,

- Fully connected gauge invariant objects.
- Objects with fermions $\psi, \bar{\psi}$ as end points.

The plaquette

The simplest gauge invariant object,

$$\begin{aligned} P_{\mu\nu}(n) &= U_\mu(n) U_\nu(n + \hat{\mu}) U_{-\mu}(n + \hat{\mu} + \hat{\nu}) U_{-\nu}(n + \hat{\nu}) \\ &= U_\mu(n) U_\nu(n + \hat{\mu}) U_\mu(n + \hat{\nu})^\dagger U_\nu(n)^\dagger, \end{aligned}$$



The Wilson gauge action is given as

$$S_G[U] = \frac{\beta}{3} \sum_{n \in \Lambda} \sum_{\mu < \nu} \text{Re tr} [1 - P_{\mu\nu}(n)], \quad (3)$$

with $\beta = 6/g_S^2$.

- Using the definition of the link we saw earlier, we can reproduce the continuum action up to an discretization error of $\mathcal{O}(a^2)$.

Developing a code for solving $SU(3)$ Yang-Mills theory on the lattice

A lattice configuration consists of $SU(3)$ matrices,

14

- The $SU(3)$ matrices are 3×3 matrices of nine complex numbers or 18 real numbers.
- This leads to an absolute **requirement of efficiency**, both in **calculations** and in **input/output**.
- When returning to what ensembles of configurations we generated this will be evident.

A lattice configuration consists of SU(3) matrices,

$$\underbrace{N^3}_{\text{Spatial}} \times \underbrace{N_T}_{\text{Temporal}} \times \underbrace{4}_{\text{Links}} \times \underbrace{9}_{\text{SU(3) matrix}} \times \underbrace{2}_{\mathbb{C}\text{-numbers}} = 72N^3N_T,$$

14

- The SU(3) matrices are 3×3 matrices of nine complex numbers or 18 real numbers.
- This leads to an absolute **requirement of efficiency**, both in **calculations** and in **input/output**.
- When returning to what ensembles of configurations we generated this will be evident.

A lattice configuration consists of SU(3) matrices,

$$\underbrace{N^3}_{\text{Spatial}} \times \underbrace{N_T}_{\text{Temporal}} \times \underbrace{4}_{\text{Links}} \times \underbrace{9}_{\text{SU(3) matrix}} \times \underbrace{2}_{\text{C-numbers}} = 72N^3N_T,$$

$\rightarrow 8 \times 72N^3N_T$ bytes.

- The SU(3) matrices are 3×3 matrices of nine complex numbers or 18 real numbers.
- This leads to an absolute **requirement of efficiency**, both in **calculations** and in **input/output**.
- When returning to what ensembles of configurations we generated this will be evident.

$$\langle O \rangle = \frac{1}{Z} \int \mathcal{D}U O[\psi, \bar{\psi}, U] e^{-S_G[U] - S_F[\psi, \bar{\psi}, U]}.$$

with

$$Z = \int \mathcal{D}U \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{-S_G[U] - S_F[\psi, \bar{\psi}, U]}.$$

$$\langle O \rangle = \frac{1}{Z} \int \mathcal{D}U O[U] e^{-S_G[U]}.$$


with

$$Z = \int \mathcal{D}U e^{-S_G[U]}.$$

$$\langle O \rangle = \frac{1}{Z} \int \mathcal{D}U O[U] e^{-S_G[U]}.$$

with

$$Z = \int \mathcal{D}U e^{-S_G[U]}.$$



An example of the discretized path integral, going from time t_0 to t_{N_T} , where the end points is taken to be equal, $x_0 = x_{N_T}$. We integrate over all of space at each time t_i finding the most likely position at a given time.

The observable becomes an average over the N_{MC} gauge configurations.

$$\langle O \rangle = \lim_{N_{\text{MC}} \rightarrow \infty} \frac{1}{N_{\text{MC}}} \sum_i^{N_{\text{MC}}} O[U_i]$$

We now need to generate configurations...

- We perform an average of the created configurations.

repeat

Randomly generate a candidate state j with probability $T_{i \rightarrow j}$.

Calculate $A_{i \rightarrow j}$ which saw on previous slide.

Generate random number $u \in [0, 1]$.

if $u \leq A_{i \rightarrow j}$ **then**

Accept new state j .

else if $u > A_{i \rightarrow j}$ **then**

Reject new state j and retain the old state i .

end if

until N_{MC} samples are generated.

- Generated state j is a gauge configuration.
- Algorithm of choice when sampling gauge configurations.
- For generating N_{MC} Monte Carlo samples.

A parameter ϵ_{rnd} controls the spread of the candidate matrices.

1. Initialize lattice with $\text{SU}(3)$ matrices close to unity(*hot start*) or at unity(*cold start*).
2. Thermalize with N_{therm} sweeps.
3. Generate N_{MC} samples,
 - i Perform N_{corr} correlation updates.
 - ii At each update, perform N_{up} single link update for every lattice link.
 - iii Store configuration and/or apply gradient flow and sample observables on it.


19

- We use **periodic boundary conditions** for all calculations.
- N_{MC} is how many configurations we will generate.
- N_{up} is how many single link updates we will perform.
- N_{corr} is how many full sweeps we shall perform in between each sampling. Needed in order to reduce the autocorrelation between the configurations.

Two methods used:

- Single link sharing used in the Metropolis algorithm.
- *shifts* used in in gradient flow and observable sampling


- Tested out **halos**, but turned out to be problematic when generating.
- We parallelized using MPI.



- An illustration of the lattice shift.
- The links U_ν of the lattice are copied over to a temporary lattice shifted in direction $\hat{\mu}$.
- The face that is shifted over to an adjacent sub-lattice is shared through a non-blocking MPI call, while we copy the links to the temporary lattice.
- Allows for a simplified syntax close to that of the equations we are working with.
- Don't have to write out any loops over the lattice positions.

We checked three types of scaling,

- **Strong scaling:** *fixed problem* and a variable N_p cores




- Strong scaling

Scaling

We checked three types of scaling,

- **Strong scaling:** *fixed problem* and a variable N_p cores
- **Weak scaling:** *fixed problem per processor* and a variable N_p cores.




22

- Strong scaling
- Weak scaling

We checked three types of scaling,

- **Strong scaling:** *fixed problem* and a variable N_p cores
- **Weak scaling:** *fixed problem per processor* and a variable N_p cores.
- **Speedup:** defined as $S(p) = \frac{t_{N_p,0}}{t_{N_p}}$.



22

- Strong scaling
- Weak scaling
- The speedup of the configuration generation, flowing, and IO. The speedup is calculated by dividing the run time of each N_p run, with the run time of the run with the least number of processors, $N_p = 8$.

We appear to have a plateau around 512 cores.

We checked three types of scaling,

- **Strong scaling:** *fixed problem* and a variable N_p cores
- **Weak scaling:** *fixed problem per processor* and a variable N_p cores.
- **Speedup:** defined as $S(p) = \frac{t_{N_p,0}}{t_{N_p}}$.


22

- Strong scaling
- Weak scaling
- The speedup of the configuration generation, flowing, and IO. The speedup is calculated by dividing the run time of each N_p run, with the run time of the run with the least number of processors, $N_p = 8$.

We appear to have a plateau around 512 cores.

Optimizing the gauge configuration generation

Generated 200 configurations for a lattice of size $N^3 \times N_T = 16^3 \times 32$ and $\beta = 6.0$.



- We run for different values for N_{up} and N_{corr} to see what gives optimizes **computational cost** and **autocorrelation**.
- The integrated autocorrelation time for topological charge $\langle Q \rangle$ for a lattice of size $N = 16$ and $N_T = 32$ with $\beta = 6.0$ for combinations of $N_{\text{corr}} \in [200, 400, 600]$ and $N_{\text{up}} \in [10, 20, 30]$, plotted against flow time $\sqrt{8}t_f$.

Optimizing the gauge configuration generation

Generated 200 configurations for a lattice of size $N^3 \times N_T = 16^3 \times 32$ and $\beta = 6.0$.

23

- We run for different values for N_{up} and N_{corr} to see what gives optimizes **computational cost** and **autocorrelation**.
- The time taking to generate 200 configurations and flowing them $N_{\text{flow}} = 250$ flow steps for a lattice of size $N = 16$ and $N_T = 32$, with $\beta = 6.0$ for combinations of $N_{\text{corr}} \in [200, 400, 600]$ and $N_{\text{up}} \in [10, 20, 30]$.

Optimizing the gauge configuration generation

Generated 200 configurations for a lattice of size $N^3 \times N_T = 16^3 \times 32$ and $\beta = 6.0$.

23

- We run for different values for N_{up} and N_{corr} to see what gives optimizes **computational cost** and **autocorrelation**.
- The time taking to generate 200 configurations and flowing them $N_{\text{flow}} = 250$ flow steps for a lattice of size $N = 16$ and $N_T = 32$, with $\beta = 6.0$ for combinations of $N_{\text{corr}} \in [200, 400, 600]$ and $N_{\text{up}} \in [10, 20, 30]$.

Gradient flow

The flow of the SU(3) gauge fields are denoted by $B_\mu(x, t_f)$ which are Lie algebra valued gauge fields,

$$\frac{d}{dt_f} B_\mu(x, t_f) = D_\nu G_{\nu\mu}(x, t_f), \quad (4)$$

$$D_\mu = \partial_\mu + [B_\mu(x, t_f), \cdot], \quad (5)$$

$$G_{\mu\nu}(x, t_f) = \partial_\mu B_\nu(x, t_f) - \partial_\nu B_\mu(x, t_f) - i[B_\mu(x, t_f), B_\nu(x, t_f)], \quad (6)$$


with the initial condition of eq. (4) being the fundamental gauge field,

$$B_\mu(x, t_f)|_{t_f=0} = A_\mu(x).$$

- Bad approx.: diffusion equation.
- Topological charge preserved and is more pronounced.
- Renormalizes the topological charge at non-zero flow time.

Results

The clover field strength definition



- We will use the clover field strength definition in gauge observables

$$E = \frac{a^4}{2|\Lambda|} \sum_{n \in \Lambda} \sum_{\mu, \nu} \left(F_{\mu\nu}^{\text{clov}}(n) \right)^2$$

- We can use this definition to set a scale.

$$Q = a^4 \sum_{n \in \Lambda} q(n),$$

with the charge density given by

$$q(n) = \frac{1}{32\pi^2} \epsilon_{\mu\nu\rho\sigma} \text{tr} [F_{\mu\nu}(n) F_{\rho\sigma}(n)].$$

- We will use the clover field strength definition.
- Symmetries will allow us to reduce the effective number of clovers need to calculate from 24 to 6.

Conclusion

Questions?

References

- [1] Hilmar Forkel. A Primer on Instantons in QCD. *arXiv:hep-ph/0009136*, September 2000. URL <http://arxiv.org/abs/hep-ph/0009136>. arXiv: hep-ph/0009136.