

Solving $SU(3)$ Yang-Mills theory on the lattice: a calculation of selected gauge observables with gradient flow

Hans Mathias Mamen Vege
April 25, 2019

Supervisor: *Andrea Shindler*
Co-supervisor: *Morten Hjorth-Jensen*
University of Oslo

Introduction

- Quantum Chromodynamics(QCD).

- **QCD.** We will go through and explain what QCD as well as motivate its existence.

- Quantum Chromodynamics(QCD).
- Lattice QCD.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.
- **GLAC.** Will briefly present the code which we developed as well as some benchmarks. We will also present the Metropolis algorithm.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.
- Results.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.
- **GLAC.** Will briefly present the code which we developed as well as some benchmarks. We will also present the Metropolis algorithm.
- **Results.** We will present the results obtained from pure gauge calculations.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.
- Results.

Quantum Chromodynamics(QCD)

- The standard model: Six quarks and eight gluons

- The standard model.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom

- The standard model.
- Asymptotic freedom.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement


- The standard model.
- Asymptotic freedom.
- Confinement.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement
- Highly nonlinear due to gluon self-interactions


- The standard model.
- Asymptotic freedom.
- Confinement.
- Nonlinearity.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement
- Highly nonlinear due to gluon self-interactions


The Standard Model



Consists of the innermost square of the six quarks and the gluons.



- The coupling constant **decreases** as we **increase** the energy
- One of the experimental proofs of QCD along with triple γ decay and muon cross section ratio R .



If we try to pull apart **two mesons**, more and more energy is required until we have enough energy to spontaneously create a **quark-antiquark** pair, forming thus **two new mesons**.


The non-linearity of QCD

The QCD Lagrangian

$$\mathcal{L}_{\text{QCD}} = \sum_{f=1}^{N_f} \bar{\psi}^{(f)} \left(i \not{D} - m^{(f)} \right) \psi^{(f)} - \frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu},$$

with action

$$S = \int d^4x \mathcal{L}_{\text{QCD}}. \quad (1)$$



- Instantons

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.

- Instantons
- Topological charge, Q

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.
- Measuring **topological charge** is a measure of the *Winding number* of the gauge field.

- Instantons
- Topological charge, Q
- Winding number




Figure 1: The figure is taken from Forkel [1, p. 32].

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.
- Measuring **topological charge** is a measure of the *Winding number* of the gauge field.
- An illustration of how one can view the winding number given a function f that parametrizes a path around a circle S^1 . Given that it starts and ends at the same point, we have that the number of times it wraps around the circle gives us the winding number.

- Instantons
- Topological charge, Q
- Winding number




Figure 1: The figure is taken from Forkel [1, p. 32].

- **Instantons** are local minimums to the Yang-Mills action in Euclidean space.
- Measuring **topological charge** is a measure of the *Winding number* of the gauge field.
- An illustration of how one can view the winding number given a function f that parametrizes a path around a circle S^1 . Given that it starts and ends at the same point, we have that the number of times it wraps around the circle gives us the winding number.

A formula connecting pure gauge theory and full QCD.

$$m_{\eta'}^2 = \frac{2N_f}{f_\pi^2} \chi_{\text{top}} \quad (2)$$

- Pion decay constant $f_\pi = 0.130(5)/\sqrt{2}$ GeV.
- η' meson mass $m_{\eta'} = 0.95778(6)$ GeV.
- χ_{top} is the *topological susceptibility*.

8

- We use the experimental values for the pion decay constant and the η' mass.
- Allows us to estimate the number of flavors in our theory N_f .
- χ_{top} is the topological susceptibility, calculated from the expectation value of Q .

Lattice Quantum Chromodynamics(LQCD)

1. Divide spacetime into a cube of size $N^3 \times N_T$.


1. Divide spacetime into a cube of size $N^3 \times N_T$.
2. Fermions live on the each *point* in the cube.

1. Divide spacetime into a cube of size $N^3 \times N_T$.
2. Fermions live on the each *point* in the cube.
3. The gauge fields live on the sites *in between* the points, and is called links.

A link

$$U_\mu(n) = \exp [iaA_\mu(n)] ,$$

connects one lattice site to another and is a $SU(3)$ matrix.



where $U_{-\mu}(n) = U_\mu(n - \hat{\mu})^\dagger$.

- Defined from the gauge transporter.
- A link in the positive $\hat{\mu}$ direction is shown in the figure to the left.
- A link in the negative $\hat{\mu}$ direction is shown in the figure to the right.

Links gauge transform as

$$\begin{aligned}U_\mu(n) &\rightarrow U'_\mu(n) = \Omega(n) U_\mu(n) \Omega(n + \hat{\mu})^\dagger, \\U_{-\mu}(n) &\rightarrow U'_{-\mu}(n) = \Omega(n) U_\mu(n - \hat{\mu})^\dagger \Omega(n - \hat{\mu})^\dagger.\end{aligned}$$

Links gauge transform as

$$\begin{aligned}U_\mu(n) &\rightarrow U'_\mu(n) = \Omega(n) U_\mu(n) \Omega(n + \hat{\mu})^\dagger, \\U_{-\mu}(n) &\rightarrow U'_{-\mu}(n) = \Omega(n) U_\mu(n - \hat{\mu})^\dagger \Omega(n - \hat{\mu})^\dagger.\end{aligned}$$

Two main types of gauge invariant objects,

Links gauge transform as

$$\begin{aligned}U_\mu(n) &\rightarrow U'_\mu(n) = \Omega(n) U_\mu(n) \Omega(n + \hat{\mu})^\dagger, \\U_{-\mu}(n) &\rightarrow U'_{-\mu}(n) = \Omega(n) U_\mu(n - \hat{\mu})^\dagger \Omega(n - \hat{\mu})^\dagger.\end{aligned}$$


Two main types of gauge invariant objects,

- Fully connected gauge invariant objects.
- Objects with fermions $\psi, \bar{\psi}$ as end points.

The plaquette

The simplest gauge invariant object,

$$\begin{aligned} P_{\mu\nu}(n) &= U_\mu(n) U_\nu(n + \hat{\mu}) U_{-\mu}(n + \hat{\mu} + \hat{\nu}) U_{-\nu}(n + \hat{\nu}) \\ &= U_\mu(n) U_\nu(n + \hat{\mu}) U_\mu(n + \hat{\nu})^\dagger U_\nu(n)^\dagger, \end{aligned}$$



The Wilson gauge action is given as

$$S_G[U] = \frac{\beta}{3} \sum_{n \in \Lambda} \sum_{\mu < \nu} \text{Re tr} [1 - P_{\mu\nu}(n)], \quad (3)$$

with $\beta = 6/g_S$.

- Using the definition of the link we saw earlier, we can reproduce the continuum action up to an discretization error of $\mathcal{O}(a^2)$.

Developing a code for solving $SU(3)$ Yang-Mills theory on the lattice

A lattice configuration consists of $SU(3)$ matrices,

14

- The $SU(3)$ matrices are 3×3 matrices of nine complex numbers or 18 real numbers.
- This leads to an absolute **requirement of efficiency**, both in **calculations** and in **input/output**.
- When returning to what ensembles of configurations we generated this will be evident.

A lattice configuration consists of SU(3) matrices,

$$\underbrace{N^3}_{\text{Spatial}} \times \underbrace{N_T}_{\text{Temporal}} \times \underbrace{4}_{\text{Links}} \times \underbrace{9}_{\text{SU(3) matrix}} \times \underbrace{2}_{\mathbb{C}\text{-numbers}} = 72N^3N_T,$$

14

- The SU(3) matrices are 3×3 matrices of nine complex numbers or 18 real numbers.
- This leads to an absolute **requirement of efficiency**, both in **calculations** and in **input/output**.
- When returning to what ensembles of configurations we generated this will be evident.

A lattice configuration consists of SU(3) matrices,

$$\underbrace{N^3}_{\text{Spatial}} \times \underbrace{N_T}_{\text{Temporal}} \times \underbrace{4}_{\text{Links}} \times \underbrace{9}_{\text{SU(3) matrix}} \times \underbrace{2}_{\mathbb{C}\text{-numbers}} = 72N^3N_T,$$

Which is $\rightarrow 8 \times 72N^3N_T$ bytes.

- The SU(3) matrices are 3×3 matrices of nine complex numbers or 18 real numbers.
- This leads to an absolute **requirement of efficiency**, both in **calculations** and in **input/output**.
- When returning to what ensembles of configurations we generated this will be evident.

Algorithm of choice when sampling gauge configurations.

$$\langle O \rangle = \frac{1}{Z} \int \mathcal{D}A \mathcal{D}\bar{\psi} \mathcal{D}\psi O[\psi, \bar{\psi}, A] e^{-S_G[A] - S_F[\psi, \bar{\psi}, A]}.$$

with

$$Z = \int \mathcal{D}A \mathcal{D}\bar{\psi} \mathcal{D}\psi e^{-S_G[A] - S_F[\psi, \bar{\psi}, A]}.$$

$$\langle O \rangle = \frac{1}{Z} \int \mathcal{D}A O[\psi, \bar{\psi}, A] e^{-S_G[A]}.$$

with


$$Z = \int \mathcal{D}A e^{-S_G[A]}.$$

$$\langle O \rangle = \frac{1}{Z} \int \mathcal{D}A O[\psi, \bar{\psi}, A] e^{-S_G[A]}.$$

with

$$Z = \int \mathcal{D}A e^{-S_G[A]}.$$

The path integral II



17

An example of the discretized path integral, going from time t_0 to t_{N_T} , where the end points is taken to be equal, $x_0 = x_{N_T}$. We integrate over all of space at each time t_i finding the most likely position at a given time.

Measuring observables on the lattice

Gradient flow

Results

Conclusion

Questions?

References

- [1] Hilmar Forkel. A Primer on Instantons in QCD.
arXiv:hep-ph/0009136, September 2000. URL
<http://arxiv.org/abs/hep-ph/0009136>. arXiv:
hep-ph/0009136.