

Solving $SU(3)$ Yang-Mills theory on the lattice: a calculation of selected gauge observables with gradient flow

Hans Mathias Mamen Vege

April 24, 2019

Supervisor: *Andrea Shindler*

Co-supervisor: *Morten Hjorth-*

Jensen

University of Oslo

Introduction

- Quantum Chromodynamics(QCD).

- **QCD.** We will go through and explain what QCD as well as motivate its existence.

- Quantum Chromodynamics(QCD).
- Lattice QCD.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.
- **GLAC.** Will briefly present the code which we developed as well as some benchmarks. We will also present the Metropolis algorithm.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.
- Results.

- **QCD.** We will go through and explain what QCD as well as motivate its existence.
- **LQCD.** We will briefly show how one discretise the lattice and perform calculations on it.
- **Gradient flow.** We will quickly introduce gradient flow and explain its effects.
- **GLAC.** Will briefly present the code which we developed as well as some benchmarks. We will also present the Metropolis algorithm.
- **Results.** We will present the results obtained from pure gauge calculations.

- Quantum Chromodynamics(QCD).
- Lattice QCD.
- Gradient flow.
- Developing a code for solving $SU(3)$ Yang-Mills theory.
- Results.

Quantum Chromodynamics(QCD)

- The standard model: Six quarks and eight gluons

- The standard model.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom

- The standard model.
- Asymptotic freedom.

- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement

- The standard model.
- Asymptotic freedom.
- Confinement.


- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement
- Highly nonlinear due to gluon self-interactions

- The standard model.
- Asymptotic freedom.
- Confinement.
- Nonlinearity.


- The standard model: Six quarks and eight gluons
- Asymptotic freedom
- Confinement
- Highly nonlinear due to gluon self-interactions

- The standard model.
- Asymptotic freedom.
- Confinement.
- Nonlinearity.


The Standard Model



Consists of the innermost square of the six quarks and the gluons.




- The coupling constant **decreases** as we **increase** the energy
- One of the experimental proofs of QCD along with triple γ decay and muon cross section ratio R .



If we try to pull apart **two mesons**, more and more energy is required until we have enough energy to spontaneously create a **quark-antiquark** pair, forming thus **two new mesons**.

The non-linearity of QCD

$$\mathcal{L}_{\text{QCD}} = \sum_{f=1}^{N_f} \bar{\psi}^{(f)} \left(i \not{D} - m^{(f)} \right) \psi^{(f)} - \frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu},$$



Measuring topological charge is a measure of the *Winding number* of the gauge field.




Figure 1: The figure is taken from [1, p. 32].

An illustration of how one can view the winding number given a function f that parametrizes a path around a circle S^1 . Given that it starts and ends at the same point, we have that the number of times it wraps around the circle gives us the winding number.

• A

Lattice Quantum Chromodynamics(LQCD)

• A

Developing a code for solving $SU(3)$ Yang-Mills theory on the lattice

Measuring observables on the lattice

Gradient flow

Results

Conclusion

Questions?

References

- [1] Hilmar Forkel. A Primer on Instantons in QCD.
arXiv:hep-ph/0009136, September 2000. URL
<http://arxiv.org/abs/hep-ph/0009136>. arXiv:
hep-ph/0009136.