

# Impact of cropping system diversification on vegetative and reproductive characteristics of waterhemp (*Amaranthus tuberculatus*)

## Abstract

Corn- and soybean-dominated cropping systems create and maintain a favorable environment for summer annual weeds whose emergence and growth phenology are similar to these annual summer crops. Cropping system diversification can be an effective approach for controlling noxious weeds without increasing reliance on chemical herbicides. Diversification may be especially important for managing waterhemp, a dioecious, summer annual weed that is becoming increasingly prevalent in the US Corn Belt due to its life history characteristics and herbicide resistance profile.

Compared to corn and soybean, alfalfa and oat emerge and establish earlier and are thus more competitive with warm-season weeds like waterhemp. Knowledge of vegetative and reproductive characteristics in a range of crop environments can be valuable for planning weed management strategies. However, most of the relevant characteristics for a population dynamics model were available in corn and soybean monocultures. We examined the relationship between waterhemp's aboveground mass and fecundity under four crop species' presence within three crop rotation systems: a 2-year sequence of corn and soybean; a 3-year sequence of corn, soybean, and oat intercropped with red clover; and a 4-year sequence of corn, soybean, oat intercropped with alfalfa, and alfalfa. All the rotation systems were treated with conventional or reduced rates of herbicides. We established eighteen linear equations to predict waterhemp's fecundity from dried aboveground mass in each crop and associated crop management program since measuring the latter allows for quicker estimation of fecundity compared to counting seeds on each individual plant.

Rotation system and crop phase within rotation system had significant effects on all the response variables but weed control regime on some. The sex ratios at maturity were slightly female-biased in oat and alfalfa. Mature waterhemp plants were larger in corn and soybean than in oat and alfalfa. Oat and alfalfa were planted earlier than corn and soybean and successfully competed for resources against waterhemp despite the absence of herbicide or interrow cultivation. Frequent hay cuts in alfalfa served as physical weed control and contributed to suppressing waterhemp and other weeds substantially.

## Introduction

Cropping system diversification can contribute to the effective suppression of noxious weeds such as velvetleaf (*Abutilon theophrasti* Medik.) (Westerman et al., 2005), giant foxtail (*Setaria faberi* Herrm.) (Liebman et al., 2014), giant ragweed (*Ambrosia trifida* L.) (Liebman and Nichols, 2020), and other species while complementing the effects of herbicides and physical weed control practices (Davis et al., 2012; Weisberger et al., 2019). Much of the effectiveness of diversified cropping systems for weed suppression can be attributed to differences among crop species in their phenologies and the management techniques applied to them. Differences in crop phenology and diverse management tactics can lead to a net loss in weed seed population density in the soil seed bank (MacLaren et al., 2020; Liebman and Gallandt, 1997) resulting from reductions in weed fecundity and increased consumption of weed seeds by granivores. The present study focuses on fecundity and other relevant individual- and population-level reproductive and vegetative characteristics of waterhemp (*Amaranthus tuberculatus* (Moq.) J. D. Sauer) in three rain-fed cropping systems differing in crop species richness and management practices.

Waterhemp is a dioecious, summer annual, dicotyledonous species that has been listed as one of the five most noxious weeds in row crops in the US based on the number of times this species appears in the literature (Johnson et al., 2009) and by growers' concern across 22 states in the U.S. (Prince et al., 2012). At least 54 waterhemp populations are resistant to up to five herbicide modes of action as of 2021 (Heap, 2021). As a dioecious species, populations of waterhemp are expected to express a 1:1 male:female ratio (Grant, 1959; Costea et al., 2005; Heneghan and Johnson, 2017). However, waterhemp has three characteristics that favor female-biasedness under conditions of no stress, according to a study of 243 dioecious species excluding waterhemp (Field et al., 2013): 1) the male sex is heterogametic (Montgomery et al., 2021); 2) the species has abiotic pollination and seed dispersal; and 3) the fruits are non-fleshy (Costea et al., 2005). A stressed waterhemp population tends to be female-biased, with up to ten females per male (Pratt and Clark, 2001), which is consistent with the general pattern of sexually differentiated stress tolerance in herbaceous plants (38 species, excluding waterhemp, Juvany and Munné-Bosch, 2015). The sex ratio plasticity of waterhemp suggests that a stressed population, which is characterized by low density, may allocate available resources to produce more female offspring as an effort to increase population density, as observed in its close relative, Palmer amaranth (*A. palmeri*) (Korres and Norsworthy, 2017; Mesgaran et al., 2019). Knowing how sex ratios may deviate from parity under different biotic and abiotic conditions could inform how a waterhemp population might progress from one generation to the next.

Waterhemp management is agronomically challenging because of a suite of life history characteristics, including a persistent soil seedbank (Davis, 2008), an extended seedling emergence pattern (Buhler and Hartzler, 2001), high relative growth rate, high fecundity (Heneghan and Johnson, 2017), and rapid herbicide resistance development (Tranel, 2021). One year of prolific seed production can replenish a declining seedbank with more seeds than existed in the seedbank (Davis, 2008). Failing to control waterhemp can cause up to 43% yield loss in soybean (*Glycine max* (L.) Merr.) (Hager et al., 2002) and 74% yield loss in corn (*Zea mays* L.) (Steckel and Sprague, 2004). The accumulated mass and density of a weed population reflect the relative competitiveness against crops and the favorability of the environment, which in turn could signal the effectiveness of weed management throughout the season. From a planning perspective, population density at maturity and plant fecundity are useful for estimating the density of new seeds produced and potentially added to the soil seedbank, and for adjusting weed management regimes accordingly (Buhler et al., 1997). Waterhemp's fecundity has been studied in corn and soybean crops (Menalled et al., 2004; Nordby and Hartzler, 2004) but not in other crops' or in an extended crop rotation system. Alfalfa (*Medicago sativa* L.), red clover (*Trifolium pratense* L.), and oat (*Avena sativa* L.) are cool-season crops that can be grown in rotation sequences with corn and soybean, whereas waterhemp is a summer annual weed. Compared to corn and soybean, alfalfa, red clover, and oat seedlings emerge and establish earlier. Alfalfa, red clover, and oat also emerge and establish earlier than a number of summer annual weed species, including waterhemp (Buhler et al., 2008; Horak and Loughin, 2000).

In the present study, we examined the population aboveground mass, density and sex ratio, and the relationship between waterhemp's female size and fecundity when the weed grew in association with five crop species (corn, soybean, oat, red clover, and alfalfa) arranged in three rain-fed cropping systems. Assessing waterhemp characteristics in the presence of oat intercropped with red clover or alfalfa and alfalfa grown as a sole crop as well as corn and soybean could help to fill the gap of information concerning waterhemp performance in extended rotations. We hypothesized that the sex ratio of a waterhemp population deviated from parity depending on the environment's favorability, but how much and to which direction the shift would occur would depend on how much the studied population was suppressed. In addition to informing management, individual- and population-level characteristics would provide useful contextual details for sex ratio comparison. Counting seeds for waterhemp fecundity assessment is time-consuming and laborious, so it would be convenient to extrapolate fecundity from plant mass. We hypothesized that regression relationships with which to predict fecundity from plant mass could be identified but that such relationships would differ among treatments, due to differences in crop phenology, crop-weed competition, and management practices.

## Materials and Methods

### Experiment Design

Empirical measurements of waterhemp biomass and fecundity were made in 2018 at Iowa State University’s Marsden Farm in Boone County, Iowa, USA, (42° 01’N, 93° 47’W, 333 m above sea level). The site description and crop management practices were described by Liebman et al. (2021). The experiment was initiated in 2001 on a 9-hectare field to compare the performance of three different crop rotations. In 2008, the experiment was reorganized to allow comparison of two contrasting weed management regimes in each of the three rotation systems. Since 2001, within each of the four blocks, plots were randomly assigned to one of the crop phases within one of three rotations.

In the present study, the experiment comprised of 72 experimental units, each 9 m x 84 m. The experimental units (eu) were arranged in a randomized complete block split-plot design with four replications. The experiment was two-way factorial, with crop rotations comprising main plots and weed management regimes comprising split plots. In any year, all crop phases within the same rotation were present in different plots in the same replicate block to avoid confounding effects of the year with those of the treatments (Payne, 2015). The 2-year, 3-year, and 4-year crop sequences included in the study are shown in Table 1. The conceptual diagram for the experiment is shown in Figure 1. Alfalfa in this experiment was sown with oat. Oat was harvested for grain in 2018 and for hay in 2019 and alfalfa was retained over the winter and used as a hay crop the following year. Oat in our experiment was intended for grain harvest but was harvested for hay in 2019 due to severe hail damage to the crop and subsequent weed infestation. In the conventional weed management regime, herbicide was broadcast on the whole area that was planted to corn, whereas in the low herbicide regime, herbicide was applied in 38-cm bands over corn rows, and interrow areas were cultivated. Even though corn was the only crop that received the contrasting weed management regimes, all other crops were identified with the weed management applied to the corn phase (hereafter referred to as *corn weed management*), to which they followed. To improve weed control efficacy, herbicides used for soybean differed between 2018 and 2019. Details concerning crop genotypes and management practices are shown in Table 2.

Table 1: Crop phases of the three rotation systems present in a replicate block in 2018 and 2019

	2-year	2-year	3-year	3-year	3-year	4-year	4-year	4-year	4-year
2018	corn	soybean	corn	soybean	oat/red clover	corn	soybean	oat/alfalfa	alfalfa
2019	soybean	corn	soybean	oat/red clover	corn	soybean	oat/alfalfa	alfalfa	corn

Table 2: Crop varieties, and dates and rates for management operations in 2018 and 2019. Soybean germination in 2018 was lower than in 2019 because of poor drainage in the soil. Oat and red clover were intercropped in the 3-year system. Oat and alfalfa were intercropped in the third year of the 4-year system and alfalfa was overwintered after oat harvest.

Rotation	Crop	Hybrid or cultivar	Planting date	Harvest date	Seed density	Crop density	Interrow	Cultivation	Herbicide (kg ai/ha)
<b>2018 season</b>									
all	corn	Epley 1420	May. 8	Oct. 30	seeds m <sup>-2</sup> 8	plants m <sup>-2</sup> 8	cm 76	low: Jun. 4; conv: none	low: tembotrione (0.054); conv: PRE thiencazone methyl (0.037), isoxaflutole (0.093); POST: mesotrione (0.105), nicosulfuron (0.053)
all	soybean	Latham 2758 R2	Jun. 3	Oct. 29	35	18	76	none	flumioxazin (0.096); POST: glyphosate as potassium salt (1.540), lactofen (0.140)
3- and 4-year	oat	INO9201	Apr. 24	Jul. 20	kg m <sup>-2</sup> 0.009	plants m <sup>-2</sup> 225 (3-year) and 236 (4-year)	cm 20	none	none
3-year	red clover	Mammoth Red	Apr. 24		0.002	187	20	none	none
4-year	alfalfa	55H94	Apr. 12, 2017	Jun. 4, Jul. 9, and Sep. 10	0.002	154	20	none	none
<b>2019 season</b>									
all	corn	Epley 1730	Jun. 3	Nov. 6	seeds m <sup>-2</sup> 8	plants m <sup>-2</sup> 8	cm 76	none	low: tembotrione (0.0054); conv: PRE: thiencazone methyl (0.037), isoxaflutole (0.093); POST: mesotrione (0.105), nicosulfuron (0.053)
all	soybean	Latham 2758 R2	Jun. 10	Oct. 18	35	31	76	none	PRE: flumioxazin (0.096); POST: glufosinate ammonium (0.594), clethodim (0.136)
3- and 4-year	oat	INO9201	Apr. 16	Jul. 22 and 24	kg m <sup>-2</sup> 0.009	plants m <sup>-2</sup> 366 (3-year) and 330 (4-year)	cm 20	none none	none none
3-year	red clover	Mammoth	Apr. 16		0.002	219	20	none	none
4-year	alfalfa	WS Leafguard	Apr. 24	Jun. 7, Jul. 12, Aug. 26,	0.002	176	20	none	none




Figure 1: Conceptual diagram of the three rotation systems compared within the experiment. A cycle of four calendar years is shown as four phases. Crops are color-coded and displayed for the approximate months that they were present. Emergence and establishment of common waterhemp plants are illustrated with black symbols. Grey plants shown in oat intercropped with red clover or alfalfa's first year were physically controlled by crop harvest operations. Red clover was not harvested but kept as a living cover crop until senescence. Grey plants shown in alfalfa's second year were physically suppressed three to four times by hay harvest. Alfalfa's hay was harvested when approximately 5% of the plants flowered.

## Sample collection

Samples for all response variables were collected at least 3m in from the border of each eu to avoid the possible edge effects. For each response variable, samples were collected from eight quadrats per eu to account for the patchiness of the weed populations. In 2018, the sex ratio was assessed by scouting the whole eu to obtain higher degrees of freedom. The date of sample collection and total sampled area for population- and individual-based measurements are shown in Table 3.

Table 3: Sampling dates and areas in 2018 and 2019

Measurement	Crop	2018		2019	
		Collection date	sampled area (m <sup>2</sup> )	Collection date	Sampled area (m <sup>2</sup> )
Population above-ground mass	corn	Sep. 11, 12, 13	148.3	Sep. 17, 18, 19	148.3
	soybean	Sep. 17, 19, 21	148.3	Sep. 30	148.3
	oat	Sep. 27; Oct. 4, 15, 16, 18, 19	16	Sep 23, 25, 26; Oct. 3, 4, 7	17.9
	alfalfa	Sep. 26, 27; Oct 16, 19	16	Sep 24, 25; Oct. 3, 7	17.9
Population density	corn	Sep. 11, 12, 13	148.3	Sep. 17, 18, 19	148.3
	soybean	Sep. 17, 19, 21	148.3	Sep. 30	148.3
	oat	Sep. 27; Oct. 4, 15, 16, 18, 19	16	Sep 23, 25, 26; Oct. 3, 4, 7	17.9
	alfalfa	Sep. 26, 27; Oct 16, 19	16	Sep 24, 25; Oct. 3, 7	17.9
Population sex ratio	corn	Sep. 10, 11, and 12	729	Sep. 17, 18, and 19	148.3
	soybean	Sep. 17 and 30	729	Sep. 30	148.3
	oat	Oct. 4, 15, and 18	729	Sep. 24, 25, 26, 30, Oct. 1 and 2	17.9
	alfalfa	Nov. 1	729	Oct. 3 and 4	17.9
Individual female above-ground mass	corn	Oct. 18, 19, 22, 24, and 25	729	Sep. 17, 18, and 19	148.3
	soybean	Oct. 4, 15, and 18	729	Sep. 30	148.3
	oat	Oct. 29, 30, 31, Nov. 1	729	Sep. 24, 25, 26, 30, Oct. 1 and 2	17.9
	alfalfa	Nov. 1	729	Oct. 3 and 4	17.9
Individual female fecundity	corn	Oct. 24 and 25	729	none	none
	soybean	Oct. 4, 15, and 18	729	none	none
	oat	Oct. 29, 30, 31, Nov. 1	729	none	none
	alfalfa	Nov. 1	729	none	none

## Population sex ratio

In 2018 we scouted each eu until 100 or all the available plants (if the number of plants available was fewer than 100) were sexed to obtain greater degrees of freedom. We determined the sex of 252, 1999, 2426, and 895 waterhemp plants in alfalfa, oat, soybean, and corn, respectively. In 2019, eight quadrats per eu were marked at the beginning of the season and fixed until crop harvest for a census. Overall, 413, 1331, 0, and 553 waterhemp plants were sexed in alfalfa, oat, soybean, and corn, respectively. Zero observations in all the soybean eu's resulted from high herbicide efficacy, so the 2019 data was imputed (Appendix B).

**Population aboveground mass and density** The quadrats were randomly placed in a 4 x 2 grid at the sampling date and were non-overlapping with the 2019 census quadrats. The number of plants and the total dried biomass was tallied by eu.

**Individual female aboveground mass and fecundity** The maturation of waterhemp seeds can take 20 days from pollination (Bell and Tranel, 2010). We harvested female waterhemp plants as close to crop harvest as possible to maximize the number of mature seeds on mother plants. Prior to sample collection in an eu, the whole area was visually inspected to estimate the difference in plant sizes. Specimens were then collected to best capture the range of within-eu variance. Given the time and labor constraints, we planned to collect eight intact plants from each eu, which was equivalent to 576 plants in total. Plants had to be identifiable per Uva et al. (1997). By the time the seeds reached maturity, 389 intact plants were collected and processed. No intact plants were collected from two eu's. Plant specimens were contained individually in tightly knitted fabric bags to prevent seed loss. The detailed procedure for seed cleaning and counting is provided in Appendix A.

## Model fitting and selection

Waterhemp survival in soybean was the greatest among all crops in the experiment in 2018 but the least in 2019 because of the use of different herbicide active ingredients. Given year-to-year differences in the sampling scheme and herbicide efficacy, the two years of data were thus analyzed separately for all the response variables in R version 4.1.2 (R Development Core Team, 2021). The data was curated with the **tidyverse** package version 1.3.1 (Wickham and RStudio, 2021).

In 2018, 2% of the sex ratio data was missing due to zero observations in one eu, so complete case analysis, in which eu's of known sex ratio were retained while unknown sex ratio were removed from the data set, was used. In 2019, 22% of the sex ratio data was missing, so the data were imputed with predictive mean matching (PMM) method with the **mice** package version 3.14.0, (van Buuren and Groothuis-Oudshoorn, 2011) to optimally replace missing data with meaningful values without altering the observed sex ratios (Appendix B). Any model that involved the imputed data was fitted on all the produced (imputed) data sets, and the results were pooled (White et al., 2011).

Block was included in all models as a fixed factor because blocks were used to control the different field conditions across sections, and thus to reduce variance between eu's (Dixon, 2016). All the models were first fitted full, with block, corn weed management and crop identity, the interaction of corn weed management and crop identity, and covariates when applicable. Crop identities are the combination of crop species and the rotation to which they belonged. The within-eu variation was random and absorbed in the random error term in each model equation. The fitted models were linear (**lm**), generalized linear (**glm**), or generalized least square (**glms**) depending on the data structure and the nature of the response variable. The response and quantitative variables were appropriately transformed as needed to obtain homogeneous variances. Half of the minimum, non-zero value among all the observations was added to all the observations before ln-transformation to replace zeros. Response variables that were all non-zero were ln-transformed without adjustment. The goodness of fit of each model was assessed with diagnosis plots and mean squared error (MSE) of the variance.

The marginal means of each response variable were estimated with the **emmeans** function from the **emmeans** package version 1.7.2 (Lenth et al., 2021) to accommodate non-integer and unequal degrees of freedom among groups. Marginal means were averaged over blocks for post-ANOVA or post-ANCOVA contrasts and over

factors whose effects were non-significant. Degree of freedom adjustment was done with the Satterthwaite method for the `gls` and Kenward-Roger method for the `glm` and `lm` models. ANCOVA (analysis of covariance) was applied to examine the effect of treatments on the relationship between female aboveground biomass and fecundity and between population sex ratio and biomass or density at maturity. ANCOVA combines regression and ANOVA (analysis of variance) to improve precision in mean estimation as compared to ANOVA estimation (Yang and Juskiw, 2011). Type III sums of squares error were calculated with the `emmeans`'s `joint_tests` function to accommodate unbalanced data with interaction when occurred and to avoid misleading assessment of factors' effects based on their sequential order in the model.

### Population sex ratio at maturity

A logistic regression model was fitted with the `glm` command and `family = quasibinomial(link = logit)` argument specification to analyze sex ratio. The `quasibinomial` family with follow-up F-test was used to accommodate overdispersion and `logit` link function transformed the sex ratio using the natural logarithm (ln) (Crawley, 2013).

$$S_{ijk} = \text{Binomial}(N_{ijk}, \pi_{ijk})$$

$$\ln \frac{\pi_{ijk}}{1 - \pi_{ijk}} = \mu + b_k + \alpha_i + \gamma_j + \alpha_i \gamma_j + \epsilon_{ijk} \quad (1)$$

where,

$S_{ijk}$  is the number of female plants among all the  $N_{ijk}$  plants in block  $k^{th}$  under crop identity  $i^{th}$  and corn weed management  $j^{th}$ ,  
 $\ln \frac{\pi_{ijk}}{1 - \pi_{ijk}}$  is the logit transformation of  $S_{ijk}$ ,  
 $\mu$  is the overall mean female proportion, the intercept,  
 $\alpha_i$  is the effect of the  $i^{th}$  crop identity,  
 $\gamma_j$  is the effect of the  $j^{th}$  corn weed management,  
 $b_k$  is the block effect,  
 $\alpha_i \gamma_j$  is the interaction effect of crop identity and corn weed management, and  
 $\epsilon_{ijk}$  is the random error.

### Population aboveground mass and density

A linear regression model was fitted with the `lm` command on each of the two variables, population aboveground mass or stand density. The general model equation for these response variables is

$$Y_{ijk} = \mu + b_k + \alpha_i + \gamma_j + \alpha_i \gamma_j + \epsilon_{ijk} \quad (2)$$

where,

$Y_{ijk}$  is either the ln-transformed population aboveground mass or ln-transformed stand density in block  $k^{th}$  under crop identity  $i^{th}$  and corn weed management  $j^{th}$ , and  
other terms as defined in Equation (1).

### ANCOVA of population sex ratio, aboveground mass, and density

The regression of sex ratio against population density or biomass was extended from the ANOVA of sex ratio (Equation (1)).

$$S_{ijk} = \text{Binomial}(N_{ijk}, \pi_{ijk})$$



$$\ln \frac{\pi_{ijk}}{1 - \pi_{ijk}} = \mu + b_k + \alpha_i + \gamma_j + \alpha_i \gamma_j + \delta D_{ijk} + \alpha_i D_{ijk} + \gamma_j D_{ijk} + (\alpha_i \gamma_j) D_{ijk} + \epsilon_{ijk} \quad (3)$$

where,

$\delta$  is the effect of the covariate,

$D_{ijk}$  is the natural log-transformed population stand density in block  $k^{th}$  under crop identity  $i^{th}$ , and corn weed management  $j^{th}$ , the covariate, and other terms as defined in Equation (1).

#### Individual female aboveground mass and fecundity

A compound symmetric linear regression model (with `nlme` package's `gls` command was first fitted for each of the response variables, individual aboveground mass and fecundity to accommodate negative variance that occurred when the within-eu variance was larger than the between-eu variance and correlated errors occurred within blocks (version 3.1-153, Pinheiro et al., 2021). The `corCompSymm` argument in the `gls` command was specified by identifying unique combinations of block and treatment. The model in this exercise is of the same form as the model in Equation (2):

$$Y_{ijkl} = \mu + b_k + \alpha_i + \gamma_j + \alpha_i \gamma_j + \epsilon_{ijkl} \quad (4)$$

where,

$Y_{ijkl}$  is either the ln-transformed aboveground mass or ln-transformed number of seeds of female plant  $l^{th}$  in block  $k^{th}$  under crop identity  $i^{th}$  and corn weed management  $j^{th}$ ,  $\epsilon_{ijkl}$  is the random error, and other terms as defined in Equation (1).

#### Individual female aboveground mass and fecundity relationship

The regression of individual plant fecundity against individual plant aboveground mass was combined from the ANOVA of each (Equation (4)) to establish a relationship between the two variables:

$$Y_{ijkl} = \mu + b_k + \alpha_i + \gamma_j + \alpha_i \gamma_j + \beta X_{ijkl} + \alpha_i X_{ijkl} + \gamma_j X_{ijkl} + (\alpha_i \gamma_j) X_{ijkl} + \epsilon_{ijkl} \quad (5)$$

where,

$Y_{ijkl}$  is the ln-transformed number of seeds of plant  $l^{th}$  in block  $k^{th}$  under crop identity  $i^{th}$  and corn weed management  $j^{th}$ ,

$X_{ijkl}$  is the ln-transformed dried aboveground mass of plant  $l^{th}$  in block  $k^{th}$  under crop identity  $i^{th}$ , and corn weed management  $j^{th}$ , the covariate, and other terms as defined in Equation (1).

If the MSE of the model described in Equation (5) were smaller than that of the model described in Equation (4), the former model would be considered better for estimating the number of seeds produced by an individual waterhemp plant.

We tested the assumption that all the regression lines were parallel. Violation of this assumption required an individual regression line for each treatment. To test model robustness, samples in each treatment were pooled across four blocks and divided into four size-based subsets. Samples from each subset were then randomly placed into the testing and training sets using the 80 testing : 20 training ratio. A model was considered to perform well if the data points in the testing set blended well with the data points from the training set. A robust model could be used to predict plant fecundity with new biomass data.

## Results

Using `ggResidpanel` version 0.3.0 (Goode and Rey, 2019), `boot` version 1.3-28 (Canty and Ripley, 2021), and two customized functions for model diagnosis, no predictable pattern in the plots of residuals versus predicted values suggests that the analysis models fit the data well (Details are provided in Nguyen and Liebman (2022)). In all rotations, all the crop yields were comparable to those of Iowa and Boone County where the experiment is situated (Hunt et al., 2020; Nguyen and Liebman, in review). Tables were compiled with `kableExtra` version 1.3.4 (Zhu et al., 2021). Figures were made with `emmeans` version 1.7.2 (Lenth et al., 2021) and `ggplot2` version 3.3.5 (Wickham et al., 2021).

### Population aboveground mass and stand density

In both years, population aboveground mass and stand density were strongly influenced by crop identity (Table 4). The rotation system in which a crop was grown also affected population aboveground mass and stand density, although not consistently between years (Table 5).

In 2018, population aboveground mass in the same crop species was comparable across rotations except for corn grown in the 2-year (C2) versus 4-year rotation (C4) (p-value = 0.043). In 2019, population aboveground mass in the same crop was different across rotations, except for corn in the 2-year (C2) versus 3-year rotation (C3) (p-value = 0.968) and soybean in the 3-year (S3) versus 4-year rotation (S4) (p-value = 1). Averaged across rotations, population aboveground mass was comparable in 2018 for corn versus alfalfa (p-value = 0.2509) and soybean versus oat (p-value = 0.504), but 10- to 5795.38-fold different in the other ten pairs of comparison (p-values < 0.01).

In 2018, population stand density in the same crop species was comparable across the rotations. In 2019, population stand density in the same crop species was comparable for soybean (p-values = 0.1256 and 1) and C2 versus C3 (p-value = 0.9284), but significantly different for the other corn comparisons and for oat in the 3-year (O3) versus 4-year rotation (O4). Averaged over rotations, population stand density was comparable in 2018 between soybean and alfalfa (p-value = 0.766), but 5- to 1330-fold different in the other eight pairs of comparison (p-values < 0.001).

In 2018, population aboveground mass was the highest in soybean and oat (Figure 2A) because soybean weed management was ineffective and herbicide was intentionally not applied in oat. The legacy of an ineffective weed management program in 2018 soybean plots was observed in 2019 oat plots where population aboveground mass and stand density were the highest among all the crop identities (Figure 2B). High stand density in 2019 oat plots was also due to uneven oat establishment. The change in 2019 in weed management for soybean substantially reduced the waterhemp pressure on soybean (Figure 2B and 2D).

Table 4: ANOVAs of crop identity and corn weed management effects on waterhemp population aboveground mass and stand density. Crop identity was the only influential factor on both population aboveground mass and stand density in 2018 and 2019.

Source of variation	df1	df2	2018				2019			
			Population aboveground mass		Population stand density		Population aboveground mass		Population stand density	
			F.value	p.value	F.value	p.value	F.value	p.value	F.value	p.value
Crop ID	8	51	21.23	<.0001	27.45	<.0001	42.14	<.0001	84.03	<.0001
Corn weed management	1	51	0.41	0.5241	0.87	0.3555	1.23	0.2730	0.30	0.5889
Crop ID x Corn weed management	8	51	0.40	0.9139	0.97	0.4736	1.35	0.2415	0.63	0.7486




Figure 2: Waterhemp population aboveground mass and stand density averaged over corn weed managements. The abbreviations on the x-axis are crop identities, which are the combinations of the first letter in crop species names and the rotation to which the crops belonged (C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation; S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, O4: oat in the 4-year rotation, and A4: alfalfa in the 4-year rotation). The black dots are estimated marginal means. The blue bars are 95% confidence intervals. The red arrows reflect the comparison among means. Overlapping arrows indicate non-significant differences.

Table 5: Rotation system and crop species effects on population aboveground mass and stand density.

Contrast	2018				2019			
	Population aboveground mass		Population stand density		Population aboveground mass		Population stand density	
	ratio	p.value	ratio	p.value	ratio	p.value	ratio	p.value
<b>(A) - Rotation system effects</b>								
C2 vs C3	12.54	0.1237	2.38	0.2990	0.84	0.9680	1.19	0.9284
C2 vs C4	23.12	0.0430	1.84	0.5435	10.61	0.0054	14.45	<.0001
C3 vs C4	1.84	0.8798	0.78	0.8983	12.65	0.0026	12.11	<.0001
S2 vs S3	6.42	0.3151	2.07	0.4258	6.27	0.0369	2.60	0.1265
S2 vs S4	2.45	0.7597	1.29	0.8976	6.27	0.0369	2.60	0.1265
S3 vs S4	0.38	0.7298	0.62	0.6963	1.00	1.0000	1.00	1.0000
O3 vs O4	0.82	0.8774	1.78	0.3235	0.14	0.0096	0.29	0.0132
<b>(B) - Crop species effects</b>								
oat vs soybean	3.15	0.5040	47.18	<.0001	2580.00	<.0001	1330.00	<.0001
oat vs corn	5795.38	<.0001	222.88	<.0001	85.30	<.0001	32.60	<.0001
oat vs alfalfa	831.33	<.0001	29.84	<.0001	8.34	0.0071	2.39	0.1712
soybean vs corn	1840.55	<.0001	4.72	0.0001	0.03	<.0001	0.02	<.0001
soybean vs alfalfa	264.02	<.0001	0.63	0.7660	0.00	<.0001	0.00	<.0001
corn vs alfalfa	0.14	0.2509	0.13	0.0005	0.10	0.0014	0.07	<.0001

*Note:* Some zero values are due to rounding. C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation; S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, and O4: oat in the 4-year rotation

## Population sex ratio

Population stand density was included to improve the precision of estimates of population sex ratios (Table 6C versus 6A). The population sex ratio in 2018 differed significantly among treatments, at different population stand densities within each treatment (p-value = 0.0155, Table 6 and Figure 3). Therefore, sex ratios in the same treatment were evaluated at four population densities, i.e., 1, 5, 50, and 500 plants/m<sup>2</sup>, to illustrate that three-way interaction (Figure 3). Female-biasedness was more likely if a waterhemp population was grown in oat and alfalfa. None of the waterhemp populations grown in corn and soybean expressed gender biasedness. It is unclear whether the corn weed management program had a significant effect on gender biasedness given the magnitude of the variance (Figure 3).

We defined a useful imputed data set to be a set that resulted in fully estimable marginal means for sex ratio comparison across all treatments, which was achievable with non-zeros in female and male categories in at least one replication among the four blocks for the missing observations in the 2019 original sex data. Unlike the 2018 data, the sex ratio in 2019 was analyzed without the covariates because none of the covariates improved the goodness of fit for the analysis model. With  $m = 24$ , five imputed data sets were useful (Appendix B). The significance and influence of treatment factors and their interaction in the imputed data sets for waterhemp sex ratio in 2019 were consistent with those of the 2018 data (Figure 4). In 21 out of 24 sets, sex ratio in 2019 was affected by crop identity (Figure 4B and 4C). Female biasedness was observed in oat and alfalfa but not in corn and soybean (Figure 4A).

Table 6: ANOVAs of crop identity, herbicide, and covariate effects on population sex ratio using 2018 data. With population aboveground mass covariate included (B), crop identity was the only influential factor on population sex ratio. With population stand density covariate included (C), sex ratio responded differently in each treatment and stand density combination.

Source of variation	df1	df2	F.value	p.value
<b>(A) no covariate. Residual deviance = 165.9, dispersion = 3.32.</b>				
Crop ID	8	Inf	8.45	<.0001
Corn weed management	1	Inf	0.01	0.9317
Crop ID x Corn weed management	8	Inf	0.46	0.8862
<b>(B) with population aboveground mass covariate. Residual deviance = 104.3, dispersion = 3.24.</b>				
Crop ID	8	Inf	1.02	0.4155
Corn weed management	1	Inf	0.00	0.9601
Population aboveground mass	1	Inf	1.51	0.2198
Crop ID x Corn weed management	8	Inf	0.71	0.6847
Crop ID x Population aboveground mass	8	Inf	1.04	0.4038
Corn weed management x Population aboveground mass	1	Inf	2.85	0.0916
Crop ID x Corn weed management x Population aboveground mass	8	Inf	1.24	0.2713
<b>(C) with population stand density covariate. Residual deviance = 82.12, dispersion = 2.54.</b>				
Crop ID	8	Inf	0.96	0.4679
Corn weed management	1	Inf	0.93	0.3346
Population stand density	1	Inf	2.46	0.1169
Crop ID x Corn weed management	8	Inf	1.46	0.1675
Crop ID x Population stand density	8	Inf	1.71	0.0896
Corn weed management x Population stand density	1	Inf	5.16	0.0231
Crop ID x Corn weed management x Population stand density	8	Inf	2.36	0.0155

## Individual female aboveground mass and fecundity

Individual female aboveground mass and fecundity were affected by rotation, crop species, and corn weed management (Table 7). Crop identity was more influential on female aboveground mass and fecundity than corn weed management regime, but the effect of crop identity differed between corn weed management regimes (Table 8A and 8B). Differences in relative female size and fecundity across rotation by herbicide treatments were attributed to the relative size and fecundity differences when the waterhemp populations grew in different crops' presence.

Individual female aboveground mass was comparable in most pairwise comparison of the same crop species in different rotations, except S2 versus S3 (p-value = 0.0076) and S3 versus S4 (p-value = 0.0268) that followed corn under conventional weed management and C2 versus C3 (p-value = 0.0064) under low weed management. Averaged over rotations, individual female aboveground mass was 3.51- to 133.62-fold different across each pair of comparison (p-values < 0.05), except for corn under low weed management versus the succeeding oat (p-value = 0.9616).

Individual fecundity was comparable in most pairwise comparison of the same crop species in different rotations, except S2 versus S3 (p-value = 0.001) and S3 versus S4 (p-value = 0.0046) that followed corn under conventional weed management and C2 versus C3 under low weed management (p-value = 0.0032), and O3 versus O4 that followed corn under low weed management (p-value = 0.0321). Averaged over rotations, individual fecundity was comparable between corn under low herbicide and oat in the same system (p-value = 0.4904) but was 6.31- to 6857.12-fold different in other pairs of comparison (p-values  $\leq$  0.0001).




Figure 3: Waterhemp population sex ratios under 54 combinations of experimental treatments and population stand densities. The abbreviations on the x-axis are crop identities, which are the combinations of the first letter in crop species names and the rotation to which the crops belonged (C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation; S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, O4: oat in the 4-year rotation, and A4: alfalfa in the 4-year rotation). The dashed lines mark sex ratio parity. The black dots are estimated marginal means. The blue bars are 95% confidence intervals. The red arrows reflect the comparisons among means. Overlapping arrows indicate non-significant differences.




Figure 4: Waterhemp population sex ratios under nine crop identities averaged over two Corn weed management regimes using 2019's 24 imputed data sets (A). The abbreviations on the panel A's sections are crop identities, which are the combinations of the first letter in crop species names and the rotation to which the crops belonged (C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation, S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, O4: oat in the 4-year rotation, and A4: alfalfa in the 4-year rotation). The dashed lines mark sex ratio parity in panel A and level of confidence in panel B, respectively. The blank spaces are nonestimable values. The triangulars and circles in panel A represent female-biased and even populations assessed at  $\alpha = 0.05$ , respectively. F-ratios for sources of variation are shown in panel C.



Table 7: Rotation system and crop species effects on individual female aboveground mass and fecundity.

Contrast	Female individual aboveground mass				Individual fecundity			
	Corn weed management: conventional		Corn weed management: low		Corn weed management: conventional		Corn weed management: low	
	ratio	p.value	ratio	p.value	ratio	p.value	ratio	p.value
<b>(A) - Rotation system effects</b>								
C2 vs C3	2.62	0.1335	4.29	0.0064	3.95	0.0819	7.29	0.0032
C2 vs C4	2.12	0.3402	2.27	0.1613	3.00	0.2367	2.55	0.2288
C3 vs C4	0.81	0.9302	0.53	0.4070	0.76	0.9240	0.35	0.2253
S2 vs S3	0.18	0.0076	0.33	0.2005	0.07	0.0010	0.47	0.6323
S2 vs S4	0.70	0.7885	0.76	0.9068	0.60	0.7451	1.19	0.9782
S3 vs S4	3.81	0.0268	2.27	0.3553	8.45	0.0045	2.51	0.4525
O3 vs O4	0.93	0.8695	0.39	0.0457	0.62	0.4363	0.27	0.0321
<b>(B) - Crop species effects</b>								
soybean vs corn	8.64	<.0001	35.10	<.0001	17.51	<.0001	96.74	<.0001
soybean vs oat	36.55	<.0001	30.29	<.0001	110.44	<.0001	55.97	<.0001
soybean vs alfalfa	128.16	<.0001	133.62	<.0001	5423.32	<.0001	6857.12	<.0001
corn vs oat	4.23	0.0001	0.86	0.9616	6.31	0.0001	0.58	0.4904
corn vs alfalfa	14.83	<.0001	3.81	0.0099	309.73	<.0001	70.88	<.0001
oat vs alfalfa	3.51	0.0324	4.41	0.0062	49.11	<.0001	122.51	<.0001

*Note:* C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation, S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, and O4: oat in the 4-year rotation

## Effects of weed management regimes and rotations on female aboveground mass and fecundity relationship

Since the treatment effects were statistically significant for both female aboveground mass and fecundity (Table 8), we proceeded with finding the slopes and intercepts for each linear regression of fecundity against biomass. Different slopes were specified by including interaction terms between the covariate and treatment factors. A regression slope for each treatment was necessary. The training and testing sets' data points were well mingled indicated that the established equations were robust (Figure 5). That the equations in Table 9 could predict waterhemp fecundity parsimoniously from dried aboveground mass using the relevant context of crop and crop management. The presented means and SEs for the estimated intercepts and slopes were established from the whole data set.

Table 8: ANOVAs for the effects of crop identity, corn weed management, and female aboveground mass on individual female aboveground mass (A), fecundity (B), and fecundity with aboveground mass covariate (C). Each combination of crop identity and corn weed management affected female aboveground mass and fecundity differently.

Source of variation	df1	df2	F.value	p.value
<b>(A) - Individual female aboveground mass. MSE = 2.02</b>				
Crop ID	8	46.56	48.83	<.0001
Corn weed management	1	158.23	13.57	0.0003
Crop ID x Corn weed management	8	73.81	2.36	0.0255
<b>(B) - Individual fecundity. MSE = 3.43</b>				
Crop ID	8	41.67	72.13	<.0001
Corn weed management	1	146.29	14.64	0.0002
Crop ID x Corn weed management	8	63.96	2.98	0.0067
<b>(C) - Individual fecundity with individual aboveground mass covariate. MSE = 1.01</b>				
Crop ID	8	67.83	16.53	<.0001
Corn weed management	1	312.06	2.92	0.0886
Biomass	1	349.07	483.09	<.0001
Crop ID x Corn weed management	8	151.00	1.66	0.1136
Crop ID x Biomass	8	300.20	2.99	0.0031
Corn weed management x Biomass	1	349.21	2.84	0.0931
Crop ID x Corn weed management x Biomass	8	333.10	2.49	0.0122

Table 9: Means and SEs for estimated linear regression of waterhemp fecundity index ( $\ln(\text{seeds} + 1)$ ) versus biomass index ( $\ln(\text{gram} + 0.005)$ ) indices intercepts and slopes, accompanied by the  $R^2$  values of each equations. The abbreviations in the Crop ID column are crop identities, which are the combinations of the first letter in crop species names and the rotation to which the crops belonged.  $R^2$  values were calculated from the whole data set (training and testing sets), and thus, were slightly different from those in Figure 5.

Effect		Intercept		Slope		$R^2$
Crop ID	Corn weed management	Estimate	Std.error	Estimate	Std.error	
C2	conventional	6.07	0.18	1.24	0.08	0.89
C2	low	5.88	0.22	1.22	0.11	0.78
S2	conventional	6.30	0.31	1.14	0.11	0.89
S2	low	7.07	0.22	0.97	0.07	0.96
C3	conventional	5.86	0.25	1.26	0.14	0.83
C3	low	5.11	0.35	0.66	0.21	0.33
S3	conventional	7.25	0.44	0.96	0.09	0.84
S3	low	4.89	0.82	1.47	0.20	0.78
O3	conventional	5.73	0.24	1.29	0.22	0.60
O3	low	5.64	0.21	0.60	0.18	0.29
C4	conventional	5.90	0.60	1.26	0.29	0.60
C4	low	6.04	0.16	1.41	0.10	0.90
S4	conventional	7.57	0.41	0.75	0.12	0.67
S4	low	7.33	0.56	0.74	0.19	0.58
O4	conventional	6.05	0.18	1.01	0.16	0.66
O4	low	6.29	0.14	0.92	0.13	0.70
A4	conventional	3.06	0.67	0.80	0.35	0.21
A4	low	1.97	0.43	0.50	0.20	0.23

*Note:* C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation, S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, O4: oat in the 4-year rotation, and A4 alfalfa in the 4-year rotation.




Figure 5: Estimation of fecundity using aboveground mass. The abbreviations on the right-hand-side y-axis are crop identities, which are the combinations of the first letter in crop species names and the rotation to which the crops belonged (C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation, S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation, S4: soybean in the 4-year rotation, O3: oat in the 3-year rotation, O4: oat in the 4-year rotation, and A4: alfalfa in the 4-year rotation). The black and red dots are values from training and testing sets, respectively. Each regression line was plotted for one crop identity by herbicide treatment using the training set.  $R^2$  values were calculated from the training set only, and thus, were slightly different from those in Table 9. Biomass index =  $\ln(\text{gram biomass} + 0.005)$  and Fecundity index =  $\ln(\text{seeds} + 1)$ .

## Discussion

Results of this study indicate that waterhemp was affected by crops and crop management in multiple ways, including a reduction in individual biomass and fecundity to the point of non-existence as occurred in 2019 soybean plots. Despite the 2018 and 2019 data being overdispersed, which resulted in high residual deviance, the significance of treatment effects was consistent. Crop identity was the most influential factor for all responses. Some covariation relationships were observed: population stand density affected sex ratio and female aboveground mass was a reliable predictor for fecundity.

Waterhemp is a small-seeded species that is more sensitive to environmental stress than larger-seeded species (Harbur and Owen, 2004). In the present study, the number of stress and mortality factors likely increased as crop diversity increased temporally and spatially (Martin and Felton, 1993). Stress and mortality factors arose from the strategic cropping system designs that employ crops of different phenology, management requirements, and relative competitiveness with weeds (Liebman and Janke, 1990; Liebman and Dyck, 1993).

The two summer annual row crops in our study, corn and soybean, differed from the cool season and perennial crops, oat, red clover, and alfalfa with regard to the strongest selection pressure against weeds: herbicides. In corn (C2, C3, and C4), weeds were controlled with broadcast herbicide (conventional), or a combination of banded herbicide (38-cm strips on top of crop rows) and interrow cultivation. In soybean (S2, S3, and S4), weeds were controlled with broadcast herbicide as in conventional corn, with different active ingredients. In contrast, in the O3, O4, and A4 treatments, no herbicide or cultivation was applied, but those three crops were strategically introduced to the 3-year and 4-year rotations for their potential allelopathic and shading effects (Liebman and Dyck, 1993; Singh et al., 2003). The spring establishment of O3 and O4 and overwintering of A4 treatments gave the crops a headstart for resource competition against waterhemp, a summer annual weed that emerged later (Hartzler et al., 2004). The timing of oat harvest in late July matched waterhemp's early reproductive stage (Buhler et al., 2008; Horak and Loughin, 2000) and the resulting mechanical damage at this stage reduced the weed's reproductive potential. Intercropping oat and alfalfa can produce stronger weed suppression than might be achieved by each species grown as a sole crop (Lanini et al., 1992), whereas the effects of intercropping oat with red clover can be more variable (Samson et al., 1990). For established alfalfa in the 4-year rotation, three to four hay cuts per crop season also served as a significant means of physical control and to reduce waterhemp reproductive potential.

High waterhemp population stand densities in oat resulted from highly productive plants in the preceding corn and soybean phases of the rotation and signaled abundant replenishment of soil seedbanks. Dyke and Barnard (1976) found that a clover and cereal intercrop substantially reduced weed emergence whereas Heggenstaller et al. (2006) found that a triticale (*x Triticosecale* Wittmack) and red clover intercrop increased weed seedling recruitment. Taking these findings with the present study's observation that higher waterhemp population stand densities and lower waterhemp aboveground mass were found in small grain and forage crops than in row crops, it is possible that cold-tolerant crops can be used to stimulate and induce fatal germination to deplete the soil seedbank (Davis and Liebman, 2003; Gallandt et al., 2005). Eventually, as more mortality and stress factors are imposed on emerged weeds via various control methods, such as allelopathy and mechanical damage via crop harvest, those emerged plants might be expected to contribute fewer seeds to the soil seedbank.

Waterhemp populations in three of the treatments, O3, O4, and A4, were slightly female-biased. Waterhemp populations in other treatments were even in sex ratio, which might be attributed to a more stressful conditions in small grain and forage crops than in row crops. A larger data set might help reducing the variance in sex ratio and provide a clearer understanding of the effect of corn weed management program on waterhemp sex ratio in subsequent oat and alfalfa phases. Systematic analysis is needed to identify the contribution of each stressor on waterhemp development and population dynamics. The 2019 sex ratio data were imputed without 2018 input but returned consistent conclusions on treatment effects, as compared to 2018. This consistency suggests an acceptable precision of the analysis model and the imputation algorithm. Since `pmm` seeks to fill in missing values with placeholders without changing the overall mean, it is reasonable to assume that the sex ratios in soybean eu's were even. The high herbicide efficacy in soybean was the strongest selection pressure on the exposed waterhemp populations.

Our analysis indicated that female aboveground mass could be used to predict fecundity parsimoniously.

The strong evidence of the significant interaction effect of weed management regime and crop identity on waterhemp fecundity justified the use of separate equations for each treatment. Since different sources of stress were introduced in the small grain and forages than in row crops, we attributed female-biasedness and lower fecundity in forages than in row crops to female herbaceous plants outperforming males under abiotic and biotic stresses (Juvany and Munné-Bosch, 2015). The stand density and sex ratio data in this study does not provide sufficient information to establish a relationship between them, as was established between individual female biomass and fecundity. It would be helpful to explore the population stand density and sex ratio relationship with a bigger data set.

In the present study, using mother plant reproduction potential (aboveground mass) gives a rough estimate of the number of seeds being added to the soil seedbank. Additionally, the total number of seeds produced at the end of the season in each treatment depended on the parent plant density and population sex ratio. The possibility of post-harvest seed loss due to seed predators under different ground cover conditions adds to the complexity of seedbank dynamics. Red clover that remained after oat harvest and alfalfa living mulch may enhance granivore activities (Davis and Liebman, 2003; Gallandt et al., 2005). Heggenstaller et al. (2006) found increased predation of velvetleaf (*Abutilon theophrasti* Medik) and giant foxtail (*Setaria faberi* Herrm) seeds in more diverse cropping systems than in shorter corn-soybean rotations. Overwintering crops such as alfalfa delayed pigweed (*Amaranthus quitensis* H.B.K.) emergence (Huarte and Arnold, 2003) and can exude allelochemicals for weed suppression (Miller, 1996). Compared to the bare ground after corn and soybean production, the post-harvest environment in oat and alfalfa may induce more seed loss due to predation (Gallandt et al., 2005). Waterhemp was not included in the Heggenstaller et al. (2006) study, but waterhemp seeds are preferred over other species' seeds by field crickets and ground beetles (van der Laet et al., 2015) so it is likely that the small grain and forage crops in the present study enhanced waterhemp seed predation.

The key take-away messages from this present study are that: 1) waterhemp reproductive potential can be effectively suppressed without herbicide with the integration of cool season crops into the cropping system, and 2) female waterhemp plants may have higher survival rate than males under the same conditions, which was reflected by higher female : male ratios.

More investigation is needed to determine how soil seedbank dynamics contribute to population dynamics in different scenarios, such as how female-biasedness could be potentially helpful to replenish a seedbank, whether sexual unevenness in a generation causes sexual unevenness in freshly produced seeds, and how those biases contribute to long-term population changes and competitiveness against crops. In the near-total control situation as occurred in the 2019 soybean plots, as the number of fresh seeds added to the soil seedbank can be considered negligible, other factors, such as sex ratio, population stand density, and plant size are of less practical concern. A more practical investigation would be to see how different levels of control efficacy translate into medium- and long-term population changes, because no herbicide is totally invulnerable to the evolution of resistance in weed populations. It would also be useful to see how populations would change once resistance occurred and how various control methods might contribute to resistance management.

## Appendix

### Appendix A:

#### *Seed cleaning and counting procedure*

The following protocol was used to separate seeds from stems and to clean seeds.

Step 1 - A whole sample was run through a brushing machine (LA-H, Westrup, Slageise, Denmark) to break the fruits from the stem. Large samples were chopped before running through the machine.

Step 2 - Chaff-covered seeds were run through a stack of two sieves (US Standard Sieve Series), one that was 710 micron on top and one that was 297 micron at the bottom in a shaker (Dayton, Seedburo Equipment Company). The sieve stack was securely closed with a top and a bottom lid. The 297-micron sieve was used to separate small stems, while the 710-micron sieve separated the dust from the seeds. Seeds at this step were still chaffy and partially enclosed in the fruits. Seeds that were still enclosed at this step did not pass the top sieve and were rubbed with rubber bands (Step 4). Small pieces of stems and dust were discarded. The material in between the two sieves was retained.

Step 3 - The material between the two sieves from step 2 was run through a custom-made dual-slope separator (Iowa State University Seed Laboratory) to separate clean seeds and chaffed seeds.

Step 4 - The pericarps from step 3 were rubbed in between rubber bands to remove capsules.

Step 5 - Seeds from steps 3 and 4 were run separately through a blower (CB-1, Agriculex Inc, Guelph, Canada) to remove the remaining capsules. The blower was started at the lowest wind speed and adjusted throughout the blowing process so that as much of the chaff and foreign materials were removed as possible while seeds were retained.

Step 6 - Steps 3 to 5 were repeated depending on the cleanliness of the material.

Step 7 - Cleaned seeds were enumerated with either an Old Mill Counter (850-3, International Marketing and Design Corp) or a Ball Coleman optical counter (Gen3, Ball Horticultural Company, Illinois, USA). The Ball Coleman optical counter was with a precision of 0.16% difference between samples several times per sample. With the Old Mill Counter, seeds were counted once. With the Old Mill Counter or by hand, each sample was counted once. With the Ball Coleman optical counter, each sample was counted several times and three counts that were the least different from one another were averaged and used as the sample's seed count.

#### *Seed sample preparation and machine calibration*

Seeds to be counted were threshed by hand or machine, depending on plant size, and then cleaned of chaff. Samples of hundreds of seeds were counted by hand, larger samples with either an Old Mill Counter or a Ball Coleman optical Counter. Before counting, each sample bag was weighed, i.e., bag, tag, and plant material, then the bag and tag. Plant weight was obtained by subtracting the bag and tag weight from the sample bag weight. Seeds must be fairly clean in order to be counted with machines. Obtaining clean seed samples without chaff and foreign materials was extremely time-consuming, so we trained an optical counter to eliminate particles whose sizes were noticeably different from the mean seed size of roughly 1 mm diameter (Bell and Tranel, 2010). The Ball Coleman optical counter was calibrated to count waterhemp seeds by running a "known sample" of 1200 seeds (a sample that was counted by hand) to establish a learning function to eliminate particles that were very much smaller and larger than the mean seed size from the final count. The "known sample" was run through the optical counter at 2300 seeds/second, uncalibrated, to obtain a distribution of all the counted particles. The counting rate of 2300 seeds/second was automated by the counter after running a small batch of seeds in such a way that a steady, single-file seed flow was maintained during the counting process. Two tails of the distribution of particle sizes, i.e., too large and too small, were trimmed in such a way that the new counts were closest to the hand-counted number. In our case, any counted particles whose surface area was larger than  $0.1963 \text{ mm}^2$  were removed from the total count in the learning function. Once the learning function was established, seeds were counted with reference to the saved settings. The known sample was recounted with the calibrated counter to confirm the result.

## Appendix B: 2019 sex ratio imputation

### Data set and imputation procedure overview

The experiment design was randomized complete block design with four blocks, nine levels of main plot (Crop ID) and two levels of split-plot (Corn weed management) effects. There were 72 experimental units (eu), 8 observational units (quadrats) per eu. Within each quadrat, six cohorts of plants were sexed. By the time that the plants were sexed, a few plants' sexes were not observable, and thus, they were marked as Unknown in the data set. This caused  $\text{Total} > \text{Male} + \text{Female}$  in some eu's.

Complete case analysis, in which missing observations are removed from the data set, is acceptable when less than 5% of the data is missing at random because removing the incomplete cases does not significantly change the outcome (Azur et al., 2011). The 2019 sex ratio is imputed because 22% of the data (755 NAs out of 3456 entries (6 cohorts/quadrat x 8 quadrats/eu x 72 eu)) was missing at random (MAR). We considered our data MAR even though the number of missing values were higher in soybean plots than in the other three crops because 1) the quadrats were placed randomly in the field before weed emergence, 2) waterhemp and other weed seedlings were present in the quadrats at the beginning of the season, and 3) waterhemp and other weed species were found in soybean plots by the time we sexed the waterhemp plants, but not in the eight fixed quadrats.

The MICE imputation method assumes that after controlling for all the observations in the original data set, all the missingness occurs at random (van Buuren and Groothuis-Oudshoorn, 2011). A good imputation model is numerically recognized by low fraction of information missing due to nonresponse ( $f_{mi}$ ), small proportion of total variance that is attributable to the missing data ( $\lambda$ ) values; and visually recognized by the similarity in kernel density estimates and data points distribution between the observed and imputed data sets (van Buuren and Groothuis-Oudshoorn, 2011). The number of imputations ( $m$ ) should be chosen such that the loss of efficiency,  $le = \frac{f_{mi}}{m} \leq 0.05$  (White et al., 2011). To give the imputation model more information, the data was imputed using 3456 data points. The analysis model on the imputed data sets was run using 72 data points.

The recommended specifications of the imputation model for this data set are: 1) uses at least 24 imputations (24 was selected because it is divisible by 4 cores in the computer processor), 2) includes all the covariates (Number of emerged seedlings, Male, Female, and Total) and predictors in the analysis model, and 3) uses an overdispersed Poisson regression model (Azur et al., 2011; White et al., 2011; and Nguyen et al., 2017). While the first two specifications were met, overdispersed Poisson regression model could not be specified. The extensions for count data mentioned in Chapter 7 (van Buuren, 2018) do not fit this data set well. Both `micemd's mice.impute.2l.glm.pois` and `mice.impute.2l.2stage.pois` functions impute the numbers of Male, Female, and Total separately (Audigier and Resche-Rigon, 2021). The predictive mean matching (`pmm`) method in the `mice` package can handle counts (van Buuren and Groothuis-Oudshoorn, 2011) and is the optimal solution for this data set at this writing (Figure 6).

### Diagnosis of the imputation model with $m = 24$

The imputation code with predicted mean matching (`pmm`) method is provided in (Nguyen and Liebman, 2022). Each round of imputation is distinguished by the number of imputations ( $m$ ). The loss of efficiency ( $le$ ) are all below the recommended value of 0.05 (White et al., 2011) for the analysis model terms using the imputed data sets, under three  $m$  values and the imputation performance improved as  $m$  increased (Table 10).  $m$  was capped at 2400 imputations for this manuscript because of limited computational power. The diagnosis of data imputation is demonstrated here with  $m = 24$  for ease of view. The distribution of the imputed data sets match that of the observation (Figure 6) so the outputs (imputed data sets) were used for further regression analyses that involved sex ratio.

### Diagnosis of the analysis model with $m = 24$

The similarity in the magnitude of reduction of deviance and dispersion values suggests comparable, small impacts of population aboveground mass and population stand density covariates on the improvement of the goodness of fit for comparing sex ratio using 2019 imputed data (Nguyen and Liebman, 2022). In addition,



496 using either of them as a covariate in the analysis models resulted in nonestimable corn weed management  
 497 effects on sex ratios in some treatments, so 2019 sex ratios were averaged over corn weed management regimes  
 498 and compared without any covariates for simplicity.




Figure 6: Number of female (A) and male (B) against the missingness probability for observed (data set numbered 0) and imputed data (data set numbered 1 to 24).

Table 10: Imputation model indices with different values of  $m$ . NAs resulted from nonestimable number of male and female in a treatment. The abbreviations are crop identities, which are the combinations of the first letter in crop species names and the rotation to which the crops belonged.

Term	n/m			riv			fmi			$\lambda$			le		
	m = 24	m = 240	m = 2400	m = 24	m = 240	m = 2400	m = 24	m = 240	m = 2400	m = 24	m = 240	m = 2400	m = 24	m = 240	m = 2400
Intercept	0.2500	0.1167	0.1233	0.3157	0.3420	0.3455	0.2461	0.2577	0.2592	0.2399	0.2549	0.2568	0.0103	0.0011	0.0001
Block: 2	0.2500	0.1167	0.1233	0.4452	0.5239	0.6026	0.3160	0.3468	0.3785	0.3081	0.3438	0.3760	0.0132	0.0014	0.0002
Block: 3	0.2500	0.1167	0.1233	0.7269	0.4931	0.5186	0.4319	0.3333	0.3440	0.4209	0.3303	0.3415	0.0180	0.0014	0.0001
Block: 4	0.2500	0.1167	0.1233	0.6642	0.6084	0.6354	0.4095	0.3814	0.3910	0.3991	0.3782	0.3885	0.0171	0.0016	0.0002
Crop ID: C3	0.2500	0.1167	0.1233	0.7145	0.4141	0.4151	0.4276	0.2957	0.2958	0.4168	0.2928	0.2934	0.0178	0.0012	0.0001
Crop ID: C2	0.2500	0.1167	0.1233	0.1766	0.2405	0.2562	0.1541	0.1965	0.2064	0.1501	0.1938	0.2039	0.0064	0.0008	0.0001
Crop ID: C4	0.2500	0.1167	0.1233	0.9428	0.9522	0.8892	0.4977	0.4912	0.4732	0.4853	0.4878	0.4707	0.0207	0.0020	0.0002
Crop ID: O3	0.2500	0.1167	0.1233	0.5597	0.6754	0.6785	0.3682	0.4063	0.4067	0.3588	0.4031	0.4042	0.0153	0.0017	0.0002
Crop ID: A4	0.2500	0.1167	0.1233	0.6102	0.5405	0.5163	0.3889	0.3539	0.3430	0.3790	0.3508	0.3405	0.0162	0.0015	0.0001
Crop ID: S2	0.2500	0.1167	0.1233	1.0551	0.0011	0.0008	0.5264	0.0036	0.0032	0.5134	0.0011	0.0008	0.0219	0.0000	0.0000
Crop ID: S3	0.2500	0.1167	0.1233	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Crop ID: S4	0.2500	0.1167	0.1233	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Corn weed management: low	0.2500	0.1167	0.1233	0.6332	0.4286	0.4564	0.3979	0.3030	0.3159	0.3877	0.3000	0.3134	0.0166	0.0013	0.0001
Crop ID: C3 x Corn weed management: low	0.2500	0.1167	0.1233	0.7540	0.6173	0.6113	0.4411	0.3848	0.3819	0.4299	0.3817	0.3794	0.0184	0.0016	0.0002
Crop ID: C2 x Corn weed management: low	0.2500	0.1167	0.1233	0.4592	0.6287	0.7774	0.3229	0.3892	0.4399	0.3147	0.3860	0.4374	0.0135	0.0016	0.0002
Crop ID: C4 x Corn weed management: low	0.2500	0.1167	0.1233	0.8062	1.0065	1.0187	0.4580	0.5051	0.5071	0.4464	0.5016	0.5046	0.0191	0.0021	0.0002
Crop ID: O3 x Corn weed management: low	0.2500	0.1167	0.1233	0.4880	0.8824	0.7931	0.3365	0.4721	0.4448	0.3279	0.4688	0.4423	0.0140	0.0020	0.0002
Crop ID: A4 x Corn weed management: low	0.2500	0.1167	0.1233	0.4903	0.4757	0.5206	0.3375	0.3253	0.3448	0.3290	0.3223	0.3424	0.0141	0.0014	0.0001
Crop ID: S2 x Corn weed management: low	0.2500	0.1167	0.1233	0.0005	NA	NA	0.0029	NA	NA	0.0005	NA	NA	0.0001	NA	NA
Crop ID: S3 x Corn weed management: low	0.2500	0.1167	0.1233	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Crop ID: S4 x Corn weed management: low	0.2500	0.1167	0.1233	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

*Note:* Some zeros are due to rounding.

C2: corn in the 2-year rotation, C3: corn in the 3-year rotation, C4: corn in the 4-year rotation,

S2: soybean in the 2-year rotation, S3: soybean in the 3-year rotation, S4: soybean in the 4-year rotation,

O3: oat in the 3-year rotation, and A4: alfalfa in the 4-year rotation

Table 11: Numerical diagnosis of model's goodness of fit with and without covariates on 2019 imputed data sets.

(A) Population aboveground mass covariate																								
Imputation	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
Deviance	53.36	48.78	25.84	40.79	59.29	40.60	48.74	56.13	67.98	59.55	51.63	61.06	40.51	53.78	36.77	69.46	38.86	52.29	55.78	55.22	53.39	54.87	49.60	44.27
Dispersion	2.26	2.19	1.03	1.66	2.33	1.82	2.04	2.43	2.71	2.49	2.15	2.69	1.69	1.95	1.58	2.55	1.64	2.09	2.36	2.10	2.17	2.20	2.16	1.57
(B) Population stand density covariate																								
Imputation	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
Deviance	64.93	58.04	21.79	47.97	56.59	37.66	40.89	44.84	68.59	52.19	43.76	52.37	44.22	48.77	40.99	79.72	47.38	43.15	54.94	66.78	43.40	70.05	48.64	48.00
Dispersion	2.74	2.65	0.87	1.97	2.24	1.69	1.73	1.93	2.75	2.16	1.83	2.72	1.89	1.77	1.78	2.94	2.09	1.72	2.33	2.58	1.80	2.90	2.12	1.73
(C) No covariate																								
Imputation	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
Deviance	82.53	90.04	48.64	66.97	82.92	71.13	67.53	76.14	104.13	81.80	73.34	102.84	72.58	98.01	58.88	99.03	74.26	92.87	83.26	81.90	83.28	97.94	71.14	72.35
Dispersion	2.12	2.47	1.21	1.75	2.07	1.88	1.75	2.05	2.74	2.17	1.94	2.73	1.84	2.37	1.56	2.33	1.92	2.33	2.17	1.99	2.07	2.60	1.88	1.73

*Note:* Bold columns are imputations that produced full sets of data (six among 24 imputed sets) that allow estimation of the effects of corn weed management on population sex ratio. In these data sets, at least one non-zero value was filled into the Male or Female column that were originally NAs. NA values in the original data set were due to high efficacy of weed control in soybean plots that left no waterhemp plants survived at sex evaluation, which resulted in non-estimable sex ratio.

## Acknowledgements

The authors thank Matt Woods, Mike Fiscus, and the Iowa State University’s Agronomy Research Farm crew for field management; Jessica Juarez-Morales, Elizabeth Oys, Ana Poznanski, Andrew Riehl, Angela Soto-Saenz, and Wyatt Westfall for field and laboratory assistance; Alan Gaul and Iowa State University’s Seed Laboratory student assistants and interns for seed cleaning laboratory space, guidance, and technical assistance; Philip Dixon and Dean Adams for data analysis assistance; Russ Lenth and other Stackoverflow community members for answering HTXN’s coding questions; Lisa Schulte-Moore, Micheal Owen, and Mark Gleason for reviewing the manuscript; Overleaf staff for L<sup>A</sup>T<sub>E</sub>X assistance in compiling the manuscript; two anonymous reviewers and the subject editor for their valuable comments.

## References

- Vincent Audigier and Matthieu Resche-Rigon. Micemd: Multiple imputation by chained equations with multilevel data, October 2021.
- Melissa J. Azur, Elizabeth A. Stuart, Constantine Frangakis, and Philip J. Leaf. Multiple imputation by chained equations: What is it and how does it work?: Multiple imputation by chained equations. *Int. J. Methods Psychiatr. Res.*, 20(1):40–49, March 2011. ISSN 10498931. doi: 10/cfbxfz.
- Michael S. Bell and Patrick J. Tranel. Time requirement from pollination to seed maturity in waterhemp (*Amaranthus tuberculatus*). *Weed Science*, 58(2):167–173, June 2010. ISSN 0043-1745, 1550-2759. doi: 10/c547b8.
- Douglas D. Buhler and Robert G. Hartzler. Emergence and persistence of seed of velvetleaf, common waterhemp, woolly cupgrass, and giant foxtail. *Weed Science*, 49(2):230–235, April 2001. ISSN 0043-1745, 1550-2759. doi: 10/dmmt6f.
- Douglas D. Buhler, Robert G. Hartzler, and Frank Forcella. Implications of weed seedbank dynamics to weed management. *Weed Science*, 45(3):329–336, 1997. ISSN 0043-1745.
- Douglas D. Buhler, Robert G. Hartzler, Frank Forcella, and Jeffrey Gunsolus. Relative emergence sequence for weeds of corn and soybeans, 2008.
- Angelo Canty and Brian Ripley. Boot: Bootstrap functions (Originally by Angelo Canty for S), May 2021.
- Mihai Costea, Susan E Weaver, and François J. Tardif. The biology of invasive alien plants in Canada. 3. *Amaranthus tuberculatus* (Moq.) Sauer var. *rudis* (Sauer) Costea & Tardif. *Can. J. Plant Sci.*, 85(2): 507–522, April 2005. ISSN 0008-4220. doi: 10/b75t54.
- Michael J. Crawley. Proportion data. In *The R Book*, pages 628–649. Wiley, second edition, 2013. ISBN 978-0-470-97392-9.
- Adam S. Davis. Weed seed pools concurrent with corn and soybean harvest in Illinois. *Weed Science*, 56(4): 503–508, August 2008. ISSN 0043-1745, 1550-2759. doi: 10/bmncpf.
- Adam S. Davis and Matt Liebman. Cropping system effects on giant foxtail (*Setaria faberi*) demography: I. Green manure and tillage timing. *Weed Science*, 51(6):919–929, December 2003. ISSN 0043-1745, 1550-2759. doi: 10/bxq7q8.
- Adam S. Davis, Jason D. Hill, Craig A. Chase, Ann M. Johanns, and Matt Liebman. Increasing cropping system diversity balances productivity, profitability and environmental health. *PLOS ONE*, 7(10):e47149, October 2012. ISSN 1932-6203. doi: 10/f39nr2.
- Philip Dixon. Should blocks be fixed or random? *Conference on Applied Statistics in Agriculture*, May 2016. ISSN 2475-7772. doi: 10/gjggh8.
- G. V. Dyke and A. J. Barnard. Suppression of couch grass by Italian ryegrass and broad red clover undersown in barley and field beans. *J. Agric. Sci.*, 87(1):123–126, August 1976. ISSN 0021-8596, 1469-5146. doi: 10/dg8cjk.

- David L. Field, Melinda Pickup, and Spencer C. H. Barrett. Comparative analyses of sex-ratio variation in dioecious flowering plants. *Evolution*, 67(3):661–672, March 2013. ISSN 1558-5646. doi: 10/f4qsnr.
- Eric R. Gallandt, Thomas Molloy, Ryan P. Lynch, and Francis A. Drummond. Effect of cover-cropping systems on invertebrate seed predation. *Weed Science*, 53(1):69–76, February 2005. ISSN 0043-1745, 1550-2759. doi: 10/bj4v3z.
- Katherine Goode and Kathleen Rey. ggResidpanel: Panels and interactive versions of diagnostic plots using 'ggplot2', May 2019.
- William F. Grant. Cytogenetic studies in *Amaranthus*: I. Cytological aspects of sex determination in dioecious species. *Can. J. Bot.*, 37(3):413–417, May 1959. ISSN 0008-4026. doi: 10/cvdprn.
- Aaron G. Hager, Loyd M. Wax, Edward W. Stoller, and Germán A. Bollero. Common waterhemp (*Amaranthus rudis*) interference in soybean. *Weed Science*, 50(5):607–610, September 2002. ISSN 0043-1745, 1550-2759. doi: 10/fp5qsm.
- Matthew M. Harbur and Micheal D. K. Owen. Light and growth rate effects on crop and weed responses to nitrogen. *Weed Science*, 52(4):578–583, August 2004. ISSN 0043-1745, 1550-2759. doi: 10/cnnw78.
- Robert G. Hartzler, Bruce A. Battles, and Dawn Nordby. Effect of common waterhemp (*Amaranthus rudis*) emergence date on growth and fecundity in soybean. *Weed Science*, 52(2):242–245, April 2004. ISSN 0043-1745, 1550-2759. doi: 10/cmhpzk.
- Heap. Herbicide resistant tall waterhemp globally (*Amaranthus tuberculatus* (=A. rudis)). <http://www.weedscience.org/Summary/Species.aspx>, 2021.
- Andrew H. Heggenstaller, Fabian D. Menalled, Matt Liebman, and Paula R. Westerman. Seasonal patterns in post-dispersal seed predation of *Abutilon theophrasti* and *Setaria faberi* in three cropping systems. *Journal of Applied Ecology*, 43(5):999–1010, October 2006. ISSN 1365-2664. doi: 10/bk6ccs.
- Joseph M. Heneghan and William G. Johnson. The growth and development of five waterhemp (*Amaranthus tuberculatus*) populations in a common garden. *Weed Science*, 65(2):247–255, March 2017. ISSN 1550-2759. doi: 10/f93hz9.
- Michael J. Horak and Thomas M. Loughin. Growth analysis of four *Amaranthus* species. *Weed Science*, 48(3):347–355, 2000. ISSN 0043-1745. doi: 10/dfvnm.
- H. R. Huarte and R. L. Benech Arnold. Understanding mechanisms of reduced annual weed emergence in alfalfa. *Weed Science*, 51(6):876–885, December 2003. ISSN 0043-1745, 1550-2759. doi: 10/fd3d6g.
- Natalie D. Hunt, Matt Liebman, Sumil K. Thakrar, and Jason D. Hill. Fossil energy use, climate change impacts, and air quality-related human health damages of conventional and diversified cropping systems in Iowa, USA. *Environ. Sci. Technol.*, 54(18):11002–11014, September 2020. ISSN 0013-936X. doi: 10/gkc28b.
- William G. Johnson, Vince M. Davis, Greg R. Kruger, and Stephen C. Weller. Influence of glyphosate-resistant cropping systems on weed species shifts and glyphosate-resistant weed populations. *European Journal of Agronomy*, 31(3):162–172, October 2009. ISSN 1161-0301. doi: 10/dxmb34.
- Marta Juvany and Sergi Munné-Bosch. Sex-related differences in stress tolerance in dioecious plants: A critical appraisal in a physiological context. *EXBOTJ*, 66(20):6083–6092, October 2015. ISSN 0022-0957, 1460-2431. doi: 10/f73m7x.
- Nicholas E. Korres and Jason K. Norsworthy. Palmer amaranth (*Amaranthus palmeri*) demographic and biological characteristics in wide-row soybean. *Weed Science*, 65(4):491–503, July 2017. ISSN 1550-2759. doi: 10/gbshrk.
- W. Thomas Lanini, Steve B. Orloff, Ronald N. Vargas, and Jack P. Orr. Fight weeds and increase forage: Using oats as a companion crop in establishing alfalfa. *California Agriculture*, 46(4):25–27, July 1992. ISSN 0008-0845. doi: 10/ghknp2.

- Russell V. Lenth, Paul Buerkner, Maxime Herve, Jonathon Love, Hannes Riebl, and Henrik Singmann. Emmeans: Estimated marginal means, aka least-squares means, November 2021.
- Matt Liebman and Elizabeth Dyck. Crop rotation and intercropping strategies for weed management. *Ecol Appl*, 3(1):92–122, February 1993. ISSN 1051-0761. doi: 10/db45cm.
- Matt Liebman and Eric R. Gallandt. Many little hammers: Ecological management of crop-weed interactions. In Louise E. Jackson, editor, *Ecology in Agriculture*, Physiological Ecology, pages 291–343. Academic Press, 1997. ISBN 978-0-12-378260-1. doi: 10.1016/B978-012378260-1/50010-5.
- Matt Liebman and R. R. Janke. Sustainable weed management practices. *Sustainable agriculture in temperate zones.*, pages 111–143, 1990.
- Matt Liebman and Virginia A. Nichols. Cropping system redesign for improved weed management: A modeling approach illustrated with giant ragweed (*Ambrosia trifida*). *Agronomy*, 10(2):262, February 2020. doi: 10/ggrj25.
- Matt Liebman, Z. J. Miller, C. L. Williams, P. R. Westerman, P. M. Dixon, A. Heggenstaller, A. S. Davis, F. D. Menalled, and D. N. Sundberg. Fates of *Setaria faberi* and *Abutilon theophrasti* seeds in three crop rotation systems. *Weed Research*, 54(3):293–306, June 2014. ISSN 1365-3180. doi: 10/f5zjjc.
- Matt Liebman, Huong T. X. Nguyen, Matthew M. Woods, Natalie D. Hunt, and Jason D. Hill. Weed seedbank diversity and sustainability indicators for simple and more diverse cropping systems. *Weed Res*, 61(3):164–177, June 2021. ISSN 0043-1737, 1365-3180. doi: 10/ghz5bj.
- Chloe MacLaren, Jonathan Storkey, Alexander Menegat, Helen Metcalfe, and Katharina Dehnen-Schmutz. An ecological future for weed science to sustain crop production and the environment. A review. *Agron. Sustain. Dev.*, 40(4):24, August 2020. ISSN 1774-0746, 1773-0155. doi: 10/gh28rg.
- R. J. Martin and W. L. Felton. Effect of crop rotation, tillage practice, and herbicides on the population dynamics of wild oats in wheat. *Australian Journal of Experimental Agriculture*, 33(2):159–165, 1993. ISSN 0816-1089.
- Fabián D. Menalled, Matt Liebman, and Douglas D. Buhler. Impact of composted swine manure and tillage on common waterhemp (*Amaranthus rudis*) competition with soybean. *Weed Science*, 52(4):605–613, August 2004. ISSN 0043-1745, 1550-2759. doi: 10/dp4572.
- Mohsen B. Mesgaran, Maor Matzrafi, and Sara Ohadi. Sex lability and dimorphism in dioecious Palmer amaranth (*Amaranthus palmeri*). Preprint, Plant Biology, September 2019.
- Darrell A. Miller. Allelopathy in forage crop systems. *Agron.j.*, 88(6):854–859, November 1996. ISSN 0002-1962, 1435-0645. doi: 10/d3t7wn.
- Jacob S. Montgomery, Darci A. Giacomini, Detlef Weigel, and Patrick J. Tranel. Male-specific Y-chromosomal regions in waterhemp (*Amaranthus tuberculatus*) and Palmer amaranth (*Amaranthus palmeri*). *New Phytol*, 229(6):3522–3533, March 2021. ISSN 0028-646X, 1469-8137. doi: 10/gjpz5c.
- Cattram D. Nguyen, John B. Carlin, and Katherine J. Lee. Model checking in multiple imputation: An overview and case study. *Emerging Themes in Epidemiology*, 14(1):8, August 2017. ISSN 1742-7622. doi: 10/gf5jh4.
- Huong T. X. Nguyen and Matt Liebman. Data for: Impact of cropping system diversification on vegetative and reproductive characteristics of waterhemp (*A. tuberculatus*), 2022.
- Huong T. X. Nguyen and Matt Liebman. Weed community composition in simple and more diverse cropping systems. *Frontiers in Agronomy*, in review.
- Dawn E. Nordby and Robert G. Hartzler. Influence of corn on common waterhemp (*Amaranthus rudis*) growth and fecundity. *Weed Science*, 52(2):255–259, April 2004. ISSN 0043-1745, 1550-2759. doi: 10/b6hsfr.
- Roger William Payne. The design and analysis of long-term rotation experiments. *Agronomy Journal*, 107(2): 772–785, March 2015. ISSN 00021962. doi: 10/f673ct.

- José Pinhero, Douglas Bates, Saikat Debroy, Deepayan Sarkar, Siem Heisterkamp, Bert Van Willigen, Johannes Ranke, and R-core. *Nlme: Linear and nonlinear mixed effects models*, February 2021.
- Donald B. Pratt and Lynn G. Clark. *Amaranthus rudis* and *A. tuberculatus*, One species or two? *The Journal of the Torrey Botanical Society*, 128(3):282–296, 2001. ISSN 1095-5674. doi: 10/drxbzj.
- Joby M. Prince, David R. Shaw, Wade A. Givens, Micheal D. K. Owen, Stephen C. Weller, Bryan G. Young, Robert G. Wilson, and David L. Jordan. Benchmark study: IV. Survey of grower practices for managing glyphosate-resistant weed populations. *Weed Technology*, 26(3):543–548, September 2012. ISSN 0890-037X, 1550-2740. doi: 10/f37vn9.
- R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, November 2021.
- Roger Samson, Chantal Foulds, and David Patriquin. Choice and management of cover crop species and varieties for use in row crop dominant rotations. Final report, University of Guelph, 1990.
- H. P. Singh, Daizy R. Batish, and R.K. Kohli. Allelopathic interactions and allelochemicals: New possibilities for sustainable weed management. *Critical Reviews in Plant Sciences*, 22(3-4):239–311, May 2003. ISSN 0735-2689, 1549-7836. doi: 10/cns78q.
- Lawrence E. Steckel and Christy L. Sprague. Common waterhemp (*Amaranthus rudis*) interference in corn. *weeds*, 52(3):359–364, May 2004. ISSN 0043-1745, 1550-2759. doi: 10/frc847.
- Patrick J Tranel. Herbicide resistance in *Amaranthus tuberculatus*. *Pest Manag Sci*, 77(1):43–54, January 2021. ISSN 1526-498X, 1526-4998. doi: 10/gjz5w.
- Richard H. Uva, Joseph C. Neal, and Joseph M. Ditomaso. *Weeds of the Northeast*. Cornell University Press, Ithaca, NY, May 1997. ISBN 978-0-8014-8334-9.
- Stef van Buuren. *Flexible imputation of missing data*. Chapman and Hall/CRC interdisciplinary statistics series. CRC Press, Taylor & Francis Group, Boca Raton, second edition edition, 2018. ISBN 978-1-138-58831-8.
- Stef van Buuren and Karin Groothuis-Oudshoorn. **Mice** : Multivariate imputation by chained equations in R. *J. Stat. Soft.*, 45(3), 2011. ISSN 1548-7660. doi: 10/gbw279.
- Rocio van der Laat, Micheal D. K. Owen, Matt Liebman, and Ramon G. Leon. Postdispersal weed seed predation and invertebrate activity density in three tillage regimes. *Weed Science*, 63(4):828–838, December 2015. ISSN 0043-1745, 1550-2759. doi: 10/gddc93.
- David Weisberger, Virginia Nichols, and Matt Liebman. Does diversifying crop rotations suppress weeds? A meta-analysis. *PLoS ONE*, 14(7):e0219847, July 2019. ISSN 1932-6203. doi: 10/gh28qh.
- Paula R. Westerman, Matt Liebman, Fabián D. Menalled, Andrew H. Heggenstaller, Robert G. Hartzler, and Philip M. Dixon. Are many little hammers effective? Velvetleaf ( *Abutilon theophrasti*) population dynamics in two- and four-year crop rotation systems. *Weed Science*, 53(3):382–392, June 2005. ISSN 0043-1745, 1550-2759. doi: 10/bgc2x2.
- Ian R. White, Patrick Royston, and Angela M. Wood. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med*, 30(4):377–399, February 2011. ISSN 1097-0258. doi: 10/cmwhtz.
- Hadley Wickham and RStudio. Tidyverse: Easily install and load the 'tidyverse', April 2021.
- Hadley Wickham, Winston Chang, Lionel Henry, Thomas Lin Pedersen, Kohske Takahashi, Claus Wilke, Kara Woo, Hiroaki Yutani, Dewey Dunnington, and RStudio. Ggplot2: Create elegant data visualisations using the grammar of graphics, June 2021.
- Rong-Cai Yang and Patricia Juskiw. Analysis of covariance in agronomy and crop research. *Can. J. Plant Sci.*, 91(4):621–641, July 2011. ISSN 0008-4220, 1918-1833. doi: 10/ft84t7.
- Hao Zhu, Thomas Trivison, Timothy Tsai, Will Beasley, Yihui Xie, GuangChuang Yu, Stéphane Laurent, Rob Shepherd, Yoni Sidi, Brian Salzer, George Gui, Yeliang Fan, Duncan Murdoch, and Bill Evans. kableExtra: Construct complex table with 'kable' and pipe syntax, February 2021.