

Notas e Listas de Prob 2

Henrique

August 14, 2026

Contents

0	Informações importantes	1
1	Martingales	1
2	Listas	3
A	Resultados clássicos de Convergência	3

0 Informações importantes

Devo guardar neste arquivo notas de aula e listas do curso. Sobre as notas, sua maior parte será somente uma enunciação dos principais teoremas vistos, provas interessantes podem ser comentadas e rascunhadas. O código fonte desse arquivo pode ser encontrado em https://en.wikipedia.org/wiki/The_Garden_of_Earthly_Delights.

1 Martingales

Definition 1.1. Let $(\Omega, \mathcal{F}, P)$ be a probability space, a martingale is a sequence $X_1, X_2, \dots$ of random variables associated with σ -algebras $\mathcal{F}_1, \mathcal{F}_2, \dots$ that satisfy:

- (i) $X_i \in L^1$ and $X_i \in \mathcal{F}_i$.
- (ii) $\mathcal{F}_i \subset \mathcal{F}_{i+1}$.
- (iii) $\mathbb{E}[X_{i+1} | \mathcal{F}_i] = X_i$.

Remark 1.2. If one changes the assumption of (iii) to be $\mathbb{E}[X_{i+1} | \mathcal{F}] \leq X_i$ (resp. $\geq$), we call such sequence a supermartingale (resp. submartingale).

Theorem 1.3. (Doob's inequality) Suppose $\{X_j\}$ is a martingale. Then

$$\mathbb{P}(\sup_{j \leq n} |X_j| \geq l) \leq \frac{1}{l} \int \mathbf{1}[\sup_{j \leq n} |X_j| \geq l] |X_n| dP \leq \frac{1}{l} \mathbb{E}|X_n|.$$

Proof. The proof is similar to Kolmogorov's maximal inequality. For $1 \leq j \leq n$, let E_j be the event $\{X_i < l, i < j\} \cap \{X_j \geq l\}$, that is the event that X_j is the first to surpass l . Note that

$$\mathbb{P}(\sup_{j \leq n} |X_j| \geq l) = \sum_{j \leq n} \mathbb{E}[\mathbf{1}[E_j]] \leq \sum_{j \leq n} \frac{1}{l} \mathbb{E}[\mathbf{1}[E_j] |X_j|] \leq \sum_{j \leq n} \frac{1}{l} \mathbb{E}[\mathbf{1}[E_j] |X_n|] = \mathbb{E}[\mathbf{1}[\sup_{j \leq n} |X_j| \geq l] |X_n|] \quad (1)$$

where the second inequality follows from the fact that $\mathbb{E}[X_n | \mathcal{F}_j] = X_j$ and $|x|$ is a convex function. $\square$

Corollary 1.4. (Doob's p-inequality) Let $\{X_j\}$ be a martingale and $p > 1$. Then if, as before, $S_n = \sup_{j \leq n} |X_j|$, we have

$$\mathbb{E}S_n^p \leq \left(\frac{p}{p-1}\right)^p \mathbb{E}|X_n|^p.$$

Proof. Follows from the previous theorem and an integration by parts. Note that for any bounded positive r.v's Y and X that satisfy $\mathbb{P}(Y \geq l) \leq \frac{1}{l} \mathbb{E}[\mathbf{1}[Y \geq l]X]$ we have

$$\begin{aligned} \int Y^p dP &= \int \int_0^Y p y^{p-1} dy dP \\ &= p \int_0^\infty y^{p-1} \mathbb{P}(Y \geq y) dy \\ &\leq p \int_0^\infty y^{p-1} \frac{1}{y} \mathbb{E}[\mathbf{1}[Y \geq y]X] dy \\ &= p \int_0^\infty y^{p-2} \int \mathbf{1}[Y \geq y] X dP \\ &= p \int X \int_0^Y y^{p-2} dy = \frac{p}{p-1} \int XY^{p-1} \\ &\leq \frac{p}{p-1} (\mathbb{E}X^p)^{1/p} (\mathbb{E}Y^p)^{\frac{p-1}{p}} \end{aligned}$$

So that $\mathbb{E}Y^p \leq (\frac{p}{p-1})^p \mathbb{E}X^p$. We can then consider a truncation argument to prove the result for unbounded positive random variables. $\square$

Remark 1.5. This corollary is very cool on its own, it says that if some X_n in the martingale belongs to L^p , then all previous r.v's belong as well.

Exercise 1.1. Consider the martingale of doubling one's bet. In the space $([0, 1], \mathcal{B}([0, 1]), dm)$, we consider the random variables $X_n \in \sigma([j/2^n, (j+1)/2^n])$

$$X_n(\omega) = \begin{cases} 2^n & \text{if } \omega \in [0, 2^{-n}] \\ 0 & \text{elsewhere} \end{cases}$$

It is easy to see that they define a martingale, all in L^1 , $\mathbb{E}X_n = 1$, but $\mathbb{E} \sup_{j \leq n} X_j = (n+1)/2$. So no alternative to [1.4] exists in L^1 .

2 Listas

A Resultados clássicos de Convergência