

# On Kernel Mengerian Orientations of Line Multigraphs

Han Xiao \*

Department of Mathematics,  
The University of Hong Kong,  
Hong Kong, China

## Abstract

We present a polyhedral description of kernels in orientations of line multigraphs. Given a digraph  $D$ , let  $FK(D)$  denote the fractional kernel polytope defined on  $D$ , and let  $\sigma(D)$  denote the linear system defining  $FK(D)$ . A digraph  $D$  is called kernel perfect if every induced subdigraph  $D'$  has a kernel, called kernel ideal if  $FK(D')$  is integral for each induced subdigraph  $D'$ , and called kernel Mengerian if  $\sigma(D')$  is TDI for each induced subdigraph  $D'$ . We show that an orientation of line multigraph is kernel perfect iff it is kernel ideal iff it is kernel Mengerian. Our result strengthens the theorem of Borodin *et al.* [3] on kernel perfect digraphs and generalizes the theorem of Király and Pap [9] on the stable matching problem.

**AMS subject classifications:** 90C10, 90C27, 90C57.

**Key words:** kernel, stable matching, line graph, polytope, total dual integrality.

---

\*hxiao.math@connect.hku.hk

## 1 Introduction

A graph is called *simple* if it contains neither loops nor parallel edges, and is called a *multigraph* if it has parallel edges. A *simple* digraph is an orientation of simple graph. A *multi-digraph* is an orientation of multigraph.

Let  $G$  be a graph. The *line graph* of  $G$ , denoted by  $L(G)$ , is a graph such that: each vertex of  $L(G)$  corresponds to an edge of  $G$ , and two vertices of  $L(G)$  are adjacent if and only if they are incident as edges in  $G$ . We call  $L(G)$  the *line multigraph* of  $G$  if any two vertices of  $L(G)$  are connected by as many edges as the number of their common ends in  $G$ . We call  $G$  a *root* of  $L(G)$ .

Let  $D = (V, A)$  be a digraph. For  $U \subseteq V$ , we call  $U$  an *independent* set of  $D$  if no two vertices in  $U$  are connected by an arc, call  $U$  a *dominating* set of  $D$  if for each vertex  $v \notin U$ , there is an arc from  $v$  to  $U$ , and call  $U$  a *kernel* of  $D$  if it is both independent and dominating. We call  $D$  *kernel perfect* if each of its induced subdigraphs has a kernel. A *clique* of  $D$  is a subset of  $V$  such that any two vertices are connected by an arc. We call  $D$  *clique-acyclic* if for each clique of  $D$  the induced subdigraph of one-way arc is acyclic, and call  $D$  *good* if it is clique-acyclic and every directed odd cycle has a (pseudo-)chord<sup>1</sup>.

**Theorem 1.1** (Borodin *et al.* [3]). *Let  $G$  be a line multigraph. The orientation  $D$  of  $G$  is kernel perfect if and only if it is good.*

A subset  $P$  of  $\mathbb{R}^n$  is called a *polytope* if it is the convex hull of finitely many vectors in  $\mathbb{R}^n$ . A point  $x$  in  $P$  is called a *vertex* or an *extreme point* if there exist no distinct points  $y$  and  $z$  in  $P$  such that  $x = \alpha y + (1 - \alpha)z$  for  $0 < \alpha < 1$ . It is well known that  $P$  is the convex hull of its vertices, and that there exists a linear system  $Ax \leq b$  such that  $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ . We say  $P$  is *1/k-integral* if its vertices are 1/k-integral vectors, where  $k \in \mathbb{N}$ . By a theorem in linear programming,  $P$  is 1/k-integral if and only if  $\max\{c^T x : Ax \leq b\}$  has an optimal 1/k-integral solution for every integral vector  $c$  for which the optimum is finite. If, instead,  $\max\{c^T x : Ax \leq b\}$  has a dual optimal 1/k-integral solution, we say  $Ax \leq b$  is *totally dual 1/k-integral* (TDI/k). It is easy to verify that  $Ax \leq b$  is TDI/k if and only if  $Bx \leq b$  is TDI, where  $B = A/k$  and  $k \in \mathbb{N}$ . Thus from a theorem of Edmonds and Giles [8], we deduce that if  $Ax \leq b$  is TDI/k and  $b$  is integral, then  $P = \{x \in \mathbb{R}^n : Ax \leq b\}$  is 1/k-integral.

---

<sup>1</sup>A pseudo-chord is an arc  $(v_i, v_{i-1})$  in a directed cycle  $v_1 v_2 \dots v_l v_1$ .

Let  $\sigma(D)$  denote the linear system consisting of the following inequalities:

$$x(v) + x(N^+(v)) \geq 1 \quad \forall v \in V, \quad (1.1)$$

$$x(Q) \leq 1 \quad \forall Q \in \mathcal{Q}, \quad (1.2)$$

$$x(v) \geq 0 \quad \forall v \in V, \quad (1.3)$$

where  $x(U) = \sum_{u \in U} x(u)$  for any  $U \subseteq V$ ,  $N^+(v)$  denotes the set of all out-neighbors of vertex  $v$ , and  $\mathcal{Q}$  denotes the set of all cliques of  $D$ . Observe that incidence vectors of kernels of  $D$  are precisely integral solutions  $x \in \mathbb{Z}^A$  to  $\sigma(D)$ . The *kernel polytope* of  $D$ , denoted by  $K(D)$ , is the convex hull of incidence vectors of all kernels of  $D$ . The *fractional kernel polytope* of  $D$ , denoted by  $FK(D)$ , is the set of all solutions  $x \in \mathbb{R}^A$  to  $\sigma(D)$ . Clearly,  $K(D) \subseteq FK(D)$ . We call  $D$  *kernel ideal* if  $FK(D')$  is integral for each induced subdigraph  $D'$ , and *kernel Mengerian* if  $\sigma(D')$  is TDI for each induced subdigraph  $D'$ .

As described in Egres Open [1], the polyhedral description of kernels remains open. Chen *et al.* [5] attained a polyhedral characterization of kernels by replacing clique constraints  $x(Q) \leq 1$  for  $Q \in \mathcal{Q}$  with independence constraints  $x(u) + x(v) \leq 1$  for  $(u, v) \in A$ . In this paper we show that kernels in orientations of line multigraph can be characterized polyhedrally.

**Theorem 1.2.** *Let  $D$  be an orientation of a line multigraph. Then the following statements are equivalent:*

- (i)  $D$  is good;
- (ii)  $D$  is kernel perfect;
- (iii)  $D$  is kernel ideal;
- (iv)  $D$  is kernel Mengerian.

The equivalence of (i) and (ii) was established by Borodin *et al.* [3] (Maffray [10] proved the case when  $D$  is perfect). Király and Pap [9] proved Theorem 1.2 for the case when the root of  $D$  is bipartite. Our result strengthens the theorem of Borodin *et al.* [3] and generalizes the theorem of Király and Pap [9] to line multigraphs.

## 2 Preliminaries

Kernels are closely related to stable matchings. Before proceeding, we introduce some notations and some theorems of the stable matching problem first. Let  $G = (V, E)$  be a graph. For  $v \in V$ , let  $\delta(v)$  denote the set of edges incident to  $v$  and  $\prec_v$  be a strict linear order on  $\delta(v)$ . We call  $\prec_v$

the *preference* of  $v$ , and for edges  $e$  and  $f$  incident to  $v$  we say  $v$  *prefers*  $e$  to  $f$  or  $e$  *dominates*  $f$  if  $e \prec_v f$ . Let  $\prec$  be the set of linear order  $\prec_v$  for  $v \in V$ . We call the pair  $(G, \prec)$  *preference system*, and call  $(G, \prec)$  *simple* if  $G$  is simple. For  $e \in E$ , let  $\varphi(e)$  denote the set consisting of  $e$  itself and edges that dominate  $e$  in  $(G, \prec)$ , and let  $\varphi_v(e)$  denote the set of edges that dominate  $e$  at vertex  $v$  in  $(G, \prec)$ . Given a matching  $M$  in  $G$ , we call  $M$  *stable* in  $(G, \prec)$  if every edge of  $G$  is either in  $M$  or is dominated by some edge in  $M$ .

Let  $\pi(G, \prec)$  denote the linear system consisting of the following linear inequalities:

$$x(\varphi(e)) \geq 1 \quad \forall e \in E, \quad (2.1)$$

$$x(\delta(v)) \leq 1 \quad \forall v \in V, \quad (2.2)$$

$$x(e) \geq 0 \quad \forall e \in E. \quad (2.3)$$

As observed by Abeledo and Rothblum [2], incidence vectors of stable matchings of  $(G, \prec)$  are precisely integral solutions  $x \in \mathbb{Z}^E$  to  $\pi(G, \prec)$ . The *stable matching polytope*, denoted by  $SM(G, \prec)$ , is the convex hull of incidence vectors of all stable matchings of  $(G, \prec)$ . The *fractional stable matching polytope*, denoted by  $FSM(G, \prec)$ , is the set of all solutions  $x \in \mathbb{R}^E$  to  $\pi(G, \prec)$ . Clearly,  $SM(G, \prec) \subseteq FSM(G, \prec)$ .

**Theorem 2.1** (Rothblum [11]). *Let  $(G, \prec)$  be a simple preference system. If  $G$  is bipartite, then  $SM(G, \prec) = FSM(G, \prec)$ .*

**Theorem 2.2** (Király and Pap [9]). *Let  $(G, \prec)$  be a simple preference system. If  $G$  is bipartite, then  $\pi(G, \prec)$  is totally dual integral.*

Given a cycle  $C = v_1 v_2 \dots v_l v_1$  in  $G$ , we call  $C$  of *cyclic preferences* in  $(G, \prec)$  if  $v_{i-1} v_i \prec_{v_i} v_i v_{i+1}$  for  $i = 1, 2, \dots, l$  or  $v_{i-1} v_i \succ_{v_i} v_i v_{i+1}$  for  $i = 1, 2, \dots, l$ , where indices are taken modulo  $l$ . For  $x \in FSM(G, \prec)$ , let  $E_\alpha(x)$  denote the set of all edges with  $x(e) = \alpha$  where  $\alpha \in \mathbb{R}$  and  $E_+(x)$  denote the set of all edges with  $x(e) > 0$ .

**Theorem 2.3** (Abeledo and Rothblum [2]). *Let  $(G, \prec)$  be a simple preference system. Then  $FSM(G, \prec)$  is  $1/2$ -integral. Moreover, for each  $1/2$ -integral point  $x$  in  $FSM(G, \prec)$ ,  $E_{1/2}(x)$  consists of vertex disjoint cycles with cyclic preferences.*

**Theorem 2.4** (Chen et al. [6]). *Let  $(G, \prec)$  be a simple preference system. Then  $\pi(G, \prec)$  is totally dual  $1/2$ -integral. Moreover,  $\pi(G, \prec)$  is totally dual integral if and only if  $SM(G, \prec) = FSM(G, \prec)$ .*

### 3 Reductions

Given a clique-acyclic orientation  $D$  of line multigraph  $L(H)$ , let  $e \prec_v f$  if  $(f, e)$  is an arc in  $D$  for any two incident edges  $e$  and  $f$  with common end  $v$  in  $H$ . Hence  $D$  is associated with a preference system  $(H, \prec)$ . Recall that  $\sigma(D)$  denotes the linear system which defines  $FK(D)$ . Consequently,  $\sigma(D)$  can be viewed as a linear system defined on preference system  $(H, \prec)$ . The equivalence of constraints (1.3) and constraints (2.3) follows directly. Constraints (1.1) can be viewed as constraints (2.1) because of the one to one correspondence between dominating vertex set  $\{v\} \cup N_D^+(v)$  for  $v \in V(D)$  and stable edge set  $\varphi(e)$  for  $e \in E(H)$ . Observe that cliques of  $D$  correspond to three types of edge set in  $H$ :

- (a)  $\delta(v)$  for  $v \in V(H)$ ,
- (b) nontrivial subsets of  $\delta(v)$  for  $v \in V(H)$ ,
- (c) complete subgraphs of  $H$  induced on three vertices,

and all three types allow parallel edges. Hence constraints (1.2) can be viewed as constraints (2.2) together with some extra constraints on  $(H, \prec)$ . Let  $\mathcal{O}(H)$  denote the set of all complete subgraphs of  $H$  induced on three vertices. Then  $\sigma(D)$  can be reformulated in terms of preference system  $(H, \prec)$ :

$$x(\varphi(e)) \geq 1 \quad \forall e \in E(H), \quad (3.1)$$

$$x(\delta(v)) \leq 1 \quad \forall v \in V(H), \quad (3.2)$$

$$x(S) \leq 1 \quad \emptyset \subset S \subset \delta(v), \quad \forall v \in V(H), \quad (3.3)$$

$$x(O) \leq 1 \quad \forall O \in \mathcal{O}(H), \quad (3.4)$$

$$x(e) \geq 0 \quad \forall e \in E(H). \quad (3.5)$$

Notice that constraints (3.1), (3.2) and (3.5) form the Rothblum system  $\pi(H, \prec)$  which defines  $FSM(H, \prec)$ . Constraints (3.3) are redundant with respect to  $\pi(H, \prec)$  due to constraints (3.2). As we shall see, constraints (3.4) are also redundant with respect to  $\pi(H, \prec)$ . Hence  $FK(D)$  is essentially defined by Rothblum system  $\pi(H, \prec)$ , or equivalently that  $FK(D) = FSM(H, \prec)$ .

Observe that  $H$  is a multigraph. To bridge the gap between simple preference system and  $(H, \prec)$ , we exploit the gadget introduced by Cechlárová and Fleiner [4]. We define a simple preference system  $(H', \prec')$  from  $(H, \prec)$  by substituting each parallel edge  $e$  with endpoints  $u$  and  $v$  in  $H$  by a 6-cycle with two hanging edges as in Figure 1 such that  $uu_0$  (*resp.*  $vv_0$ ) has the same order with  $uv$  in  $\prec_u$  (*resp.*  $\prec_v$ ). Notice that the construction preserves the parity of cycles with cyclic preferences in  $H$ .




Figure 1: The gadget introduced for parallel edges

**Lemma 3.1.**  $FSM(H, \prec)$  is a projection of  $FSM(H', \prec')$ .

*Proof.* Take  $x \in FSM(H, \prec)$ . For each parallel edge  $e$  with endpoints  $u$  and  $v$  in  $H$ , we set the value of edges in the gadget corresponding to  $e$  as follows:

- (a)  $x'_{uu_0} = x'_{vv_0} := x_e$ ;
- (b)  $x'_{u_0u_1} = x'_{v_0v_2} := 1 - x_e - x(\varphi_u(e))$ ;
- (c)  $x'_{u_0u_2} = x'_{v_0v_1} := x(\varphi_u(e))$ ;
- (d)  $x'_{u_1v_2} := x_e + x(\varphi_u(e))$ ;
- (e)  $x'_{u_2v_1} := 1 - x(\varphi_u(e))$ .

For each edge  $f$  without parallel edges in  $H$ , set  $x'_f := x_f$ . It is easy to check that  $x' \in FSM(H', \prec')$ . Hence  $FSM(H, \prec)$  is a projection of  $FSM(H', \prec')$ .  $\square$

By Theorem 2.3 and Lemma 3.1,  $FSM(H, \prec)$  is  $1/2$ -integral since  $FSM(H', \prec')$  is  $1/2$ -integral. Hence vertices in  $FSM(H, \prec)$  with  $x(O) = 3/2$  where  $O \in \mathcal{O}(H)$  are the only possible vertices of  $FSM(H, \prec)$  that violate constraints (3.4). However each  $O$  with  $x(O) = 3/2$  leads to a 3-cycle with cyclic preferences which arises from a directed 3-cycle in  $D$ , contradicting to the assumption that  $D$  is clique-acyclic. It follows that constraints (3.4) are unbinding for all vertices of  $FSM(H, \prec)$ . Hence constraints (3.4) are redundant with respect to  $\pi(H, \prec)$

**Lemma 3.2.** If  $\pi(H', \prec')$  is totally dual integral, then so is  $\pi(H, \prec)$ .

*Proof.* We show that  $\pi(H, \prec)$  can be obtained from  $\pi(H', \prec')$  after a series of Fourier-Motzkin eliminations. It suffices to demonstrate one elimination process from a gadget to an edge. Given a gadget as in Figure 1 arising from edge  $e$ , eliminate  $u_1v_2$  from  $\pi(H', \prec')$  first. Then all constraints involving  $x_{u_1v_2}$  are replaced by equality  $x_{u_0u_1} = x_{v_0v_2}$ . Similarly, eliminating  $u_2v_1$  yields equality

$x_{v_0v_1} = x_{u_0u_2}$ . Next eliminating  $uu_0$  gives  $x(\delta(u) \setminus \{uu_0\}) \leq x_{u_0u_1} + x_{u_0u_2}$  and  $x_{u_0u_2} \leq x(\varphi_u(uv))$ . After eliminating  $u_0u_1$  and  $u_0u_2$ , we arrive at  $x(\delta(u) \setminus \{uu_0\}) \leq x_{v_0v_1} + x_{v_0v_2}$  and  $x_{v_0v_1} \leq x(\varphi_u(uv))$ . In the end, canceling  $v_0v_1$  and  $v_0v_2$  gives  $x_{vv_0} + x(\varphi_v(vv_0) + x(\varphi_u(uu_0))) \geq 1$  and  $x_{vv_0} + x(\delta(u) \setminus \{uu_0\}) \leq 1$ . Besides,  $x_{vv_0} + x(\delta(v) \setminus \{vv_0\}) \leq 1$  is unchanged. Hence all constraints involving edges from the gadget are reduced to three constraints  $x_{vv_0} + x(\varphi_v(vv_0) + x(\varphi_u(uu_0))) \geq 1$ ,  $x_{vv_0} + x(\delta(u) \setminus \{uu_0\}) \leq 1$ , and  $x_{vv_0} + x(\delta(v) \setminus \{vv_0\}) \leq 1$ , which can be viewed as  $x(\varphi(e)) \geq 1$ ,  $x(\delta(u)) \leq 1$ , and  $x(\delta(v)) \leq 1$  respectively. Performing Fourier-Motzkin eliminations in such an order for each gadget in  $H'$  leads to a linear system defined on  $(H, \prec)$ , which is precisely the same with  $\pi(H, \prec)$  (renaming variables and removing redundant constraints if necessary). As proved by Cook [7], total dual integrality is preserved under Fourier-Motzkin elimination of a variable if it occurs in each constraint with coefficient 0 or  $\pm 1$ . Hence the lemma follows.  $\square$

By Lemma 3.1 and Lemma 3.2, we generalize the theorem of Chen *et al.* [6] on the stable matching problem to multigraphs.

**Corollary 3.3.** *Let  $(G, \prec)$  be a preference system, where  $G$  is a multigraph. Then  $\pi(G, \prec)$  is 1/2-totally dual integral. Moreover,  $\pi(G, \prec)$  is totally dual integral if and only if  $FSM(G, \prec) = SM(G, \prec)$ .*

Moreover, we have the following corollary for kernels in clique-acyclic orientations of line multigraph.

**Corollary 3.4.** *Let  $D$  be a clique-acyclic orientation of line multigraph. Then  $\sigma(D)$  is 1/2-totally dual integral. Moreover,  $\sigma(D)$  is totally dual integral if and only if  $FK(D) = K(D)$ .*

## 4 Proofs

Observe that when  $D$  is good, both  $(H, \prec)$  and  $(H', \prec')$  admit no odd cycles with cyclic preferences. In the following we exhibit some properties of simple preference systems admitting no odd cycles with cyclic preferences.

**Lemma 4.1.** *Let  $(G, \prec)$  be a simple preference system. If  $(G, \prec)$  admits no odd cycles with cyclic preferences, then  $SM(G, \prec) = FSM(G, \prec)$ .*

By Theorem 2.4, integrality of  $FSM(G, \prec)$  is equivalent to total dual integrality of  $\pi(G, \prec)$ , where  $(G, \prec)$  is a simple preference system. A corollary follows directly.

**Corollary 4.2.** *Let  $(G, \prec)$  be a simple preference system. If  $(G, \prec)$  admits no odd cycles with cyclic preferences, then  $\pi(G, \prec)$  is totally dual integral.*




Figure 2: Case 2

*Proof of Lemma 4.1.* By Theorem 2.3,  $FSM(G, \prec)$  is  $1/2$ -integral as  $(G, \prec)$  is a simple preference system. Let  $x$  be a  $1/2$ -integral point in  $FSM(G, \prec)$ . Since  $(G, \prec)$  admits no odd cycles with cyclic preferences,  $E_{1/2}(x)$  consists of even cycles  $C_1, C_2, \dots, C_r$  with cyclic preferences. For  $i = 1, 2, \dots, r$ , label vertices and edges of  $C_i \in E_{1/2}(x)$  such that  $C_i = v_1^i v_2^i \dots v_l^i$  and  $e_k^i \prec_{v_{k+1}^i} e_{k+1}^i$  for  $k = 1, 2, \dots, l$ , where  $e_k^i = v_k^i v_{k+1}^i$  and indices are taken modulo  $l$ . We remark that the parity of vertices and edges refers to the parity of their indices. Define  $z \in \mathbb{R}^{E(G)}$  by

$$z(e) := \begin{cases} 1 & e \text{ is an even edge in } C \in E_{1/2}(x), \\ -1 & e \text{ is an odd edge in } C \in E_{1/2}(x), \\ 0 & \text{otherwise.} \end{cases}$$

We are going to exclude  $x$  from vertices of  $FSM(G, \prec)$  by adding perturbation  $\epsilon z$  for small  $\epsilon$  to  $x$  and showing that  $x \pm \epsilon z \in FSM(G, \prec)$ . Tight constraints in (2.1)-(2.3) under perturbation  $\epsilon z$  play a key role here. Observe that tight constraints in (2.2) and (2.3) are invariant under perturbation  $\epsilon z$ . It remains to show that perturbation  $\epsilon z$  does not affect tight constraints in (2.1) either. Let  $e$  be an edge with  $x(\varphi(e)) = 1$ . Clearly,  $|\varphi(e) \cap E_+(x)| \in \{1, 2\}$ . When  $|\varphi(e) \cap E_+(x)| = 1$ ,  $x(e) = 1$  follows, which is trivial. When  $|\varphi(e) \cap E_+(x)| = 2$ , we claim that the parity of dominating edges in  $E_{1/2}(x)$  of  $e$  does not agree (relabeling vertices and edges in  $E_{1/2}(x)$  if necessary). Hence corresponding tight constraints in (2.1) are also invariant under perturbation  $\epsilon z$ . To justify this claim, we distinguish four cases.

**Case 1.** Edge  $e$  is an edge from some  $C \in E_{1/2}(x)$ . This case is trivial since  $C$  admits cyclic preferences.

**Case 2.** Edge  $e$  is a chord in some  $C \in E_{1/2}(x)$ . We first show that endpoints of  $e$  have different parity in  $C$ . We prove it by contradiction. Without loss of generality, let  $e = v_{2i} v_{2j}$ .






Figure 3: Case 4

If  $e_{2i} \prec e$ , then  $e_{2i-1} \prec e$ . Since  $x(\varphi(e)) = 1$ , it follows that  $e \prec e_{2j-1}$  and  $e \prec e_{2j}$ . However,  $v_{2i}ev_{2j}e_{2j}v_{2j+1} \dots v_{2i-1}e_{2i-1}v_{2i}$  form an odd cycle with cyclic preferences, a contradiction. Hence  $e \prec e_{2i}$ .

Similarly, if  $e_{2j} \prec e$ , then  $e_{2j-1} \prec e$ . Equality  $x(\varphi(e)) = 1$  implies that  $e \prec e_{2i}$  and  $e \prec e_{2i-1}$ . However,  $v_{2i}e_{2i}v_{2i+1} \dots v_{2j-1}e_{2j-1}v_{2j}ev_{2i}$  form an odd cycle with cyclic preferences, a contradiction. Hence  $e \prec e_{2j}$ .

Now  $e \prec e_{2i}$  and  $e \prec e_{2j}$ , it follows that  $e_{2i-1} \prec e$  and  $e_{2j-1} \prec e$  since  $x(\varphi(e)) = 1$ . But in this case two odd cycles with cyclic preferences mentioned above occur at the same time. Therefore, endpoints of  $e$  have different parity in  $C$ . Hence let  $e = v_{2i}v_{2j+1}$ . If  $e_{2i} \prec e$  (*resp.*  $e_{2j+1} \prec e$ ), it follows that  $e_{2i-1} \prec e$  (*resp.*  $e_{2j} \prec e$ ). Then  $e$  is dominated by two consecutive edges from  $C$ , which is trivial. So assume that  $e \prec e_{2i}$  and  $e \prec e_{2j+1}$ . Since  $x(\varphi(e)) = 1$ , it follows that  $e_{2i+1} \prec e$  and  $e_{2j} \prec e$ . Therefore  $e$  is dominated by two edges with different parity.

**Case 3.** Edge  $e$  is a hanging edge of some  $C \in E_{1/2}(x)$  and dominated by two edges from  $C$ . This case is trivial.

**Case 4.** Edge  $e$  is a connecting edge between  $C_i$  and  $C_j$  and dominated by one edge from  $C_i$  and one edge from  $C_j$  respectively, where  $C_i, C_j \in E_{1/2}(x)$ . For  $k = 1, 2, \dots, r$ , let  $F_k$  be the subset of edges in this case and incident to  $C_k$ . Then  $\cup_{i=1}^r F_i \cup C_i$  induces a subgraph of  $G$ . It suffices to work on a component of the induced subgraph. We apply induction on the number  $\alpha$  of cycles from  $E_{1/2}(x)$  in a component.

When  $\alpha = 1$ , it is trivial. Hence assume the claim holds for components with  $\alpha \geq 1$  cycles from  $E_{1/2}(x)$ . We consider a component with  $\alpha + 1$  cycles  $C_1, \dots, C_\alpha, C_{\alpha+1}$  from  $E_{1/2}(x)$ . Without loss of generality, assume that deleting  $C_{\alpha+1}$  yields a new component with  $\alpha$  cycles. By induction hypothesis, the claim holds for the resulting component. It remains to check edges in  $F_{\alpha+1}$ . If there exists an edge in  $F_{\alpha+1}$  violating the claim, relabel vertices and edges in  $C_{\alpha+1}$ . After at most one relabeling, all edges in  $F_{\alpha+1}$  satisfy the claim. We prove it by contradiction. Let  $f_1, f_2 \in F_{\alpha+1}$  be edges such that  $f_1$  satisfies the claim but  $f_2$  violates the claim. For  $i = 1, 2$ , let  $f_i = u_iw_i$ , where  $u_i$  is the endpoint in the resulting component and  $w_i$  is the endpoint in  $C_{\alpha+1}$ .

By assumption,  $u_1$  and  $w_1$  have different parity and  $u_2$  and  $w_2$  have the same parity. Analogous to the definition of cycles with cyclic preferences, we call path  $P = v_1v_2 \dots v_l$  a  $v_1v_l$ -path with linear preferences if  $v_iv_{i+1} \prec_{v_{i+1}} v_{i+1}v_{i+2}$  for  $i = 1, 2, \dots, l-2$ . Clearly, for any two vertices in the same component, there exists a path with linear preferences between them. Hence there exist a  $u_1u_2$ -path  $P_\alpha$  and a  $w_2w_1$ -path  $P_{\alpha+1}$ , both of which admit linear preferences. Moreover,  $u_1P_\alpha u_2 f_2 w_2 P_{\alpha+1} w_1 f_1 u_1$  form a cycle with cyclic preferences. We justify this cycle is odd by showing that the  $u_1u_2$ -path  $P_\alpha$  is even (*resp.* odd) if  $u_1$  and  $u_2$  have the same (*resp.* different) parity.

If  $u_1$  and  $u_2$  belong to the same cycle from  $E_{1/2}(x)$ , it is trivial. Hence assume  $u_1 \in C_s$  and  $u_2 \in C_t$ , where  $s, t \in \{1, 2, \dots, \alpha\}$  and  $s \neq t$ . We apply induction on the number  $\tau$  of cycles from  $E_{1/2}(x)$  involved in  $P_\alpha$ . Clearly,  $\tau \geq 2$ . When  $\tau = 2$ . Take  $v^s v^t \in F_s \cap F_t$  on  $P_\alpha$ . Let  $P_s$  be the part of  $P_\alpha$  from  $u_1$  to  $v^s$  in  $C_s$  and  $P_t$  be the part of  $P_\alpha$  from  $v^t$  to  $u_2$  in  $C_t$ . It follows that  $u_1 P_s v^s v^t P_t u_2$  form  $P_\alpha$ . By primary induction hypothesis,  $v^s$  and  $v^t$  have different parity since  $v^s v^t \in F_s \cap F_t$ . If  $u_1$  and  $u_2$  have the same parity, then  $P_s$  and  $P_t$  have different parity, implying that  $P_\alpha$  is even; if  $u_1$  and  $u_2$  have different parity, then  $P_s$  and  $P_t$  have the same parity, implying that  $P_\alpha$  is odd. Now assume  $\tau \geq 2$ . Let  $C_{k_1}, \dots, C_{k_\tau}, C_{k_{\tau+1}}$  be cycles from  $E_{1/2}(x)$  involved along  $P_\alpha$ . Take  $v^{k_\tau} v^{k_{\tau+1}} \in F_{k_\tau} \cap F_{k_{\tau+1}}$  on  $P_\alpha$ . Let  $P_{s, k_\tau}$  denote the part of  $P_\alpha$  from  $u_1$  to  $v^{k_\tau}$  and  $P_{k_\tau, t}$  denote the part of  $P_\alpha$  from  $v^{k_\tau}$  to  $u_2$ . Clearly,  $P_\alpha = u_1 P_{s, k_\tau} v^{k_\tau} P_{k_\tau, t} u_2$ . Since  $P_{s, k_\tau}$  involves  $\tau$  cycles and  $P_{k_\tau, t}$  involves two cycles, both length depend on the parity of endpoints. It follows that  $P_\alpha$  is even when  $u_1$  and  $u_2$  have the same parity, and  $P_\alpha$  is odd when  $u_1$  and  $u_2$  have different parity.

Hence when  $u_1$  and  $u_2$  have the same parity,  $w_1$  and  $w_2$  have different parity, implying that  $P_\alpha$  is even and  $P_{\alpha+1}$  is odd; when  $u_1$  and  $u_2$  have different parity,  $w_1$  and  $w_2$  have the same parity, implying that  $P_\alpha$  is odd and  $P_{\alpha+1}$  is even. Either case yields an odd cycle with cyclic preferences, a contradiction.

Therefore  $1/2$ -integral points are not vertices of  $FSM(G, \prec)$  as they can be perturbed by  $\epsilon z$  for small  $\epsilon$  without leaving  $FSM(G, \prec)$ . By Theorem 2.3,  $SM(G, \prec) = FSM(G, \prec)$  follows.  $\square$

Now we are ready to present a proof of our main theorem.

*Proof of Theorem 1.2.* It suffices to show the equivalence of (i), (iii) and (iv). Let  $D$  be a good orientation of line multigraph  $L(H)$ . Construct preference system  $(H, \prec)$  from  $D$ , and construct simple preference system  $(H', \prec')$  from  $(H, \prec)$  by substituting each parallel edge with a gadget as in Figure 1. By the construction,  $(H', \prec')$  admits no odd cycles with cyclic preferences. Now  $\sigma(D)$  can be viewed as a linear system defined on preference system  $(H, \prec)$  and consisting of constraints (3.1)-(3.5). Observe that constraints (3.1), (3.2) and (3.5) form the Rothblum

system  $\pi(H, \prec)$ , and constraints (3.3)-(3.4) are redundant with respect to  $\pi(H, \prec)$ . Hence  $FK(D) = FSM(H, \prec)$  follows.

By Lemma 4.1,  $FSM(H', \prec')$  is integral. Integrality of  $FSM(H, \prec)$  follows from Lemma 3.1, implying  $FK(D)$  is integral. Similar arguments apply to any induced subdigraphs of  $D$ . Hence (i)  $\implies$  (iii).

By Corollary 4.2,  $\pi(H', \prec')$  is TDI. Total dual integrality of  $\pi(H, \prec)$  follows from Lemma 3.2. Since  $\pi(H, \prec)$  is part of  $\sigma(D)$  and the other constraints (3.3)-(3.4) are redundant in  $\sigma(D)$  with respect to  $\pi(H, \prec)$ , total dual integrality of  $\sigma(D)$  follows. Similar arguments apply to any induced subdigraphs of  $D$ . Hence (iii)  $\implies$  (iv).

By a theorem of Edmonds and Giles [8], implication (iv)  $\implies$  (iii) follows directly.

To prove implication (iii)  $\implies$  (i), we assume the contrary. Observe that  $D$  being kernel ideal implies the existence of kernels for any induced subdigraphs of  $D$ . Let  $D$  be a digraph such that  $D$  is kernel ideal but not good. Then there either exists a clique containing directed cycles or exists a directed odd cycle without (pseudo-)chords. We show that neither case is possible. If  $D$  has a clique containing directed cycles, we consider the subdigraph induced on this clique. There is no kernel for this induced subdigraph, a contradiction. If  $D$  contains a directed odd cycle without (pseudo-)chords, we restrict ourselves to the subdigraph induced on this directed odd cycle. There is no kernel for this induced subdigraph either, a contradiction.  $\square$

## 5 Discussions

It is natural to consider whether our result applies to superorientations of line multigraph. However, the connection between kernels and stable matchings seems to be broken by superorientations. Indeed, consider the digraph  $D = (V, A)$ , where  $V = \{1, 2, 3, 4\}$  and  $A = \{(1, 2), (2, 1), (3, 1), (3, 4), (4, 2)\}$ . Clearly,  $D$  is a good superorientation of a 4-cycle which is a line graph. However we cannot construct a preference system with strict linear preferences from  $D$  such that kernels in  $D$  correspond to stable matchings in the preference system. Besides, the fractional kernel polytope  $FK(D)$  has three vertices  $(1, 0, 0, 1)$ ,  $(1/2, 1/2, 0, 1/2)$  and  $(0, 1, 1, 0)$  including a fractional one.

## Acknowledgments

The author is grateful to Prof. Wenan Zang for his valuable suggestions.

## References

- [1] Egres Open. Available from <http://lemon.cs.elte.hu/egres/open>.
- [2] H. G. ABELEDO AND U. G. ROTHBLUM, *Stable matchings and linear inequalities*, Discrete Appl. Math., 54 (1994), pp. 1–27.
- [3] O. V. BORODIN, A. V. KOSTOCHKA, AND D. R. WOODALL, *On kernel-perfect orientations of line graphs*, Discrete Math., 191 (1998), pp. 45–49.
- [4] K. CECHLÁROVÁ AND T. FLEINER, *On a generalization of the stable roommates problem*, ACM Trans. Algorithms, 1 (2005), pp. 143–156.
- [5] Q. CHEN, X. CHEN, AND W. ZANG, *A polyhedral description of kernels*, Math. Oper. Res., to appear.
- [6] X. CHEN, G. DING, X. HU, AND W. ZANG, *The maximum-weight stable matching problem: duality and efficiency*, SIAM J. Discrete Math., 26 (2012), pp. 1346–1360.
- [7] W. COOK, *Operations that preserve total dual integrality*, Oper. Res. Lett., 2 (1983), pp. 31–35.
- [8] J. EDMONDS AND R. GILES, *A min-max relation for submodular functions on graphs*, Ann. of Discrete Math., 1 (1977), pp. 185–204.
- [9] T. KIRÁLY AND J. PAP, *Total dual integrality of Rothblum’s description of the stable-marriage polyhedron*, Math. Oper. Res., 33 (2008), pp. 283–290.
- [10] F. MAFFRAY, *Kernels in perfect line-graphs*, J. Combin. Theory Ser. B, 55 (1992), pp. 1–8.
- [11] U. G. ROTHBLUM, *Characterization of stable matchings as extreme points of a polytope*, Math. Programming, 54 (1992), pp. 57–67.
- [12] A. SCHRIJVER, *Theory of Linear and Integer Programming*, Wiley-Interscience Series in Discrete Mathematics, John Wiley & Sons, 1986.