

VELOX PEPTIDES

THE RESEARCHER'S HANDBOOK

The Researcher's Handbook

Research Peptide Reference — 2026 Edition

A plain-English reference covering reconstitution, storage, certificates of analysis and 14 research compounds across five categories. Quality and documentation, explained clearly.

≥ 99% HPLC

Janoshik tested

CoA per batch

2026 EDITION

veloxpeps.com

Velox Peptides · CRP Labs Ltd (UK, NI738125) · Hollywood, Northern Ireland
For in vitro research use only. Not for human or veterinary consumption.

Contents

01 Part 1 · Research Fundamentals

What research peptides are	p3
Reconstitution — basics & worked example	p4
Storage & handling	p5
How to read a Certificate of Analysis	p6

02 Part 2 · The Compound Reference

Cognitive — Semax, Selank, DSIP	p8
Metabolic — Retatrutide, MOTS-c, NAD+	p12
Recovery — BPC-157, TB-500, KPV, GHK-Cu	p16
Growth — CJC-1295, Tesamorelin	p21
Antioxidant — Glutathione, Melanotan II	p24

03 Glossary

04 Why Velox · How to Order · Compliance

What Research Peptides Are

A peptide is a short chain of amino acids — the same building blocks that make up proteins. Research peptides are synthetic (lab-made) versions of these chains, supplied as a freeze-dried (lyophilised) powder. They are studied in vitro (test-tube and laboratory work) and in animal models to understand biological pathways — the chains of chemical signals that cells use to communicate.

Everything in this handbook describes preclinical research observations and product quality only. These are not medicines. Nothing here is a health, medical or benefit claim, and no compound is intended for human or veterinary use, consumption, injection, dosing, diagnosis or treatment.

HOW VELOX SUPPLIES RESEARCH MATERIAL

- $\geq 99\%$ HPLC purity — batch-verified by high-performance liquid chromatography.
- Janoshik third-party testing — independent verification, not in-house only.
- Certificate of Analysis (CoA) supplied per batch, available on request.
- 24h Royal Mail Tracked UK dispatch · free UK shipping over £80 · Pay by Bank.
- Supplied to qualified researchers aged 21+ for in vitro research use only.

Reconstitution — The Basics

Reconstitution means mixing a freeze-dried powder back into a liquid. The usual diluent is bacteriostatic water — sterile water containing a small amount of benzyl alcohol (about 0.9%) that suppresses bacterial growth, so a reconstituted vial stays stable in the fridge for longer than plain sterile water would allow.

$$\text{mg of peptide} \div \text{mL of water} = \text{mg/mL concentration}$$

the core reconstitution formula

WORKED EXAMPLE

A 10 mg vial reconstituted with 2 mL of bacteriostatic water gives $10 \div 2 = 5$ mg/mL. Drawing 0.1 mL from that vial therefore contains 0.5 mg (500 mcg) of peptide for the research preparation.

CONCENTRATION REFERENCE — VIAL SIZE × WATER VOLUME

Vial size	1 mL	2 mL	3 mL	5 mL
5 mg	5.0 mg/mL	2.5 mg/mL	1.67 mg/mL	1.0 mg/mL
10 mg	10.0 mg/mL	5.0 mg/mL	3.33 mg/mL	2.0 mg/mL
15 mg	15.0 mg/mL	7.5 mg/mL	5.0 mg/mL	3.0 mg/mL
20 mg	20.0 mg/mL	10.0 mg/mL	6.67 mg/mL	4.0 mg/mL

Reconstitution calculator: veloxpeps.com/tools/reconstitution-calculator/

Storage & Handling

- Lyophilised (freeze-dried) powder is the most stable form. Stored sealed at -20°C and kept dry, it remains stable for extended periods.
- Once reconstituted, keep the vial refrigerated at $2-8^{\circ}\text{C}$ and use within a research-appropriate window; the liquid form is less stable than the powder.
- Minimise freeze-thaw cycles. Repeated freezing and thawing stresses peptide bonds and can degrade the material; aliquoting before freezing avoids re-thawing a whole vial.
- Use sterile technique — clean surfaces, an alcohol-wiped vial stopper, and a fresh sterile needle — to avoid introducing contaminants.
- Protect from light and heat. Degradation is driven by temperature, light, repeated thawing and moisture; cloudiness or colour change can indicate a compromised sample.

STATE • TEMPERATURE • NOTES

State	Temperature	Notes
Lyophilised powder	-20°C (sealed, dry)	Most stable; long-term storage form.
Reconstituted vial	$2-8^{\circ}\text{C}$ (refrigerated)	Use within research window; avoid light.
Aliquots	-20°C	Freeze once; thaw only what is needed.
In use / bench	Brief, cool	Return to fridge promptly; avoid heat.

How to Read a CoA

A Certificate of Analysis (CoA) is the laboratory report for a specific batch. It documents what is in the vial and how it was verified. Here is what each part means and why it matters.

HPLC purity (%)

High-performance liquid chromatography separates a sample into its components and measures the target peptide as a percentage. A figure of $\geq 99\%$ means highly pure material with minimal related impurities.

Mass spectrometry / molecular weight

Mass spec confirms the molecule's measured mass matches the expected molecular weight for that peptide — confirming identity, not just purity.

Appearance

A simple physical check (for example, a white lyophilised powder) recorded so the researcher knows what a correct sample should look like.

PASS

An overall result indicating the batch met every specification on the certificate for that test panel.

WHY THIRD-PARTY (JANOSHIK) TESTING MATTERS

Janoshik is an independent analytical lab. Third-party testing means purity and identity are verified by an outside party rather than self-reported, giving researchers documentation they can trust. Velox supplies a CoA per batch on this basis.

Cognitive

Neuropeptide research tools studied for brain-signalling pathways.

COMPOUNDS IN THIS SECTION

Semax ACTH(4-7) analogue · BDNF/NGF

Selank Tuftsin analogue · GABAergic

DSIP Endogenous · sleep modulation

Semax

CAS: 80714-61-0

·

Class: ACTH(4-7) analogue

·

Seq: Met-Glu-His-Phe-Pro-Gly-Pro

·

HPLC: ≥99.0%

WHAT IT IS

Semax is a synthetic heptapeptide (seven amino acids, sequence Met-Glu-His-Phe-Pro-Gly-Pro) copied from a small fragment of the natural hormone ACTH, with an added Pro-Gly-Pro tail that slows enzymatic breakdown so it persists far longer than the plain fragment. It is a neuropeptide research reagent that retains nerve-related signalling without the stress-hormone-triggering part of the full hormone, making it a clean tool for studying those signals in isolation. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Linked in rodent studies to increased BDNF and NGF, proteins that help nerve cells survive, grow and form new connections.
- Associated with region-specific, multidirectional regulation of neurotrophin gene activity across brain regions over time.
- Studied in stroke (ischaemia) and oxidative-stress models for nerve-cell survival and activation of nerve-growth genes.
- Its Pro-Gly-Pro metabolite linked to switching on neurotrophic genes while dialling down inflammation- and cell-death-related genes.
- Examined for effects on attention, learning, and the dopamine and serotonin systems. Research observations only.

Selank

CAS: 129954-34-3

Class: Tuftsin analogue

Seq: Thr-Lys-Pro-Arg-Pro-Gly-Pro

HPLC: $\geq 98.9\%$

WHAT IT IS

Selank is a synthetic heptapeptide (Thr-Lys-Pro-Arg-Pro-Gly-Pro) derived from the natural immune peptide tuftsin, with an added Pro-Gly-Pro tail that extends its stability against enzymatic breakdown. Developed in Russia at the Institute of Molecular Genetics, it is studied as a neuropeptide research tool, often paired with Semax, for how it relates to the calming GABA system, brain chemistry and stress responses in animal models. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Best studied for modulating the GABA system indirectly (allosterically) rather than at the benzodiazepine site.
- In rodent anxiety models, associated with a calming effect without the heavy sedation or motor impairment of classical anxiolytics.
- Linked to increased BDNF in the hippocampus and to serotonin signalling, pathways tied to memory, mood and coping with stress.
- Reported to slow the breakdown of enkephalins (natural pain-relieving signals), reflecting its tuftsin origin.
- Studied for effects on the immune system and cytokines, linking nervous and immune signalling. Research observations only.

DSIP

CAS: 62568-57-4

Length: 9 amino acids

Origin: Endogenous neuropeptide

HPLC: ≥98.7%

WHAT IT IS

DSIP (Delta Sleep-Inducing Peptide) is a nonapeptide (nine amino acids, sequence Trp-Ala-Gly-Gly-Asp-Ala-Ser-Gly-Glu) first identified in rabbit blood during sleep experiments and named after the slow delta brain waves of deep sleep. It is an endogenous peptide — one the body makes naturally. Its exact mechanism and receptor remain unresolved, which is part of why it is studied as a tool for how small peptides participate in sleep and hormone signalling. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Best known for its link to slow-wave (delta) sleep, the deepest sleep stage, in rat and rabbit studies.
- Associated with more deep sleep and REM sleep; blocking the animal's own DSIP reduced rebound deep sleep after sleep loss.
- Studied for links to sleep-time hormone release, including growth hormone, via a somatostatin/dopamine pathway.
- Examined for relationships to calming the stress response and to antioxidant activity in animal studies.
- Investigated for a broader role in keeping the body's internal systems balanced. Research observations only.

Metabolic

Compounds studied for cellular energy, fuel handling and incretin pathways.

COMPOUNDS IN THIS SECTION

Retatrutide Triple agonist · GLP-1/GIP/glucagon

MOTS-c Mitochondrial-derived · AMPK

NAD⁺ Coenzyme · sirtuins / redox

Retatrutide

Code: LY3437943 · Class: GIP/GLP-1/glucagon triple agonist · Form: Lyophilised powder · HPLC: $\geq 99\%$

WHAT IT IS

Retatrutide (LY3437943, developed by Eli Lilly) is a synthetic peptide built to act on three receptors at once — GLP-1, GIP and glucagon — making it a triple agonist (a molecule that switches on all three 'locks'). Each receptor helps control metabolism: how the body handles energy, blood sugar and fat. Because it engages all three simultaneously, it works in a more complex way than single- or dual-receptor compounds, and is one of the most studied peptides in incretin and receptor research. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- GLP-1 receptor activation linked in animal studies to glucose-dependent insulin release, slower gastric emptying and appetite signals.
- GIP receptor activation associated with insulin release and changes in how fat tissue stores and breaks down fat in mice.
- Glucagon receptor activation studied in the liver for glucose output and for increased energy expenditure (thermogenesis).
- Characterised in rodent and non-human primate models with binding at all three receptors and once-weekly pharmacokinetics.
- Investigated in Phase 1 and Phase 2 clinical trials (human data, distinct from preclinical work) published in major journals.

MOTS-c

Length: 16 amino acids

·

Origin: Mitochondrial 12S rRNA

·

Pathway: AMPK signalling

·

HPLC: ≥99%

WHAT IT IS

MOTS-c is a 16-amino-acid mitochondrial-derived peptide — unusually, its instructions are encoded inside the mitochondria (the cell's energy 'power plants') rather than the nucleus. It acts as a messenger from the mitochondria to the rest of the cell and is studied mainly for how it switches on AMPK, the cell's low-fuel energy sensor. First described in 2015, it is one of the first mitochondrial messengers shown to have a clear role in controlling energy and fuel. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Activates the AMPK energy sensor indirectly by interfering with the folate cycle, mainly in skeletal muscle.
- Under metabolic stress, observed moving from the mitochondria into the nucleus to help regulate defence and antioxidant genes.
- A rare clear example of a peptide that physically carries a signal from the mitochondria to the cell's control centre.
- In rodent studies, linked to maintaining insulin sensitivity with age or a high-fat diet and to less diet-induced weight gain.
- Described as exercise-responsive and linked to fitness and age-related decline. Research observations only.

NAD⁺

CAS: 53-84-9 · Class: Pyridine dinucleotide coenzyme · Pathway: Sirtuins / redox · HPLC: ≥99.5%

WHAT IT IS

NAD⁺ (nicotinamide adenine dinucleotide) is a coenzyme — a small helper molecule — found in every living cell. It has two main roles: carrying energy (electrons) during the reactions that turn food into ATP, cycling between its loaded form (NAD⁺) and unloaded form (NADH); and acting as fuel for enzymes called sirtuins and PARPs. This dual role links a cell's energy state to research on ageing, metabolism and stress, and is why NAD⁺ and its precursors are widely studied. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Central to redox metabolism: the NAD⁺/NADH balance is used as a quick read on a cell's energy state.
- Required fuel for sirtuins, enzymes that link protein modification to metabolism, stress responses and ageing.
- Consumed by PARP enzymes during DNA repair, creating a studied tug-of-war with sirtuin demand under stress.
- Reported to decline with age in lab animals, a change linked to reduced energy production and mitochondrial function.
- Used as the reference reagent when studying precursors such as NMN and NR. Research observations only.

Recovery

Peptides studied in tissue-repair, angiogenesis and anti-inflammatory models.

COMPOUNDS IN THIS SECTION

BPC-157 Pentadecapeptide · tissue repair

TB-500 Thymosin β 4 · actin / migration

KPV α -MSH fragment · anti-inflammatory

GHK-Cu Copper tripeptide · collagen / repair

BPC-157

CAS: 137525-51-0

·

Seq: GEPPPGKPADDAGLV (15 aa)

·

Class: Synthetic pentadecapeptide

·

HPLC: ≥99.4%

WHAT IT IS

BPC-157 (Body Protection Compound-157) is a synthetic pentadecapeptide — fifteen amino acids, sequence GEPPPGKPADDAGLV — based on a small fragment of a protein found in gastric juice. It is notably stable, staying intact in water and stomach-like lab conditions where many peptides break down, which is one reason it is used so often in studies of how cells heal and move. Researchers study it in relation to tissue repair, angiogenesis and gut protection. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Studied in cells and rodents for angiogenesis, with links to the VEGFR2 signal and to nitric oxide in blood-vessel-lining cells.
- Lab work on tendon and ligament fibroblasts reported changes in cell growth, survival, movement and growth-hormone receptors.
- Because it derives from a gastric protein, much early research examined the gut lining and the gut-brain axis in injury models.
- Often studied alongside TB-500 in tissue-repair models as a complementary, blood-vessel-focused pathway.
- Its in-solution stability makes it a frequently used probe for cell healing and migration. Research observations only.

TB-500

CAS: 77591-33-4

Class: Actin-binding peptide

Parent: Thymosin β 4 (43-aa)HPLC: $\geq 99\%$

WHAT IT IS

TB-500 is a synthetic peptide built around the active region of Thymosin β 4, a natural 43-amino-acid protein found in almost every mammalian cell and abundant in platelets and wound fluid. Its core job centres on actin — the protein scaffolding cells use to move and hold their shape. Thymosin β 4 sequesters loose actin subunits, helping control the balance cells need to rapidly rebuild that scaffolding when migrating, which puts it at the centre of tissue-repair research. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Sequesters loose (G-)actin, keeping a ready supply cells use to build the leading-edge structures that pull them forward.
- Linked to blood-vessel cells moving, forming tubes and growing new vessels (angiogenesis) in wound-healing studies.
- In rodent studies, reported to grow more new vessels and speed skin-wound repair in both young and old animals.
- Studied for helping inflammation settle and for affecting matrix-remodelling enzymes during healing.
- Examined in heart and corneal tissue models for cell survival and migration after injury. Research observations only.

CAS: 13147-97-2

·

Seq: Lys-Pro-Val

·

Origin: C-terminal α -MSH fragment

·

HPLC: $\geq 98.5\%$

WHAT IT IS

KPV is a tripeptide — just three amino acids, lysine-proline-valine — forming the tail-end fragment of the hormone alpha-melanocyte-stimulating hormone (α -MSH). It retains much of α -MSH's anti-inflammatory character while dropping the pigment-changing part, making it a clean tool for studying inflammation. Notably, it calms inflammation without engaging the melanocortin receptor at all, instead entering cells through a different doorway. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Best studied for turning down NF- κ B, a master inflammation 'on switch' inside cells, and the MAP-kinase pathway.
- In cell tests, even nanomolar amounts lowered NF- κ B activity and reduced pro-inflammatory cytokine release.
- Enters cells via the PepT1 transporter on gut-lining and immune cells, acting independently of melanocortin receptors.
- In mouse colitis models, oral KPV reported to reduce colitis severity and lower inflammation messengers.
- Also studied in skin and wound inflammation alongside other repair peptides. Research observations only.

GHK-Cu

CAS: 49557-75-7 · Seq: Gly-His-Lys · Cu(II) · Class: Copper-binding tripeptide · HPLC: ≥99%

WHAT IT IS

GHK-Cu (copper-tripeptide-1) is a tripeptide — glycine-histidine-lysine — bound to a single copper ion. First identified in human blood plasma in the 1970s by Pickart and Thaler, who observed it changing how liver cells from older donors behaved. Natural GHK levels fall with age, which is one reason it is studied as a research tool in tissue-repair, skin-cell biology and antioxidant pathways. The research-grade version is chemically identical to the natural one, with the copper attached. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- In dermal-fibroblast studies, linked to switching on genes for type I collagen, elastin, decorin and glycosaminoglycans.
- Studied as a tool for how copper moves within tissue and for relationships to antioxidant enzymes under oxidative stress.
- In rodent and lab-dish wound models, observed alongside angiogenesis (new blood-vessel growth).
- Associated with movement of skin-building fibroblast cells and rebuilding of the tissue's support structure.
- Studied for the age-related decline of natural GHK and what restoring it may do in models. Research observations only.

Growth

GHRH-analogue tools studied in somatotropic (growth-hormone) research.

COMPOUNDS IN THIS SECTION

CJC-1295 GHRH analogue · GH pulse

Tesamorelin GHRH analogue · GH/IGF-1 axis

CJC-1295

Seq: Modified GHRH(1-29)

·

Class: GHRH analogue

·

Target: GHRH receptor

·

HPLC: $\geq 99.1\%$

WHAT IT IS

CJC-1295 without DAC — also called Modified GRF(1-29) — is a synthetic peptide that acts like GHRH, the natural hormone that prompts the body to release its own growth hormone in short bursts. It starts from the active GHRH(1-29) fragment and swaps in four amino acids that protect it from enzymes such as DPP-IV, so it lasts longer than plain GHRH(1-29). The 'without DAC' form omits the albumin-binding handle, so it acts for about 30 minutes rather than days. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Docks onto the GHRH receptor on pituitary somatotroph cells, raising cAMP to prompt growth-hormone production and release.
- Works one step before growth hormone, so the gland keeps its own natural feedback controls in research models.
- The no-DAC form's short ~30-minute action is studied for producing separate GH bursts resembling the body's natural rhythm.
- Studies report preserved pulsatile GH release with higher mean GH and IGF-1 rather than a flattened rhythm.
- DAC and no-DAC forms used as a complementary pair to separate signalling pattern from simple activation. Research observations only.

Tesamorelin

CAS: 218949-48-5

·

Class: GHRH analogue

·

Target: GHRH receptor

·

HPLC: $\geq 99\%$

WHAT IT IS

Tesamorelin is a GHRH analogue — a synthetic copy of the natural GHRH signal that tells the pituitary to release growth hormone. Chemically it is the full 44-amino-acid GHRH chain with an added trans-3-hexenoyl cap that armours the peptide's weak end, so it lasts much longer than plain GHRH, which enzymes chop up within minutes. It works one step before growth hormone, prompting the pituitary to release its own in natural bursts, which makes it a research tool for studying that earlier step. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Binds the GHRH receptor on pituitary somatotrophs, raising cAMP to make the cell build and release growth hormone.
- Prompts pulsatile GH release in the body's natural rhythm, keeping normal feedback in place — unlike giving GH directly.
- The released growth hormone triggers IGF-1, extending research to the wider GH/IGF-1 axis.
- Most studied for metabolic context, especially relationships to visceral fat and how the body handles fats.
- Characterised pharmacologically (as TH9507) as a GH-axis research tool. Research observations only.

Antioxidant

Redox and melanocortin reference reagents for oxidative-stress research.

COMPOUNDS IN THIS SECTION

Glutathione Tripeptide · master antioxidant

Melanotan II Cyclic α -MSH · melanocortin

Glutathione

CAS: 70-18-8

Seq: γ -Glu-Cys-Gly

Pathway: Cellular redox / antioxidant

HPLC: $\geq 99.2\%$

WHAT IT IS

Glutathione (GSH) is a tripeptide — glutamate, cysteine and glycine — and the most abundant antioxidant inside cells. Its working part is a sulphur-containing thiol (–SH) group on its cysteine, which donates electrons to neutralise reactive oxygen species, switching between its fresh form (GSH) and used-up form (GSSG). Found in nearly every cell and called the 'master antioxidant', the GSH:GSSG ratio is a standard measure of cellular stress, making it both a study subject and a reference reagent. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- Drives the GSH/GSSG redox cycle: it neutralises reactive oxygen species, then glutathione reductase recycles it using NADPH.
- Acts as fuel for glutathione peroxidase enzymes that break hydrogen peroxide and damaged fats into safe products.
- Supports detoxification: glutathione S-transferases attach it to foreign substances so cells can clear them.
- The GSH:GSSG balance shifts measurably under stress, so it is used to track how stressed a cell is.
- Valued as a steady reference reagent in oxidative-stress studies. Research observations only.

Melanotan II

Class: Cyclic α -MSH analogue · Targets: MC1R–MC5R · Key receptors: MC1R, MC3R, MC4R · HPLC: $\geq 99.3\%$

WHAT IT IS

Melanotan II (MT-2) is a synthetic cyclic peptide copy of the hormone alpha-melanocyte-stimulating hormone (α -MSH). First made at the University of Arizona in the 1980s–90s, its ring shape makes it more stable than natural α -MSH. In the lab it activates four of the five melanocortin switches (MC1R, MC3R, MC4R, MC5R), largely sparing MC2R, making it one of the broadest melanocortin reference tools. It is not a medicine and not for tanning, cosmetic, human or veterinary use. Supplied strictly for in vitro research use.

WHAT'S BEEN STUDIED

- A non-selective melanocortin agonist activating MC1R, MC3R, MC4R and MC5R at once — its main studied feature.
- Used to switch on MC1R in melanocytes to study how that switch controls melanin (pigment) production.
- Used in rodent studies to probe MC3R and MC4R, brain switches involved in energy use and feeding behaviour.
- Serves as a 'yardstick' reference agonist against which newer, more selective MC3R/MC4R compounds are compared.
- Studied via site-specific brain microinjection for central melanocortin pathways. Research observations only.

Glossary

Peptide	A short chain of amino acids, the building blocks of proteins.
Lyophilised	Freeze-dried into a dry, stable powder form.
Reconstitution	Mixing a freeze-dried powder back into a liquid.
Bacteriostatic water	Sterile water with ~0.9% benzyl alcohol that suppresses bacterial growth.
HPLC	High-performance liquid chromatography; measures purity as a percentage.
Mass spectrometry	Confirms a molecule's mass matches its expected molecular weight (identity).
CoA	Certificate of Analysis — the lab report for a specific batch.
In vitro	Test-tube / laboratory work, outside a living organism.
Preclinical	Research in cells and animals, before any human trials.
Agonist	A molecule that switches a receptor on.
Receptor	A 'lock' on or in a cell that a molecule fits to send a signal.
Angiogenesis	The growing of new blood vessels.
BDNF	Brain-derived neurotrophic factor; helps nerve cells grow and survive.
AMPK	A cellular energy sensor that responds to low fuel.
Antioxidant	A molecule that neutralises cell-damaging reactive particles.
Pharmacokinetics	How a compound is processed over time (e.g. half-life).

CLOSING

Why Velox

- $\geq 99\%$ HPLC purity, batch-verified — every compound documented.
- Janoshik third-party testing — independent verification, not in-house only.
- Certificate of Analysis supplied per batch, available on request.
- 24h Royal Mail Tracked UK dispatch · free UK shipping over £80.
- Pay by Bank checkout · supplied to qualified researchers aged 21+.

HOW TO ORDER

Browse the catalogue at veloxpeps.com, add research compounds to your order and check out with Pay by Bank. Need help choosing a vial size or working out concentration? Use the reconstitution calculator at veloxpeps.com/tools/reconstitution-calculator/. Request a batch CoA any time at support@veloxpeps.com.

COMPLIANCE

All Velox products are supplied as research reagents for in vitro research use only. They are NOT for human or veterinary use, consumption, injection, dosing, diagnosis or treatment, and nothing in this handbook is a health, medical or benefit claim. Everything described is preclinical research observation or product quality and documentation. Products are not medicines and have not been evaluated or approved by the MHRA or FDA.

Start your research at veloxpeps.com

VELOX PEPTIDES · 2026

Velox Peptides · CRP Labs Ltd (UK, NI738125) · Hollywood, Northern Ireland · support@veloxpeps.com