

EEL 4930 Stats – Lecture 19

EEL 4930 Stats – Lecture 19

RANDOM VARIABLES (RVs)

EEL 4930 Stats – Lecture 19

RANDOM VARIABLES (RVs)

- What is a random variable?

numeric occurrence
that is random

EEL 4930 Stats – Lecture 19

RANDOM VARIABLES (RVs)

- What is a random variable?
- We define a random variable is defined on a probability space (S, \mathcal{F}, P) as a **function**

EEL 4930 Stats – Lecture 19

RANDOM VARIABLES (RVs)

- What is a random variable?
- We define a random variable is defined on a probability space (S, \mathcal{F}, P) as a function from S

EEL 4930 Stats – Lecture 19


RANDOM VARIABLES (RVs)

- What is a random variable?
- We define a random variable is defined on a probability space (S, \mathcal{F}, P) as a function from S to \mathcal{R}

Δ 5

EX

Create a binary RV from tossing a fair coin



$$X(\Delta): \begin{aligned} X(H) &= 1 \\ X(T) &= 0 \end{aligned}$$


$$P[X(\Delta) = 1] = P[\{ \Delta \mid X(\Delta) = 1 \}] = P[\{ H \}] = \frac{1}{2}$$

$$P[X(\Delta) = 0] = P[\{ T \}] = \frac{1}{2}$$

$$P[X(\Delta) = x] = 0, x \neq 0, 1$$

Create a binary RV from tossing a fair coin

EX twice



$$X(\omega) : \quad \begin{aligned} X(HH) &= X(HT) = X(TH) = 1 \\ X(TT) &= 0 \end{aligned}$$

$$P[X(\omega) = 1] = P[\omega \mid X(\omega) = 1] \quad \text{such that, not given!}$$

$$= P[\{HH, HT, TH\}] = 3/4$$

$$P[X=0] = P[\{\omega \mid X(\omega) = 0\}] = 1/4$$



EX

Create another RV from tossing a fair coin twice

DISCRETE RANDOM VARIABLES

A discrete random variable



DISCRETE RANDOM VARIABLES



A **discrete random variable** has nonzero probability at a countable number of values.

PROBABILITY MASS FUNCTION

PROBABILITY MASS FUNCTION

For a discrete RV, the *probability mass function*



PROBABILITY MASS FUNCTION

For a discrete RV, the *probability mass function* (pmf)



PROBABILITY MASS FUNCTION



For a discrete RV, the *probability mass function* (pmf) is

$$P(X = x) = P[X \leq x] - P[X < x]$$

PROBABILITY MASS FUNCTION



For a discrete RV, the *probability mass function* (pmf) is

$$P(X = x) = P[X \leq x] - P[X < x]$$

EX: Roll a fair 6-sided die

EX: Roll a fair 6-sided die

$X = \#$ on top face

EX: Roll a fair 6-sided die

$X = \#$ on top face

$$P(X = x) =$$

EX: Roll a fair 6-sided die

$X = \#$ on top face

$$P(X = x) = \begin{cases} 1/6, & x = 1, 2, \dots, 6 \\ \end{cases}$$

EX: Roll a fair 6-sided die

$X = \#$ on top face

$$P(X = x) = \begin{cases} 1/6, & x = 1, 2, \dots, 6 \\ 0, & \text{o.w.} \end{cases}$$

EX: Flip a fair coin until heads occurs

EX: Flip a fair coin until heads occurs

X = # of flips

EX: Flip a fair coin until heads occurs

X = # of flips

$$P(X = x) =$$

EX: Flip a fair coin until heads occurs

X = # of flips

$$P(X = x) = \begin{cases} \left(\frac{1}{2}\right)^x, & x = 1, 2, \dots \\ 0, & \text{o.w.} \end{cases}$$

EX: Flip a fair coin until heads occurs

X = # of flips

$$P(X = x) = \begin{cases} \left(\frac{1}{2}\right)^x, & x = 1, 2, \dots \\ 0, & \text{o.w.} \end{cases}$$

CUMULATIVE DISTRIBUTION FUNCTION

If (S, \mathcal{F}, P) is a prob. space with $X(\omega)$ a real RV on S , the **cumulative distribution function**



CUMULATIVE DISTRIBUTION FUNCTION

If (S, \mathcal{F}, P) is a prob. space with $X(\omega)$ a real RV on S , the **cumulative distribution function** (cdf)



CUMULATIVE DISTRIBUTION FUNCTION

If (S, \mathcal{F}, P) is a prob. space with $X(\omega)$ a real RV on S , the **cumulative distribution function** (cdf), denoted $F_X(x)$



CUMULATIVE DISTRIBUTION FUNCTION

If (S, \mathcal{F}, P) is a prob. space with $X(\omega)$ a real RV on S , the **cumulative distribution function** (cdf), denoted $F_X(x)$ is



$$F_X(x) = P[\{\omega | X(\omega) \in (-\infty, x]\}]$$

and $\omega \in S$

CUMULATIVE DISTRIBUTION FUNCTION

If (S, \mathcal{F}, P) is a prob. space with $X(\omega)$ a real RV on S , the **cumulative distribution function** (cdf), denoted $F_X(x)$ is

$$\begin{aligned} F_X(x) &= P[\{\omega | X(\omega) \in (-\infty, x]\}] \\ &\quad \text{and } \omega \in S] \\ &= P(X \leq x). \end{aligned}$$



CUMULATIVE DISTRIBUTION FUNCTION

If (S, \mathcal{F}, P) is a prob. space with $X(\omega)$ a real RV on S , the **cumulative distribution function** (cdf), denoted $F_X(x)$ is

$$\begin{aligned} F_X(x) &= P[\{\omega | X(\omega) \in (-\infty, x]\}] \\ &\quad \text{and } \omega \in S] \\ &= P(X \leq x). \end{aligned}$$



- $F_X(x)$ is also sometimes called the *probability distribution function (PDF)*, but I will avoid this terminology to avoid confusion with another function we will use, called the probability density function (pdf)

- $F_X(x)$ is a prob. measure

- $F_X(x)$ is a prob. measure
 - Thus $F_X(x)$ inherits all the properties of a probability measure (axioms and corollaries still apply)



EX

Find and plot the cdfs for the previous two examples

