

# EEL 4930 Stats – Lecture 21

# EEL 4930 Stats – Lecture 20

## **IMPORTANT RANDOM VARIABLES**

# EEL 4930 Stats – Lecture 20

## **IMPORTANT RANDOM VARIABLES**

### **Discrete RVs**

# EEL 4930 Stats – Lecture 20

## IMPORTANT RANDOM VARIABLES

### Discrete RVs

#### 1. Bernoulli RV

# EEL 4930 Stats – Lecture 20

## IMPORTANT RANDOM VARIABLES

### Discrete RVs

#### 1. Bernoulli RV

- An event  $A \in \mathcal{F}$  is considered a “success”

# EEL 4930 Stats – Lecture 20

## IMPORTANT RANDOM VARIABLES

### Discrete RVs

#### 1. Bernoulli RV

- An event  $A \in \mathcal{F}$  is considered a “success”
- A Bernoulli RV  $X$  is defined by

$$X = \begin{cases} 1, & s \in A \\ 0, & s \notin A \end{cases}$$

# EEL 4930 Stats – Lecture 20

## IMPORTANT RANDOM VARIABLES

### Discrete RVs

#### 1. Bernoulli RV

- An event  $A \in \mathcal{F}$  is considered a “success”
- A Bernoulli RV  $X$  is defined by

$$X = \begin{cases} 1, & s \in A \\ 0, & s \notin A \end{cases}$$

- The pmf for a Bernoulli RV  $X$  can be found formally as



- The pmf for a Bernoulli RV  $X$  can be found formally as

$$P(X = 1)$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$P(X = 1) = P(X(s) = 1)$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$\begin{aligned} P(X = 1) &= P\left(X(s) = 1\right) \\ &= P\left(\{s | s \in A\}\right) \end{aligned}$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$\begin{aligned} P(X = 1) &= P\left(X(s) = 1\right) \\ &= P\left(\{s | s \in A\}\right) = P(A) \end{aligned}$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$\begin{aligned} P(X = 1) &= P\left(X(s) = 1\right) \\ &= P\left(\{s | s \in A\}\right) = P(A) \triangleq p \end{aligned}$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$\begin{aligned} P(X = 1) &= P\left(X(s) = 1\right) \\ &= P\left(\{s | s \in A\}\right) = P(A) \triangleq p \end{aligned}$$

So,

$$P(X = x) = \begin{cases} p & x = 1 \\ 1 - p & x = 0 \end{cases}$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$\begin{aligned} P(X = 1) &= P\left(X(s) = 1\right) \\ &= P\left(\{s | s \in A\}\right) = P(A) \triangleq p \end{aligned}$$

So,

$$P(X = x) = \begin{cases} p & x = 1 \\ 1 - p & x = 0 \end{cases}$$

- The pmf for a Bernoulli RV  $X$  can be found formally as

$$\begin{aligned} P(X = 1) &= P\left(X(s) = 1\right) \\ &= P\left(\{s | s \in A\}\right) = P(A) \triangleq p \end{aligned}$$

So,

$$P(X = x) = \begin{cases} p & x = 1 \\ 1 - p & x = 0 \\ 0 & x \neq 0, 1 \end{cases}$$



- **Engineering examples:** Whether a bit is 0 or 1

- **Engineering examples:** Whether a bit is 0 or 1, whether a bit is in error

- **Engineering examples:** Whether a bit is 0 or 1, whether a bit is in error, whether a component has failed, whether something has been detected

- **Engineering examples:** Whether a bit is 0 or 1, whether a bit is in error, whether a component has failed, whether something has been detected
- **Examples using Jupyter Notebook**

## 2. Binomial RV

## 2. Binomial RV

- A Binomial RV represents the number of successes on  $n$  independent Bernoulli trials

## 2. Binomial RV

- A Binomial RV represents the number of successes on  $n$  independent Bernoulli trials
- Thus, a Binomial RV can also be defined as the sum of  $n$  independent Bernoullis RVs

## 2. Binomial RV

- A Binomial RV represents the number of successes on  $n$  independent Bernoulli trials
- Thus, a Binomial RV can also be defined as the sum of  $n$  independent Bernoullis RVs
- Let  $X = \#$  of successes



## 2. Binomial RV

- A Binomial RV represents the number of successes on  $n$  independent Bernoulli trials
- Thus, a Binomial RV can also be defined as the sum of  $n$  independent Bernoullis RVs
- Let  $X = \#$  of successes  
Then the pmf of  $X$  is given by

## 2. Binomial RV

- A Binomial RV represents the number of successes on  $n$  independent Bernoulli trials
- Thus, a Binomial RV can also be defined as the sum of  $n$  independent Bernoullis RVs
- Let  $X = \#$  of successes

Then the pmf of  $X$  is given by

$$P[X = k] = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k}, & k = 0, 1, \dots, n \\ 0, & \text{o.w.} \end{cases}$$

- **Engineering examples:** The number of bits in error in a packet

- **Engineering examples:** The number of bits in error in a packet, the number of defective items in a manufacturing run

- **Engineering examples:** The number of bits in error in a packet, the number of defective items in a manufacturing run
- **Examples using Jupyter Notebook**

### 3. Geometric RV

### 3. Geometric RV

- A Geometric RV occurs when independent Bernoulli trials are conducted until the first success

### 3. Geometric RV

- A Geometric RV occurs when independent Bernoulli trials are conducted until the first success
- $X = \#$  number of trials required



### 3. Geometric RV

- A Geometric RV occurs when independent Bernoulli trials are conducted until the first success
- $X = \#$  number of trials required  
Then the pmf of  $X$  is given by

$$P[X = k] = \begin{cases} (1 - p)^{k-1} p, & k = 1, 2, \dots \\ 0, & \text{o.w.} \end{cases}$$

- **Engineering examples:** The number of retransmissions required for a packet

- **Engineering examples:** The number of retransmissions required for a packet, number of white dots between black dots in the scan of a black and white document

- **Engineering examples:** The number of retransmissions required for a packet, number of white dots between black dots in the scan of a black and white document
- **Examples using Jupyter Notebook**

## 4. Poisson RV

## 4. Poisson RV

- Models events that occur randomly in space or time

## 4. Poisson RV

- Models events that occur randomly in space or time
- Let  $\lambda$  = the # of events/(unit of space or time)

## 4. Poisson RV

- Models events that occur randomly in space or time
- Let  $\lambda$  = the # of events/(unit of space or time)
- Consider observing some period of time or space of length  $t$  and let  $\alpha = \lambda t$



## 4. Poisson RV

- Models events that occur randomly in space or time
- Let  $\lambda$  = the # of events/(unit of space or time)
- Consider observing some period of time or space of length  $t$  and let  $\alpha = \lambda t$
- Let  $N$  = the # events in time (or space)  $t$

## 4. Poisson RV

- Models events that occur randomly in space or time
- Let  $\lambda$  = the # of events/(unit of space or time)
- Consider observing some period of time or space of length  $t$  and let  $\alpha = \lambda t$
- Let  $N$  = the # events in time (or space)  $t$
- The pmf of the Poission random variable is

$$P_N(n) = \begin{cases} \frac{\alpha^n}{n!} e^{-\alpha}, & n = 0, 1, \dots \\ 0, & \text{o.w.} \end{cases}$$

- For large  $\alpha$ , the Poisson pmf has a bell shape. For example, see the pmf when  $\alpha = 20$

- For large  $\alpha$ , the Poisson pmf has a bell shape. For example, see the pmf when  $\alpha = 20$
- **Engineering examples:**
  - calls coming in to a switching center

- For large  $\alpha$ , the Poisson pmf has a bell shape. For example, see the pmf when  $\alpha = 20$
- **Engineering examples:**
  - calls coming in to a switching center
  - packets arriving at a queue in a network

- For large  $\alpha$ , the Poisson pmf has a bell shape. For example, see the pmf when  $\alpha = 20$
- **Engineering examples:**
  - calls coming in to a switching center
  - packets arriving at a queue in a network
  - processes being submitted to a scheduler

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book



*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book
- # of people in a community that live to be 100 years old

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book
- # of people in a community that live to be 100 years old
- # of wrong telephone numbers that are dialed in a day

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book
- # of people in a community that live to be 100 years old
- # of wrong telephone numbers that are dialed in a day
- # of  $\alpha$ -particles discharged in a fixed period of time from some radioactive material

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book
- # of people in a community that live to be 100 years old
- # of wrong telephone numbers that are dialed in a day
- # of  $\alpha$ -particles discharged in a fixed period of time from some radioactive material
- # of earthquakes per year

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book
- # of people in a community that live to be 100 years old
- # of wrong telephone numbers that are dialed in a day
- # of  $\alpha$ -particles discharged in a fixed period of time from some radioactive material
- # of earthquakes per year
- # of computer crashes in a lab in a week

*The following examples are adopted from A First Course in Probability by Sheldon Ross:*

- # of misprints on a group of pages in a book
- # of people in a community that live to be 100 years old
- # of wrong telephone numbers that are dialed in a day
- # of  $\alpha$ -particles discharged in a fixed period of time from some radioactive material
- # of earthquakes per year
- # of computer crashes in a lab in a week
- **Examples using Jupyter Notebook**

# EEL 4930 Stats – Lecture 21

## **PROPERTIES OF DISTRIBUTION FUNCTIONS**

1.  $0 \leq F_X(x) \leq 1$

# EEL 4930 Stats – Lecture 21

## PROPERTIES OF DISTRIBUTION FUNCTIONS

1.  $0 \leq F_X(x) \leq 1$

Pf:  $F_X(x)$  is a prob. measure



2.  $F_X(-\infty) = 0$  and  $F_X(\infty) = 1$

2.  $F_X(-\infty) = 0$  and  $F_X(\infty) = 1$

Proof is technical.

2.  $F_X(-\infty) = 0$  and  $F_X(\infty) = 1$

Proof is technical.

Basically,  $F_X(-\infty)$  and  $F_X(\infty)$  are defined as limits, and the corresponding subsets of the samples space  $\{s \in S : X \leq x\}$  are either shrinking to  $\emptyset$  or  $S$

3.  $F_X(x)$  is monotonically nondecreasing,

3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

$$\text{Pf: } P\{X \in (-\infty, b]\}$$

3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

$$\text{Pf: } P\{X \in (-\infty, b]\} = P(X \in (-\infty, a])$$

3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

$$\begin{aligned} \text{Pf: } P\{X \in (-\infty, b]\} &= P(X \in (-\infty, a]) \\ &\quad + P(X \in (a, b]) \end{aligned}$$



3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

$$\begin{aligned} \text{Pf: } P\{X \in (-\infty, b]\} &= P(X \in (-\infty, a]) \\ &\quad + P(X \in (a, b]) \end{aligned}$$

$$\Rightarrow F_X(b) = F_X(a) + P(a < X \leq b)$$

3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

$$\begin{aligned} \text{Pf: } P\{X \in (-\infty, b]\} &= P(X \in (-\infty, a]) \\ &\quad + P(X \in (a, b]) \end{aligned}$$

$$\Rightarrow F_X(b) = F_X(a) + P(a < X \leq b) \quad (1)$$

$$4. P(a < X \leq b) = F_X(b) - F_X(a)$$

3.  $F_X(x)$  is monotonically nondecreasing,  
i.e.,  $F_X(a) \leq F_X(b)$  iff  $a \leq b$

$$\begin{aligned} \text{Pf: } P\{X \in (-\infty, b]\} &= P(X \in (-\infty, a]) \\ &\quad + P(X \in (a, b]) \end{aligned}$$

$$\Rightarrow F_X(b) = F_X(a) + P(a < X \leq b) \quad (1)$$

4.  $P(a < X \leq b) = F_X(b) - F_X(a)$

Pf: rewriting equation (1)

5.  $F_X(x)$  is continuous on the right,

5.  $F_X(x)$  is continuous on the right,  
i.e.,  $F_X(b) = \lim_{h \rightarrow 0} F_X(b + h) = F_X(b)$

5.  $F_X(x)$  is continuous on the right,  
i.e.,  $F_X(b) = \lim_{h \rightarrow 0} F_X(b + h) = F_X(b)$

*(The value at a jump discontinuity is the value after the jump.)*

5.  $F_X(x)$  is continuous on the right,  
i.e.,  $F_X(b) = \lim_{h \rightarrow 0} F_X(b + h) = F_X(b)$

*(The value at a jump discontinuity is the value after the jump.)*

**Proof is rather technical and will be omitted.**

6.  $P(X > x) = 1 - F_X(x)$



$$6. P(X > x) = 1 - F_X(x)$$

$$\text{Pf: } \{X > x\} = \{X \leq x\}^c$$

$$6. P(X > x) = 1 - F_X(x)$$

$$\begin{aligned} \text{Pf: } \{X > x\} &= \{X \leq x\}^c \\ \Rightarrow P(X > x) &= 1 - P(X \leq x) \end{aligned}$$

$$6. P(X > x) = 1 - F_X(x)$$

$$\text{Pf: } \{X > x\} = \{X \leq x\}^c$$

$$\Rightarrow P(X > x) = 1 - P(X \leq x) = 1 - F_X(x)$$