




Phase is shown in radians; amplitude shows relative object amplitude. Recon n is the displayed acquisition; train n is the recorded training count. High half selects checkpoints (spatial validation, not an independent test; no guard band). Reference is an experimental estimate; dashed contour marks scored interior. Prior fragments: Vong et al., Contrast-invariant deep ptychography neural networks, arXiv:2608.02869, synthetic-only whole-field crops, cropped and rescaled (CC BY 4.0). Their original scales are shown in-cell; the shared computed-method bar excludes prior fragments. Raster registration and zoom are unknown. TP2 probe/reference-histogram provenance applies only to its Velociprobe family. Whole-acquisition images include the training region.



Phase is shown in radians; amplitude shows relative object amplitude. Recon n is the displayed acquisition; train n is the recorded training count. High half selects checkpoints (spatial validation, not an independent test; no guard band). Reference is an experimental estimate; dashed contour marks scored interior. Prior fragments: Vong et al., Contrast-invariant deep ptychography neural networks, arXiv:2608.02869, synthetic-only whole-field crops, cropped and rescaled (CC BY 4.0). Their original scales are shown in-cell; the shared computed-method bar excludes prior fragments. Raster registration and zoom are unknown. TP2 probe/reference-histogram provenance applies only to its Velociprobe family. Whole-acquisition images include the training region.



Phase is shown in radians; amplitude shows relative object amplitude. Recon n is the displayed acquisition; train n is the recorded training count. High half selects checkpoints (spatial validation, not an independent test; no guard band). Reference is an experimental estimate; dashed contour marks scored interior. Prior fragments: Vong et al., Contrast-invariant deep ptychography neural networks, arXiv:2608.02869, synthetic-only whole-field crops, cropped and rescaled (CC BY 4.0). Their original scales are shown in-cell; the shared computed-method bar excludes prior fragments. Raster registration and zoom are unknown. TP2 probe/reference-histogram provenance applies only to its Velociprobe family. Whole-acquisition images include the training region.