

**IT University of Copenhagen**  
**Foundations of Computing—Discrete Mathematics BSc**  
**Midterm Assignment**

October 8, 2019

**Instructions** (*read carefully*)

**Contents:** The midterm contains 8 questions for the total of 19 points. The midterm is divided into two parts: The first part has 4 multiple choice questions and the second part has 4 open ended questions (some of the open ended questions have subquestions).

**What to check:** In the multiple choice questions, there is one and only one correct answer. You should only check 1 box.

**Definitions and theorems:** At the end of this document (page 5) you can find some definitions and theorems that could be useful for answering some of the questions.

**Info about you:** Write *clearly* your *full name*, your date of birth (DoB) and the room that you normally go to exercises in (i.e. 2A52, 2A54, or 4A16) on every page (top-right) including the front page.

**—IMPORTANT—**

*Only information written on the pages 1–4 will be evaluated.  
Anything else that you hand-in will NOT be considered for the final evaluation!*

**Part I.** Answer the following multiple choice questions.

1. (2 pts) Which of the following statements is *true*?

☐  $\{(2, 4), (2, 6), (2, 8), (3, 6), (3, 9), (4, 8)\} \subset | \cap (\{2 \dots 9\} \times \{2 \dots 9\})$   
where  $|$  is the “divides” relation

☐  $\mathcal{P}(A \cup B) \subseteq \mathcal{P}(A) \cup \mathcal{P}(B)$  for arbitrary sets  $A, B$

☐  $\{3, \sqrt{10}, 24 \bmod 7\} \subseteq \{8 \bmod 5\}$

☐  $(A - C) \cap (B - C) = \emptyset$  for arbitrary sets  $A, B, C$

2. (2 pts) Which one is a valid representation of  $(E74)_{16}$ ?

☐  $(3900)_{10}$

☐  $(111001110100)_2$

☐  $(7174)_8$

☐  $(11011011)_2$

3. (2 pts) Let  $a_k = a_{k-1} + 2a_{k-2}$ ,  $a_0 = -1$ ,  $a_1 = 2$ . Which of the following statements is *true*?


☐  $a_0 = a_3$

☐  $a_k + a_{k+1} = a_k - a_{k-1}$  for all  $k$

☐  $a_k \cdot a_{k+1} = a_k^2 + 2 \cdot a_k \cdot a_{k-1}$  for all  $k$

☐  $\sum_{i=1}^4 a_k = a_5$

4. (2 pts) Consider the following directed graph representing the relation  $R$  on the set  $\{a, b, c, d, e\}$ .



Which of the following statements is *true*?

- ☐  $R$  is antisymmetric and transitive, but not reflexive
- ☐  $R$  is transitive, but neither reflexive nor antisymmetric
- ☐  $R$  is antisymmetric, but neither reflexive nor transitive
- ☐  $R$  is reflexive and antisymmetric and transitive

**Part II.** Answer the following questions. Be brief but precise, your correct use of mathematical notation is an important aspect of your answer.

5. (a) (3 pts) Construct a valid judgment (derivation) for the following predicate:

$$\exists x(\neg Q(x)) \rightarrow (\forall y(P(y) \rightarrow Q(y))) \rightarrow \neg \forall z(P(z))$$

- (b) (1 pt) Is this predicate valid in intuitionistic logic, classical logic, or both? Explain your answer.

6. (3 pts) Assume that  $A$  and  $B$  are non-empty sets and that  $f : A \rightarrow B$  is a bijective function. Let  $g : A \rightarrow \mathcal{P}(B)$  be the function defined as  $g(a) = \{f(a)\}$  for all  $a \in A$ . One of the following statements is true, and one is false.

1.  $g$  is injective.
2.  $g$  is surjective.

Write which statement is true and prove it. Write which statement is false and find a counter example. (Just answering which is true and which is false gives no points.)

7. (2 pts) Let  $A$ ,  $B$  and  $C$  be sets. Show that if  $A \subseteq B$  and  $B \subseteq \overline{C}$ , then  $A \cap C = \emptyset$ .

8. (2 pts) Let  $a, b \in \mathbb{Z}$  be integers, and assume that  $a \mid b$ . Prove that  $a \mid 5b + a$ .

## Definitions and theorems

### Logic

The logic rules for propositional and predicate logics are given below.

#### Conjunction:

$$\frac{A \text{ true} \quad B \text{ true}}{A \wedge B \text{ true}} \wedge I \quad \frac{A \wedge B \text{ true}}{A \text{ true}} \wedge E_1 \quad \frac{A \wedge B \text{ true}}{B \text{ true}} \wedge E_2$$

#### Implication:

$$\frac{\overline{A \text{ true}}^u \quad \vdots \quad B \text{ true}}{A \rightarrow B \text{ true}} \rightarrow I^u \quad \frac{A \rightarrow B \text{ true} \quad A \text{ true}}{B \text{ true}} \rightarrow E$$

#### Disjunction:

$$\frac{A \text{ true}}{A \vee B \text{ true}} \vee I_1 \quad \frac{B \text{ true}}{A \vee B \text{ true}} \vee I_2 \quad \frac{A \vee B \text{ true} \quad \overline{A \text{ true}}^u \quad \vdots \quad C \text{ true} \quad \overline{B \text{ true}}^v \quad \vdots \quad C \text{ true}}{C \text{ true}} \vee E^{u,v}$$

#### True and false:

$$\overline{\mathbf{T} \text{ true}} \mathbf{T}I \quad \frac{\mathbf{F} \text{ true}}{C \text{ true}} \mathbf{F}E$$

#### Negation:

$$\frac{\overline{A \text{ true}}^u \quad \vdots \quad \mathbf{F} \text{ true}}{\neg A \text{ true}} \neg I^u \quad \frac{\neg A \text{ true} \quad A \text{ true}}{C \text{ true}} \neg E$$

#### Classical rules:

$$\overline{A \vee \neg A \text{ true}} LEM \quad \frac{\neg \neg A \text{ true}}{A \text{ true}} \neg \neg C \quad \frac{\overline{\neg A \text{ true}}^u \quad \vdots \quad \mathbf{F} \text{ true}}{A \text{ true}} \mathbf{F}_C^u$$

#### Quantifiers:

$$\frac{A[a/x] \text{ true}}{\forall x(A) \text{ true}} \forall I^a \quad \frac{\forall x(A) \text{ true}}{A[t/x] \text{ true}} \forall E \quad \frac{\overline{A[a/x] \text{ true}}^u \quad \vdots \quad C \text{ true}}{\exists x(A) \text{ true}} \exists E^{a,u} \quad \frac{A[t/x] \text{ true}}{\exists x(A) \text{ true}} \exists I$$

## Sets

**Size of a set**  $|A|$  denotes the number of elements of  $A$

**Emptyset**  $\forall x(x \notin \emptyset)$

**Equality**  $A = B$  iff  $\forall x(x \in A \leftrightarrow x \in B)$

**Subset**  $A \subseteq B$  iff  $\forall x(x \in A \rightarrow x \in B)$

**Union**  $A \cup B = \{x \mid x \in A \vee x \in B\}$  Property:  $\forall x(x \in A \cup B \leftrightarrow x \in A \vee x \in B)$

**Intersection**  $A \cap B = \{x \mid x \in A \wedge x \in B\}$  Property:  $\forall x(x \in A \cap B \leftrightarrow x \in A \wedge x \in B)$

**Difference**  $A - B = \{x \mid x \in A \wedge x \notin B\}$  Property:  $\forall x(x \in A - B \leftrightarrow x \in A \wedge x \notin B)$

**Complement**  $\bar{A} = U - A$  Property:  $\forall x(x \in \bar{A} \leftrightarrow x \in U \wedge x \notin A)$

**Cartesian product**  $A \times B = \{(a, b) \mid a \in A \wedge b \in B\}$

## Relations

**Binary relation** from  $A$  to  $B$  is a subset of  $A \times B$

**Relation on a set**  $A$  is a relation from  $A$  to  $A$  (subset of  $A \times A$ )

**Reflexive relation**  $R$  on set  $A$  satisfies  $\forall x \in A((x, x) \in R)$

Digraphs: loops on every vertex

**Symmetric relation** satisfies  $\forall x, y \in A((x, y) \in R \leftrightarrow (y, x) \in R)$

Digraphs: every edge has the corresponding edge in the opposite direction

**Antisymmetric relation** satisfies  $\forall x, y \in A((x, y) \in R \wedge (y, x) \in R \rightarrow x = y)$

Digraphs: never two edges in the opposite direction

**Transitive relation** satisfies  $\forall x, y, z \in A((x, y) \in R \wedge (y, z) \in R \rightarrow (x, z) \in R)$

Digraphs: an edge  $a \rightarrow b$  and an edge  $b \rightarrow c$  implies an edge  $a \rightarrow c$

**Composing relations**  $S \circ R = \{(x, y) \mid (x, z) \in R \wedge (z, y) \in S\}$

Property:  $\forall x, y((x, y) \in S \circ R \leftrightarrow \exists z((x, z) \in R \wedge (z, y) \in S))$

**Partial ordering**  $R$  on  $S$  iff  $R$  is reflexive, antisymmetric and transitive.

**Poset**  $(S, \preceq)$  is a poset iff  $\preceq$  is a partial ordering on  $S$ .

**Comparable**  $a$  and  $b$  are comparable iff  $a \preceq b$  or  $b \preceq a$

**Total order**  $(S, \preceq)$  iff all elements are comparable

**Maximal element**  $a$  is maximal iff  $\neg \exists x \in S(a \prec x)$

**Minimal element**  $a$  is minimal iff  $\neg \exists x \in S(x \prec a)$

**Greatest element**  $a$  is greatest iff  $\forall x \in S(x \preceq a)$

**Least element**  $a$  is least iff  $\forall x \in S(a \preceq x)$

**Upper bound**  $u$  of  $A$  iff  $\forall x \in A(x \preceq u)$

**Lower bound**  $l$  of  $A$  iff  $\forall x \in A(l \preceq x)$

## Functions

**Function**  $f : A \rightarrow B$  assigns exactly one element of  $B$  to each element of  $A$ .  
 $A$  is the **domain** and of  $f$   $B$  is the **codomain** of  $f$ .

**Image, preimage** Let  $f(a) = b$ , then  $b$  is the **image** and  $a$  is the **preimage**.

**Image of a set**  $S \subseteq A$ :  $f(S) = \{y \in B \mid x \in S, y = f(x)\}$

**One-to-one/injective function**  $\forall x, y \in A(f(x) = f(y) \rightarrow x = y)$

**Onto/surjective function**  $\forall y \in B \exists x \in A(f(x) = y)$

**One-to-one correspondence / bijection** is both one-to-one and onto.

**Function composition**  $\forall x((f \circ g)(x) = f(g(x)))$   
 $\forall x, y((f \circ g)(x) = y \leftrightarrow \exists z(f(x) = z \wedge g(z) = y))$

## Sequences and Summations

**Geometric progression** A sequence of the form  $a, ar, ar^2, \dots, ar^n, \dots$ , where  $a \in \mathbb{R}$  is the *initial term* and  $r \in \mathbb{R}$  is the *common ratio*.

**Arithmetic progression** A sequence of the form  $a, a + d, a + 2d, \dots, a + nd, \dots$ , where  $a \in \mathbb{R}$  is the *initial term* and  $d \in \mathbb{R}$  is the *common difference*.

### Some Useful Summation Formulae

Sum	Closed form	Sum	Closed form
$\sum_{k=0}^n ar^k (r \neq 0)$	$\frac{ar^{n+1} - a}{r - 1}, r \neq 1$	$\sum_{k=1}^n k$	$\frac{n(n+1)}{2}$
$\sum_{k=1}^n k^2$	$\frac{n(n+1)(2n+1)}{6}$	$\sum_{k=1}^n k^3$	$\frac{n^2(n+1)^2}{4}$
$\sum_{k=1}^{\infty} x^k,  x  < 1$	$\frac{1}{1-x}$	$\sum_{k=1}^{\infty} kx^{k-1},  x  < 1$	$\frac{1}{(1-x)^2}$



## Number Theory

Given two integers  $a$  and  $b$ , with  $a \neq 0$ , we say that  $a$  *divides*  $b$  if there exist an integer  $c$  such that  $b = ac$ , or equivalently, if  $\frac{b}{a}$  is an integer. If  $a$  divides  $b$  then  $a$  is a *factor* (or *divisor*) of  $b$ , and  $b$  is said to be a *multiple* of  $a$ .

**Theorem 1** (Division algorithm). *Let  $a$  be an integer and  $d$  a positive integer. Then there exist unique integers  $q$  and  $r$ , with  $0 \leq r < d$ , such that  $a = dq + r$ .*

In theorem 1 the value  $d$  is called the *divisor*,  $a$  is the *dividend*,  $q$  is the *quotient*, and  $r$  is the *remainder*. Then  $q = a \text{ div } d$ ,  $r = a \text{ mod } d$ . Remember that the remainder cannot be negative.

If  $a$  and  $b$  are integers and  $m$  is a positive integer, then  $a$  is *congruent to  $b$  modulo  $m$*  if  $m \mid a - b$  and we write  $a \equiv b \pmod{m}$ .

The *greatest common divisor* of two integers  $a$  and  $b$ , not both zero, is denoted by  $\gcd(a, b)$  and is the largest integer that both divides  $a$  and divides  $b$ .

The *Euclidean algorithm* provides a efficient way to compute the greatest common divisor of two integers. The algorithm is based on the following lemma.

**Lemma 1.** *Let  $a = bq + r$  where  $a$ ,  $b$ ,  $q$  and  $r$  are integers. Then  $\gcd(a, b) = \gcd(b, r)$ .*

Hexadecimal, octal and binary representation of integers 0 through 15																
Decimal	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
Hexadecimal	0	1	2	3	4	5	6	7	8	9	A	B	C	D	E	F
Octal	0	1	2	3	4	5	6	7	10	11	12	13	14	15	16	17
Binary	0	1	10	11	100	101	110	111	1000	1001	1010	1011	1100	1101	1110	1111

## Graph Theory

A *graph*  $G = (V, E)$  consists of  $V$ , a nonempty set of *vertices* (or *nodes*) and  $E$ , a set of *edges*. Each edge has either one or two vertices associated with it, called its *endpoints*. An edge is said to *connect* its endpoints.

**Theorem 2** (Handshaking theorem). *Let  $G = (V, E)$  be an undirected graph with  $m$  edges, then*

$$2m = \sum_{v \in V} \deg(v)$$

*Note that this applies even if multiple edges and loops are present.*

**Theorem 3.** *Let  $G = (V, E)$  be a graph with directed edges, then  $\sum_{v \in V} \deg^-(v) = \sum_{v \in V} \deg^+(v) = |E|$ .*

<b>Graph terminology</b>			
<i>Type</i>	<i>Edges</i>	<i>Multiple edges are allowed?</i>	<i>Loops are allowed?</i>
Simple graph	Undirected	No	No
Multigraph	Undirected	Yes	No
Pseudograph	Undirected	Yes	Yes
Simple directed graph	Directed	No	No
Directed multigraph	Directed	Yes	Yes
Mixed graph	Directed and undirected	Yes	Yes