

Exact dynamics of Eratosthenes sieve beyond $2p$: coincident fusions, births of gaps and missing differences

Tomasz Hojka

Independent researcher

Preprint — October 2026

DOI: 10.5281/zenodo.23007183

Abstract

At each stage of Eratosthenes sieve the integers coprime to a modulus N form a periodic cycle of gaps $\mathcal{G}(N)$. F. B. Holt showed that the populations of gaps and constellations, together with their driving terms, evolve by an exact linear system as long as the span is smaller than twice the next prime q . Beyond that threshold several points of one driving term may be removed in the same image; Holt showed that this happens exactly when q divides their distance, but the lengths of the resulting terms were not determined. We give the exact transition law for all spans: by this criterion, applied to all points of a driving term including the interior ones, the points split into residue classes modulo q ; each class is removed in exactly one of the q images, and this determines every image. Consequences include a ladder of thresholds (classes of 2, 3, 4, 5 points appear from spans $2q, 6q, 8q, 12q$), a form in binomial moments with nonnegative sources, an exact birth law for gaps and constellations, a criterion for the first appearance of a constellation, and a sufficient condition for the exactness of Holt's Markov chain. As applications we extend Holt's exact models to gaps $84 \leq g \leq 104$, reproduce the pre-linear rows of his worked example, classify all 20 missing differences in Ziller's table up to $k = 30$ (all are delays; 17 holes and 3 shadows; none is caused by a blocked birth), and describe how the maximal gaps of $\mathcal{G}(p_n\#)$ arise for $n \leq 64$. All theorems are also verified computationally; code and data accompany the paper.

1 Introduction

For a prime p let $p\#$ denote the product of all primes up to p . The integers coprime to $p\#$ — the candidates that survive Eratosthenes sieve up to p — form a periodic set; the sequence of differences between consecutive elements over one period is the *cycle of gaps* $\mathcal{G}(p\#)$, of length $\varphi(p\#)$ and span $p\#$. Holt and Rudd [1] observed that $\mathcal{G}(p_{k+1}\#)$ arises from $\mathcal{G}(p_k\#)$ by a simple recursion: concatenate p_{k+1} copies of the cycle and fuse adjacent gaps at the positions of the multiples of p_{k+1} . In a series of papers [1, 2, 3, 4] Holt developed this into a discrete dynamical system. A central notion is that of a *driving term*: a run of consecutive gaps whose sum equals a given gap g (or which contains a given constellation s), and which can turn into g (or into s) at a later stage by fusions. The populations $n_{g,j}(p\#)$ of driving terms of span g and length j obey the linear recursion

$$n_{g,j}(qN) = (q - j - 1) n_{g,j}(N) + j n_{g,j+1}(N)$$

with explicit eigenstructure, which yields exact models for the relative populations of gaps and constellations at all later stages of the sieve.

This recursion holds under a restriction: the span must be smaller than twice the next prime. For larger spans two fusions can fall into the same image of a driving term, and Holt

characterized when this happens: two fusions fall into the same image if and only if the next prime divides their distance [3, Lemma 3.1]. Holt notes that the minimum distance between fusions is $2p_{k+1}$ [4, Obs. 2.3], that for spans below $2p_1$ all fusions of a constellation fall into different images [4, Lemma 2.4], and that for gaps beyond $2p_1$ the counts of driving terms of various lengths could not be determined [1, 2]; the single case $g = 2p_1$ was handled by a special first iteration [2]. Several fusions at the *boundary* of one image are accounted for in his aggregate count through the number $\nu_s(p)$ of residue classes covered by the boundary points [4, Lemma 2.5, Lemma 2.7]. What remained open is the case of several *interior* fusions in the same image, which changes the distribution of lengths.

Main result. We show that the transition from $\mathcal{G}(N)$ to $\mathcal{G}(qN)$ is determined exactly, for every span, by the partition of the points of a driving term into residue classes modulo q : each class is removed in exactly one of the q images, and the remaining images are untouched (Theorem 3.4). The partition into classes is Holt’s criterion [3, Lemma 3.1] applied to all points of a driving term, interior points included; what is new is that this partition determines the lengths of all images, together with the consequences drawn from it. Holt’s recursion is the special case in which all classes are singletons.

Consequences. Section 3 derives a ladder of thresholds for the size of the classes (Corollary 3.6) and a form of the dynamics in binomial moments, in which the nonlinearity appears only as a nonnegative source (Theorem 3.7). Section 4 gives an exact birth law for gaps and constellations (Theorem 4.1), a criterion for the first appearance of a constellation (Corollary 4.3), and the description of the set of gap lengths $\mathcal{D}(qN)$ in terms of $\mathcal{G}(N)$. Section 5 identifies the exact process on shapes of driving terms and gives a sufficient condition for Holt’s Markov chain on lengths to be exact (Theorem 5.1).

Applications (part 2 of this draft). We extend Holt’s exact models from gaps $g \leq 82$ to $g \leq 104$, which includes the gaps $84 \leq g \leq 90$ named in [4] as a current goal; reproduce the rows of Holt’s worked example [4, Fig. 6] that lie outside the linear model; classify all missing differences in Ziller’s table [5] up to $k = 30$; and describe how the maximal gaps of $\mathcal{G}(p_n\#)$ — the values of Jacobsthal’s function for primorials, OEIS A048670 — arise from the previous cycle for $n \leq 64$.

What is not claimed. All results concern the cycles $\mathcal{G}(N)$; nothing is claimed about the actual gaps between primes. Ziller’s Conjecture 4.1 [5] is reformulated, not proved. The condition in Theorem 5.1 is sufficient; its necessity is only observed. Statements marked as computed or observed rest on computation and are labelled as such.

2 Setting and notation

Throughout, N is a squarefree integer with $2 \mid N$, and q is an odd prime with $q \nmid N$. The main case is $N = p_k\#$ and $q = p_{k+1}$. Let $\Omega(N) = \{x \in \mathbb{Z} : \gcd(x, N) = 1\}$, a set periodic modulo N . The cycle of gaps $\mathcal{G}(N)$ is the sequence of differences of consecutive elements of $\Omega(N)$ over one period; $\mathcal{D}(N)$ is the set of gap lengths occurring in $\mathcal{G}(N)$ and

$$H(N) := \max \mathcal{D}(N)$$

is the largest gap, the value of Jacobsthal’s function at N . For primorials we write $\mathcal{D}(k) := \mathcal{D}(p_k\#)$ and $h(k) := H(p_k\#)$, as in [5].

Definition 2.1 (Constellations and driving terms). A *constellation* is a finite sequence $s = (s_1, \dots, s_m)$ of even positive integers with span $|s| = s_1 + \dots + s_m$; its points are $C = \{c_0 = 0 < c_1 < \dots < c_m = |s|\}$, $c_i = s_1 + \dots + s_i$. We assume $0 < |s| < N$. A *driving term* for s in $\mathcal{G}(N)$ is a residue $x \bmod N$ with $x + c \in \Omega(N)$ for all $c \in C$. Its point set is

$$P(x) = \{t \in \{0, 1, \dots, |s|\} : x + t \in \Omega(N)\} \supseteq C,$$

its *length* is $J = |P(x)| - 1$ (the number of gaps of $\mathcal{G}(N)$ it spans), and $E(x) = P(x) \setminus C$ is the set of its *extra points*. We write $n_{s,J}(N)$ for the number of driving terms for s of length J ; the occurrences of s itself are those with $J = m$. For a single gap ($m = 1$, $s = (g)$) we write $n_{g,J}(N)$.

The condition $|s| < N$ is a convention ensuring that a driving term lies within one period of the cycle; none of the proofs below uses it (Remark 4.5). In Holt's terms, a driving term for s becomes s through *interior fusions* (removal of extra points), while a *boundary fusion* (removal of a point of C) eliminates it as a driving term for s [4, §2.1].

Images and classes. For $x \bmod N$ the q residues $X \bmod qN$ with $X \equiv x \pmod{N}$ are $X_i = x + iN$, $0 \leq i < q$; we call them the *images* of x . For a driving term x we partition $P(x)$ into classes of congruence modulo q and denote the set of classes by $\mathcal{C}(x)$. For a finite set A , $\nu_q(A)$ denotes the number of residue classes modulo q occupied by A ; thus $|\mathcal{C}(x)| = \nu_q(P(x))$, and $\nu_q(C)$ is Holt's $\nu_s(q)$.

3 The transition law

Lemma 3.1. For $t, t' \in P(x)$ we have $t \equiv t' \pmod{q}$ if and only if $2q \mid t - t'$.

Proof. Since $2 \mid N$, all elements of $\Omega(N)$ are odd, so $t - t' = (x + t) - (x + t')$ is even; for even d and odd prime q , $q \mid d \iff 2q \mid d$. $\square$

This is the quantitative content of Holt's observation that the minimum distance between fusions is $2p_{k+1}$ [4, Obs. 2.3].

Lemma 3.2. The images $X_0, \dots, X_{q-1}$ of x run through all residues modulo q , each exactly once.

Proof. $\gcd(N, q) = 1$, so $i \mapsto iN \bmod q$ is a bijection of $\mathbb{Z}/q$. $\square$

Lemma 3.2 is [3, Lemma 4.1], [4, Lemma 2.6]. It gives Holt's criterion [3, Lemma 3.1]: two points of a driving term are removed in the same image if and only if q divides their distance, or equivalently, by Lemma 3.1, $2q$ divides it. Holt applies it to the boundary points, to count admissible instances; Theorem 3.4 applies it to all points.

Lemma 3.3. If $X \equiv x \pmod{N}$, then $X + t \in \Omega(qN)$ if and only if $x + t \in \Omega(N)$ and $q \nmid X + t$.

Proof. $\gcd(X + t, qN) = 1$ iff $\gcd(X + t, N) = 1$ and $q \nmid X + t$, and $X + t \equiv x + t \pmod{N}$. $\square$

Theorem 3.4 (Transition law). Let x be a driving term for s of length J in $\mathcal{G}(N)$, with classes $\mathcal{C} = \mathcal{C}(x)$. Among the q images of x :

1. for each class $K \in \mathcal{C}$ there is exactly one image X_K in which $q \mid X_K + t$ for all $t \in K$ and for no $t \in P(x) \setminus K$;
2. the remaining $q - |\mathcal{C}|$ images contain no point divisible by q .

Consequently the images of type 2 are driving terms for s of length J in $\mathcal{G}(qN)$; the image X_K with $K \cap C = \emptyset$ is a driving term for s of length $J - |K|$; and the image X_K with $K \cap C \neq \emptyset$ is not a driving term for s . Every driving term for s in $\mathcal{G}(qN)$ arises in this way from exactly one driving term in $\mathcal{G}(N)$ and exactly one of its images. Hence

$$n_{s,J'}(qN) = \sum_x \left([J(x) = J'] (q - |\mathcal{C}(x)|) + \#\{K \in \mathcal{C}(x) : K \cap C = \emptyset, J(x) - |K| = J'\} \right), \quad (1)$$

the sum running over all driving terms for s in $\mathcal{G}(N)$.

Proof. $q \mid X + t$ iff $X \equiv -t \pmod{q}$. The set $\{-t \bmod q : t \in P(x)\}$ has exactly $|\mathcal{C}|$ elements, one for each class. By Lemma 3.2 each residue modulo q is taken by exactly one image; the image with residue $-t$ removes exactly the points of the class of t , and the other $q - |\mathcal{C}|$ images remove none. By Lemma 3.3 the points of $\Omega(qN)$ in $[X, X + |s|]$ are the points of $P(x)$ minus the removed class. If the removed class meets C , a point of s is lost; otherwise $|K|$ extra points are removed and the length drops by $|K|$. Conversely, if X is a driving term for s in $\mathcal{G}(qN)$, then by Lemma 3.3 $x = X \bmod N$ has $x + c \in \Omega(N)$ for all $c \in C$, so x is a driving term in $\mathcal{G}(N)$ and X is one of its images; x is determined by X . $\square$

Remark 3.5 (Holt's recursion). If no two points of $P(x)$ are congruent modulo q — by Lemma 3.1 this holds whenever $|s| < 2q$ — all classes are singletons, $|\mathcal{C}(x)| = J + 1$, and there are $J - m$ extra points. Then (1) becomes

$$n_{s,J}(qN) = (q - J - 1) n_{s,J}(N) + (J + 1 - m) n_{s,J+1}(N),$$

which for gaps ($m = 1$) is the recursion of Holt and Rudd [1, Thm. 3.2], and for constellations the recursion underlying [4, Fig. 7]. Theorem 3.4 determines the lengths also when several interior fusions fall into one image, the case left open in [1, 2].

Corollary 3.6 (Ladder of thresholds). *A class with $r \geq 2$ points has its extreme points at distance at least $2q(r - 1)$, a positive multiple of $2q$. If moreover $3 \mid N$ and $q \neq 3$, a class with three points spans at least $6q$.*

Proof. The first statement follows from Lemma 3.1. For the second, let $t_1 < t_2 < t_3$ lie in one class, $t_2 = t_1 + 2qa$, $t_3 = t_2 + 2qb$ with $a, b \geq 1$. If $a = b = 1$, the residues of $t_1, t_1 + 2q, t_1 + 4q$ modulo 3 are $t_1, t_1 - q, t_1 + q$, all three residues since $3 \nmid q$; one of the points would be divisible by 3, contradicting $3 \mid N$. Hence $a + b \geq 3$. $\square$

Hagedorn's bounds for the number of positions a prime can remove [6, Prop. 3.3] translate into further thresholds: in the presence of the primes 3 and 5 and for $q > 5$, classes of four points require span at least $8q$ and classes of five points at least $12q$. In all our tests (Section 7, part 2) the first classes of 2, 3, 4, 5 points occur exactly at the spans $2q, 6q, 8q, 12q$. In particular, for $3 \mid N$ and $|s| < 6q$ all classes have at most two points, at distance $2q$ or $4q$.

Theorem 3.7 (Moment form). *For $k \geq 0$ let $S_k(s, N) = \sum_J \binom{J-m}{k} n_{s,J}(N)$, the number of pairs (driving term, k -subset of its extra points), and for a finite set A let $W_N(A) = \#\{x \bmod N : x + a \in \Omega(N) \forall a \in A\}$. Then*

$$S_k(s, N) = \sum_{\substack{T \subseteq \{1, \dots, |s|-1\} \setminus C \\ |T|=k}} W_N(C \cup T), \quad S_k(s, qN) = (q - m - 1 - k) S_k(s, N) + \Pi_k(s, N),$$

where

$$\Pi_k(s, N) = \sum_{|T|=k} (m + 1 + k - \nu_q(C \cup T)) W_N(C \cup T) \geq 0.$$

Moreover $n_{s,m}(N) = \sum_{k \geq 0} (-1)^k S_k(s, N)$.

Proof. A pair (x, T) with $x + a \in \Omega(N)$ for $a \in C \cup T$ is exactly a driving term x together with a k -subset T of its extra points, which gives the first identity. By the Chinese remainder theorem $W_{qN}(A) = W_N(A)(q - \nu_q(A))$, since the condition modulo q excludes exactly $\nu_q(A)$ residues. Writing $q - \nu_q(A) = (q - |A|) + (|A| - \nu_q(A))$ with $|A| = m + 1 + k$ gives the recursion; $|A| - \nu_q(A) \geq 0$ counts coincidences modulo q . The last identity is binomial inversion, $\sum_k (-1)^k \binom{J-m}{k} = [J = m]$. $\square$

For $k = 0$ the theorem gives $S_0(s, qN) = (q - \nu_q(C)) S_0(s, N)$, the aggregate count of Holt [3, Cor. 1.2], [4, Thm. 1.1, Lemma 2.7] (for gaps [1, Cor. 5.3]). The coefficients $q - m - 1 - k$ are the eigenvalues of Holt's linear system and the S_k are its left-eigenvector coordinates [4, §3]; in these coordinates the nonlinearity is a nonnegative source, and decreases of a population can arise only through the alternating sum for $n_{s,m}$.

4 Births

Theorem 4.1 (Birth law).

$$n_{s,m}(qN) = (q - \nu_q(C)) n_{s,m}(N) + B_s^{(q)}(N),$$

where $B_s^{(q)}(N)$ is the number of driving terms for s in $\mathcal{G}(N)$ whose extra points form a single residue class modulo q that is disjoint from the classes of the points of C .

Proof. Apply Theorem 3.4 with target length m . Occurrences of s in $\mathcal{G}(N)$ have $P = C$ and keep $q - \nu_q(C)$ untouched images. A longer driving term yields an occurrence of s exactly when the removed class equals its whole set of extra points and contains no point of C , and then it yields exactly one. $\square$

For a gap $(s = (g))$ we write $e_g = [2q \mid g]$, so that $q - \nu_q(\{0, g\}) = q - 2 + e_g$. In the language of constellations of consecutive gaps $(a_1, \dots, a_j)$, $j \geq 2$, of $\mathcal{G}(N)$ with sum g , the condition of Theorem 4.1 reads: $2q \nmid a_1$, $2q \nmid a_j$, and $2q \mid a_i$ for $1 < i < j$ (Lemma 3.1). For $3 \mid N$ and $g < 6q$ only pairs (a, b) and triples (u, v, w) with $v \in \{2q, 4q\}$ occur. A pair with $2q \mid a$ or $2q \mid b$ does not produce a gap: we call this a *blocked birth*. A triple whose middle gap is a multiple of $2q$ and whose end gaps are not produces a gap of length $u + v + w$ in one image: a *coincident birth*.

Lemma 4.2 (Persistence). *If $\nu_q(C) < q$, then $n_{s,m}(qN) \geq n_{s,m}(N)$. If $\nu_q(C) = q$, then s occurs in no $\mathcal{G}(M)$ with $q \mid M$.*

Proof. The first part follows from Theorem 4.1; the second because every x then has a point $x + c$, $c \in C$, divisible by q . $\square$

Holt's Theorem 1.1 [3] shows that every admissible instance of s , driving terms included, arises and persists; the lemma is the corresponding statement for the occurrences of s itself, in substance Holt's Lemma 2.6 [4].

Corollary 4.3 (First appearance). *Let $N_0 \mid N_1 \mid \dots$ with $2 \mid N_0$ and $N_{i+1} = q_i N_i$ for odd primes $q_i \nmid N_i$. The constellation s occurs for the first time in this chain at N_{i+1} if and only if $n_{s,m}(N_i) = 0$ and $B_s^{(q_i)}(N_i) > 0$.*

Proof. If s occurs in $\mathcal{G}(N_{i+1})$ it is admissible modulo every prime dividing N_{i+1} ; had it occurred at some N_r with $r < i$, Lemma 4.2 applied to $q_r, \dots, q_{i-1}$ would give an occurrence in $\mathcal{G}(N_i)$. The converse follows from Theorem 4.1. $\square$

Corollary 4.4 (Calm regime). *If all gaps of $\mathcal{G}(N)$ are smaller than $2q$, then for every even g with $2 \leq g < N$*

$$n_{g,1}(qN) = (q - 2 + e_g) n_{g,1}(N) + n_{g,2}(N),$$

$$\mathcal{D}(qN) = \mathcal{D}(N) \cup \{a + b : (a, b) \text{ adjacent gaps of } \mathcal{G}(N)\}, \quad H(qN) = \max\{a + b\}.$$

Proof. No gap is a multiple of $2q$, so every pair qualifies and no constellation with a middle gap divisible by $2q$ exists. $\mathcal{D}(N) \subseteq \mathcal{D}(qN)$ because $q - 2 + e_g \geq 1$. $\square$

Remark 4.5 (Domain). The corollaries concern gaps up to $H(qN)$ while the convention of Section 2 requires spans below N . This is consistent: $H(qN) \leq N$ for every even squarefree N with an odd prime factor. Indeed, by Kanold's bound $H(M) \leq 2^{\omega(M)}$ [8]; if $\omega(N) \geq 3$ then $2^{\omega(N)+1} \leq N$; if $N = 2p$ with $p \geq 5$ then $H(qN) \leq 8 \leq 2p$; and for $N = 6$ the gaps of $\mathcal{G}(6)$ alternate 4, 2 and a prime $q \geq 5$ cannot remove two adjacent points, so $H(6q) = 6$. (For $N = 2$ one has $H(2q) = 4 > 2$.) The proofs of Theorems 3.4 and 4.1 do not use the convention at all, and our tests confirm the laws also for spans $\geq N$.

For primorials, the known values of Jacobsthal's function give $h(k) < 2p_{k+1}$ for all $k \leq 18$ ($h(18) = 132 < 134$), while $h(19) = 152 \geq 142 = 2p_{20}$. Hence for $k \leq 18$

$$\mathcal{D}(k+1) = \mathcal{D}(k) \cup \{\text{sums of two adjacent gaps of } \mathcal{G}(p_k\#)\}, \quad h(k+1) = \max\{\text{such sums}\},$$

and $k+1 = 20$ is the first stage at which blocked and coincident births are possible. Ziller observed that $h(k) > 2p_k$ for $k > 17$ [5, §4]; the condition relevant here compares $h(k)$ with $2p_{k+1}$.

Corollary 4.6 (Missing differences). *An even g with $2 \leq g < H(qN)$ is missing from $\mathcal{D}(qN)$ if and only if $n_{g,1}(N) = 0$ and $B_g^{(q)}(N) = 0$. Let $M_2^{(q)}(N)$ be the largest sum $a + b$ over pairs of adjacent gaps of $\mathcal{G}(N)$ with $2q \nmid a$, $2q \nmid b$. A missing difference g is*

- *by cause:* a delay if no pair of adjacent gaps of $\mathcal{G}(N)$ sums to g , and a blocked birth if such pairs exist but each contains a gap divisible by $2q$;
- *by position:* a hole if $g < M_2^{(q)}(N)$, and a shadow if $M_2^{(q)}(N) < g < H(qN)$; in the latter case the maximum $H(qN)$ arises from a coincident birth.

In the calm regime every missing difference is a hole and a delay.

Corollary 4.7 (Maximal gaps). *The largest gap $H(qN)$ is the maximum of $H(N)$ and of all spans g with $B_g^{(q)}(N) > 0$. If it is attained by a qualifying constellation of length j , then in each maximal interval of $\mathcal{G}(qN)$ produced by it the prime q alone removes exactly $j - 1$ points. For $3 \mid N$ and $H(qN) < 6q$ this is one point for a pair birth and two points, at distance $2q$ or $4q$, for a coincident birth.*

5 The process on shapes and Holt's chain

Let $m_P(N)$ be the number of driving terms for s with $P(x) = P$, the population of the shape P .

Theorem 5.1. *For every span the shape populations evolve by the exact linear system*

$$m_{P'}(qN) = \sum_P m_P(N) T_q(P, P'),$$

where $T_q(P, P) = q - |\mathcal{C}(P)|$, $T_q(P, P \setminus K) = 1$ for every class $K \in \mathcal{C}(P)$ with $K \cap C = \emptyset$, and all other coefficients are zero. Holt's chain on lengths is the aggregation of shapes of equal length. If no populated shape has two points congruent modulo q — in particular if $|s| < 2q$ — the coefficients depend only on the length, $T(J \rightarrow J) = q - J - 1$ and $T(J \rightarrow J - 1) = J - m$, and Holt's chain is exact.

The condition extends Holt's condition $|s| < 2p_1$ [3, §5], [4, Lemma 2.4], under which all fusions fall into separate images.

Proof. The system is Theorem 3.4. With singleton classes $|\mathcal{C}(P)| = J + 1$ and there are $J - m$ classes disjoint from C , each of one point. $\square$

Remark 5.2. The condition is sufficient, not claimed necessary: contributions of different shapes have different signs and could in principle cancel. In all tested cases in which some populated shape contained two congruent points, Holt's chain was inexact (44 of 44 in the main test, over 200 cases in the audit).

6 Applications

6.1 Holt's exact models for gaps $84 \leq g \leq 104$

Holt's exact models require initial conditions $n_{g,j}(p_0\#)$ at a stage after which every step is linear for the gap considered, that is $g < 2p_1$ for the prime p_1 following p_0 . With initial conditions from $\mathcal{G}(37\#)$ this gives the models for $g \leq 82$; the gaps $84 \leq g \leq 90$ are named in [4, §5] as a current goal.

We computed the complete matrices $n_{g,j}(p\#)$ for $p = 41$ ($g \leq 110$), $p = 43$ ($g \leq 104$) and $p = 47$ ($g \leq 104$). Since $2 \cdot 53 = 106 > 104$, every step from $47\#$ on is linear for $g \leq 104$, so $n_{g,j}(47\#)$ are exact initial conditions, with $p_0 = 47$, for all gaps up to 104. They pass three independent checks.

1. The transitions $37\# \rightarrow 41\# \rightarrow 43\# \rightarrow 47\#$, all of which contain spans beyond $2q$, are reproduced exactly by Theorem 3.4 from Holt's data for $\mathcal{G}(37\#)$ and our pair structures (309, 225 and 153 cells).
2. The populations of gaps in $\mathcal{G}(53\#)$ obtained from $n_{g,j}(47\#)$ by one linear step agree with an independently computed profile of $\mathcal{G}(53\#)$ for all even $g \leq 104$ (52 of 52).
3. The leading coefficient of Holt's model, $\ell_1 = \sum_j w_{g,j}(47\#)$, where $w_{g,j}(47\#)$ is $n_{g,j}(47\#)$ divided by the number $\prod_{3 \leq p \leq 47} (p-2)$ of gaps 2 in $\mathcal{G}(47\#)$, equals the Hardy–Littlewood constant $\prod_{q|g, q>2} \frac{q-1}{q-2}$ [4, eq. (1.3)], [10] exactly, as a rational number, for all eleven gaps. By Holt's Theorem 1.1 [4] (Corollary 1.2 of [3]) this must hold whenever the aggregate counts $\sum_j n_{g,j}(47\#)$ are right, so this check concerns the totals; the distribution over lengths is checked by the first two items.

Table 1 lists the first coefficients $\ell_j = \sum_{i \geq j} \binom{i-1}{j-1} w_{g,i}(47\#)$ of Holt's expansion [4, eq. (1.2)]; the complete matrices are in the accompanying data. Evaluating the models shows how slowly large gaps approach their limits: at $p \approx 10^5$ the relative populations of the gaps 84–104 have reached only between 0.4% and 1.2% of ℓ_1 , in line with the parameter $\lambda = \Theta(1/\ln p)$ of [4].

Holt's worked example. For the constellation $s = 2, 10, 2, 10, 2$ ($|s| = 26$) Holt lists the populations of s and its driving terms of lengths 5, 6, 7 at $p = 5, 7, 11, 13$ [4, Fig. 6]: $(0, 0, 1)$, $(1, 0, 2)$, $(6, 4, 8)$, $(52, 44, 48)$, and notes that the linear system holds from $p = 13$ on. The three transitions $5 \rightarrow 7$, $7 \rightarrow 11$ and $11 \rightarrow 13$ lie outside his model ($|s| \geq 2q$). Theorem 3.4 derives each of them exactly from the previous stage; they involve 3, 3 and 18 removals of several points in one image, respectively.

g	ℓ_1 (= HL constant)	ℓ_2	ℓ_3
84	2.400000	25.8583	129.4360
86	1.024390	11.4282	59.3230
88	1.111111	12.5253	65.7569
90	2.666667	31.1566	169.9885
92	1.047619	12.6201	71.1408
94	1.022222	12.4851	71.4527
96	2.000000	24.9092	145.4980
98	1.200000	15.3165	91.7893
100	1.333333	17.2173	104.4522
102	2.133333	28.4414	178.3882
104	1.090909	14.8365	95.0951

Table 1: Coefficients of Holt’s exact models for the gaps $84 \leq g \leq 104$, initial conditions at $p_0 = 47$.

A remark on quadratic density. Holt shows that the expected quadratic density of a gap increases along the sieve [4, Thm. 4.3], under the assumption $g < 2p_1$. The proof uses the recursion only through the inequality $n_{g,1}(p_k\#) \geq (p_k - 2)n_{g,1}(p_{k-1}\#)$. By Theorem 4.1 this inequality holds for every gap, since $n_{g,1}(qN) = (q - 2 + e_g)n_{g,1}(N) + B_g^{(q)}(N)$ with $e_g \geq 0$ and $B_g^{(q)}(N) \geq 0$. Hence the conclusion of [4, Thm. 4.3] holds for every gap once it occurs, without the restriction $g < 2p_1$.

6.2 Ziller’s table of missing differences

Ziller [5] tabulated, for $k \leq 44$, the even numbers below $h(k)$ that do not occur as gaps in $\mathcal{G}(p_k\#)$ — the *missing differences* of $\mathcal{D}(k)$. Writing $N_{\min}(k)$ for the largest even number such that all even numbers up to it belong to $\mathcal{D}(k)$, he conjectured that $h(k-1) \leq N_{\min}(k)$ for $k > 1$ [5, Conj. 4.1]: every even number up to $h(k-1)$ occurs as a gap of $\mathcal{G}(p_k\#)$, or equivalently, every missing difference of $\mathcal{D}(k)$ exceeds $h(k-1)$. We reproduced his table independently for $18 \leq k \leq 30$ by an exhaustive search (Section 7); all entries agree.

A missing difference of $\mathcal{D}(k)$ arises from $\mathcal{G}(p_{k-1}\#)$ with $q = p_k$, and Corollary 4.6 classifies it. For $k \leq 19$ the calm regime applies ($h(k-1) < 2p_k$), so all missing differences are delays and holes by Corollary 4.4. For $20 \leq k \leq 30$ we searched $\mathcal{G}(p_{k-1}\#)$ exhaustively for the constellations $(2q, g-2q)$ whose existence would make g a blocked birth; none exists, so all are delays. Ten of these fourteen negative results were confirmed independently with a SAT model solved by CP-SAT; four of them (170, 230, 256, 296) are impossible already modulo 3. The position (hole or shadow) is determined by the reach of pairs $M_2^{(q)}$, obtained from a complete classification of the new differences for $20 \leq k \leq 27$ (all 54 present differences in these bands: 50 pair births and 4 coincident births, namely 190, 216, 234 and 282, all of them maxima), from the pair sum 326 at $k = 30$, and from the type of the maxima (Section 6.3).

Result. All 20 missing differences up to $k = 30$ are delays; none is caused by a blocked birth. Seventeen are holes in the set of qualifying pair sums, and three — 186, 188 at $k = 21$ and 328 at $k = 30$ — lie in the shadow of a maximum produced by a coincident birth. Blocked births do occur as individual events, but in this range they never remove a difference from the table.

Observations in other sieves. Replacing the primorial chain by sieves built from random subsets and orders of the primes up to 37, we examined 3689 such sieves (moduli up to $3 \cdot 10^7$, computed by brute force; the survey uses a fixed random seed and is reproducible). The birth

k	missing g	$M_2^{(q)}$	$h(k)$, birth	basis
6	20	22	22, pair	theorem
8	32	34	34, pair	theorem
14	86, 88	90	90, pair	theorem
15	98	100	100, pair	theorem
19	146	152	152, pair	theorem
20	166, 170, 172	174	174, pair	search + SAT
21	182; 186, 188	184	190, coincident	search + SAT
22	194	200	200, pair and coincident	search + SAT
24	230	232	234, coincident	search + SAT
25	254, 256	258	258, pair	search; SAT for 256
27	278	280	282, coincident	search
28	296, 298	300	300, pair	search; SAT for 296
30	328	326	330, coincident	search

Table 2: All missing differences of $\mathcal{D}(k)$ for $k \leq 30$. All are delays. Shadows in bold; all others are holes. “Theorem”: calm regime; “search”: exhaustive search for the blocking constellation; “SAT”: independent confirmation by CP-SAT (for 254 the solver did not finish within one hour).

law was exact in all of them; among their 734 missing differences, 356 were shadows, none was caused by a blocked birth, and every one of them was born at the next stage. These are observations on small moduli, not evidence about large k .

6.3 How maximal gaps arise

By Corollary 4.7, a maximal gap of $\mathcal{G}(p_n\#)$ arises from $\mathcal{G}(p_{n-1}\#)$ either by a pair birth, in which p_n alone removes one point of the maximal interval, or by a coincident birth, in which it removes two points at distance $2p_n$ (up to $n = 64$ one has $h(n) < 4p_n$, so no other route is possible). Counting the points removed by p_n alone in maximal intervals classifies the route. We used the complete lists of maximal configurations for $n \leq 54$ published by Ziller and Morack [9] (ancillary file `remainders.txt`) and the start positions of one maximal interval for each $n \leq 64$ from the OEIS table of A048670 (R. Gerbicz; the values for $n = 62$ – 64 computed by A. Bożek) [12]; see also [6, 7].

- For $n \leq 19$ all maxima are pair births, as the calm regime requires; $n = 20$ is a pair birth as well.
- For $21 \leq n \leq 54$ (all configurations): coincident births only for 22 values of n , pair births only for 6 ($n = 25, 28, 29, 31, 35, 42$), both routes for 6 ($n = 22, 26, 33, 34, 38, 46$).
- For $55 \leq n \leq 64$ (one interval each): coincident births at $n = 56, 58, 59, 61, 62, 64$ and pair births at $n = 55, 57, 60, 63$.
- In every coincident birth the two points are exactly $2p_n$ apart; the two sources agree wherever both are available.

Thus from $n = 21$ on both routes occur, the coincident one more often. The data do not support a simple rule deciding the route: an extrapolation from $n \leq 54$, where $47 \leq n \leq 54$ are all coincident births, predicted coincident births for $n = 55$ – 57 and failed at $n = 55$ and 57 ; nor does the ratio $2p_n/h(n-1)$ decide it, since pair births occur even at its smallest values in the range.

7 Computations and verification

Programs. The population matrices were computed by an exact counting engine that enumerates the windows of a wheel of small primes and treats the remaining primes by an exact covering computation; its results agree with Holt’s published data for $\mathcal{G}(37\#)$ (415 cells) and with his recursion across $37\# \rightarrow 41\# \rightarrow 43\#$ (408 cells). Ziller’s table was verified for $18 \leq k \leq 30$ by an exhaustive search over the sectors of a wheel, with every positive answer certified by an explicit witness checked with gcd. The same machinery, restricted to prescribed interior points, decides the existence of constellations for the classification of Section 6.2. The independent check models the existence of a constellation as a covering problem — one residue class for each prime, all interior points except the prescribed ones covered, the prescribed points and the endpoints not covered — solved by Google OR-Tools CP-SAT; the model was validated against brute force on 70 constellations in $\mathcal{G}(19\#)$.

Brute-force verification. Independently of the engine, all statements of Sections 3–5 were checked by listing complete cycles for moduli up to 510510, including sieves that omit a small prime, in which multi-point removals are frequent. Table 3 summarises the checks.

Statement	Test	Result
Thm. 3.4, gaps	full cycles; $7\# \rightarrow 11$, $11\# \rightarrow 13$, $13\# \rightarrow 17$ and sieves without 5, 7, 11	1585 cells, 0 errors; classes up to 5 points
Thm. 3.4, gaps, engine	$23\# \rightarrow 29\#$, $37\# \rightarrow 41\#$, $41\# \rightarrow 43\#$, $43\# \rightarrow 47\#$	805 cells, 0 errors
Thm. 3.4, constellations Cor. 3.6	160 constellations, spans $2q$ – $5q$ same tests	425 cells, 0 errors first 2-, 3-, 4-, 5-point classes at $2q, 6q, 8q, 12q$
Thm. 3.7	$k = 0, \dots, 3$, 180 constellations; engine for gaps	720 of 720; 609 cells
Thm. 4.1	all births of constellations with at most 4 gaps	783 of 783
Lemma 4.2, Cor. 4.3	chain $2 \rightarrow \dots \rightarrow 30030$, 389 constellations	no exception; 1945 of 1945
Thm. 5.1	$180 + 600$ constellations	sufficient condition never violated
Convention $ s < N$	spans $\geq N$, including $N = 2$	laws hold (214 + 60 cases)

Table 3: Computational verification.

Data availability. Code, data (the matrices $n_{g,j}$ for $41\#$, $43\#$, $47\#$, the pair structures, the classification results and the verification logs) and the programs used for all computations are available at <https://github.com/hojka1987t-dot/sieve-dynamics-beyond-2p>; the version used in this paper (release v1.0.0) is archived at Zenodo as doi:10.5281/zenodo.23009305. Table 1 and the census of Section 6.3 are reproduced by the scripts in `code/analysis`, Table 3 and the survey of Section 6.2 by those in `code/brute_force`. The classification in Table 2 rests on the result files in `data/`, produced by the exhaustive searches and the SAT check, whose programs are included; these runs take from hours to about a day.

8 Open questions

1. *Ziller’s Conjecture 4.1.* By Theorem 4.1 it states that every even number up to $h(k-1)$ is a gap of $\mathcal{G}(p_{k-1}\#)$ or the span of a qualifying constellation; in the calm regime, a gap

or a sum of two adjacent gaps. Is there a structural reason why the holes of this set stay near its top? In all small sieves we examined no missing difference persisted for more than one stage.

2. *Which route produces the maximum?* Both pair and coincident births of maxima occur for $n \geq 21$, with no simple rule deciding between them.
3. *Necessity in Theorem 5.1.* Is Holt's chain exact only if no populated shape has two points congruent modulo q ?
4. *Higher thresholds.* Up to $n = 64$ the maxima stay below $4p_n$. For larger n they will pass the thresholds $4p_n$ and $6p_n$, where births through pairs at distance $4q$ and through classes of three points become possible for maximal gaps.
5. *Paired progressions.* The same method applies to sieves removing two residue classes per prime, as in the paired Jacobsthal function of Ziller and Morack [11]; the classes of coincidence and the thresholds then change.

Acknowledgements

Parts of the code, the computations and the drafting were carried out with the assistance of Claude, an AI model developed by Anthropic. The author takes full responsibility for the content.

Note on versions. A preliminary version of this text is included in release v1.0.0 of the accompanying repository (doi:10.5281/zenodo.23009305). The present text differs from it in attributing the residue-class criterion explicitly to Holt [3, Lemma 3.1], and the identity for ℓ_1 to [4, Thm. 1.1] and [3, Cor. 1.2]; no result has changed.

References

- [1] F. B. Holt and H. Rudd, *Combinatorics of the gaps between primes*, Connections in Discrete Mathematics, Simon Fraser Univ. (2015); arXiv:1510.00743.
- [2] F. B. Holt, *Models for gaps $g = 2p_1$* , preprint (2023), arXiv:2309.16833.
- [3] F. B. Holt, *Eratosthenes sieve supports the k -tuple conjecture*, preprint (2025), arXiv:2502.20470.
- [4] F. B. Holt, *Discrete dynamics of Eratosthenes sieve*, preprint (2026), arXiv:2608.26384; cited by the numbering of versions 3 and 4, which agree.
- [5] M. Ziller, *On differences between consecutive numbers coprime to primorials*, preprint (2020), arXiv:2007.01808.
- [6] T. Hagedorn, *Computation of Jacobsthal's function $h(n)$ for $n < 50$* , Math. Comp. **78** (2009), no. 266, 1073–1087.
- [7] L. Hajdu and N. Saradha, *Disproof of a conjecture of Jacobsthal*, Math. Comp. **81** (2012), no. 280, 2461–2471.
- [8] H.-J. Kanold, *Über eine zahlentheoretische Funktion von Jacobsthal*, Math. Ann. **170** (1967), 314–326.

- [9] M. Ziller and J. F. Morack, *Algorithmic concepts for the computation of Jacobsthal's function*, preprint (2016), arXiv:1611.03310; ancillary file `remainders.txt`.
- [10] G. H. Hardy and J. E. Littlewood, *Some problems of 'Partitio numerorum' III: on the expression of a number as a sum of primes*, Acta Math. **44** (1923), 1–70.
- [11] M. Ziller and J. F. Morack, *Divisibility in paired progressions, Goldbach's conjecture, and the infinitude of prime pairs*, preprint (2017), arXiv:1706.00317.
- [12] OEIS Foundation, *The On-Line Encyclopedia of Integer Sequences*, sequence A048670.