

# 1 Birthweight of babies

TOPICS: Linear normal model

The `births` data in the `Epi` package contains recordings of birthweights of babies in a London hospital together with recordings of several possible explanatory variables.

The purpose of this exercise is to establish a model for understanding how the birthweight is related to the explanatory variables.

```
library(Epi)
data(births)
head(births)
```

	id	bweight	lowbw	gestwks	preterm	matage	hyp	sex
1	1	2974	0	38.52	0	34	0	2
2	2	3270	0	NA	NA	30	0	1
3	3	2620	0	38.15	0	35	0	2
4	4	3751	0	39.80	0	31	0	1
5	5	3200	0	38.89	0	33	1	1
6	6	3673	0	40.97	0	33	0	2

```
dim(births)
```

```
[1] 500 8
```


```
sapply(births, class)
```

	id	bweight	lowbw	gestwks	preterm	matage	hyp	sex
	"numeric"	"numeric"	"numeric"	"numeric"	"numeric"	"numeric"	"numeric"	"numeric"

1. (a) Plot the birth weights of babies and observe the distribution. Comments?

Solution:

```
hist(births$bweight)
```



- (b) See how birth weight of babies are related to gestation period, and maternal age by scatter plot. Differentiate between males and females in the plot. (Hint: `xypplot()` could be your friend.) Comments?
  - (c) Make a scatterplot of all pair of variables. (Hint: `pairs()` could be your friend). Comments?
2.
  - (a) Assume that number of boys follows a  $\text{bin}(N = 500, \theta)$ -distribution.
  - (b) Estimate  $\theta$  from observed data. Estimate the variance of  $\hat{\theta}$  from data.
  - (c) What is the probability of observing a male child?
  - (d) What is the probability of observing at least 26 babies of the same sex in 44 births? (Use above estimated value of  $\theta$ .)
3. Fit a linear normal model  $M_1$  for  $bweight \sim gestwks + matage$  to the `births` data.
  - (a) Write down in precise mathematical terms what your model is.
  - (b) Write down in your own words what the assumptions behind a linear normal model are. Investigate the residuals of  $M_1$ . Comment on whether you find that these assumptions are likely to be satisfied for these data.
  - (c) Write down in your own words what a least squares fit of a model to data is.
  - (d) From model  $M_1$  report the estimated parameters and their standard deviations.
  - (e) In model  $M_1$ , calculate the values of fitted values (use function `predict()`) and plot the fitted line over the observations.
  - (f) Test the effect of each term in the model  $M_1$  using the function `drop1()`.

- (g) Predict the birth weights at weeks 10, 15, 20, 25, 30, 35, 40, 45, 50 for women of age 20, 30 and 40. Include confidence intervals in your prediction. Present the results as a table and as illustrative graphics. Comment on these predictions.
4. Fit an interaction model  $M_2$  for  $bweight \sim gestwks + matage + gestwks * matage$  to the `births` data.
  5. Write down  $M_2$  in precise mathematical terms.
  6. Investigate whether  $M_2$  fits well to data.
  7. Test whether the additive model is more appropriate or the interactive model.
  8. Fit a model  $M_3$  that differentiate the two slope for sexes.
  9. Write down  $M_3$  in precise mathematical terms.
  10. Investigate whether  $M_3$  fits well to data.
  11. Investigate whether it is necessary to have two slopes for each sex by comparing the models  $M_1$  and  $M_3$ .