

Topology

Summary with my Intuitions

Hojune Lee

School of Computing, KAIST

Contents

1	Definition of Topology	3
2	Some Basic Famous Topologies	6
3	Closed Sets and Limit Points	8
4	Basic notion of Hausdorff Space	10
5	Continuous Functions and Homeomorphisms	11
6	The Product Topology	14
7	The Quotient Topology	19
8	Connected Spaces	24
9	Connected Subspaces of The Real Line	27
10	Compact Space	29

1 Definition of Topology

In many times, we ‘draw’ some mathematical objects especially sets. One of the key intuition of studying topology is how can we ‘draw’ such things. To draw somethings formally, we need to define bunch of things like *shape of sets*. As we describe *shape of groups* with its inner structures like subgroups, we describe *shape of sets* with its inner structures like subsets.

Definition 1.1. Topology

A topology $\mathcal{T}$ on a set X is collection of subsets of X having following axioms.

1. $\emptyset, X \in \mathcal{T}$
2. $\mathcal{T}$ is closed under arbitrary unions.
3. $\mathcal{T}$ is closed under finite intersections.

Then we denote the pair $(X, \mathcal{T})$ is *topological space* (or just *space*) and call every $U \in \mathcal{T}$ by is *open set* in X . Then now X is shaped by $\mathcal{T}$ with certain axioms such that related subsets of X . This is very similar to defining groups.

Definition 1.2. Group

Let G be a set and $\star : G \times G \rightarrow G$ is binary operation. $(G, \star)$ is group when have following axioms.

1. $\forall a, b, c \in G, \quad a \star (b \star c) = (a \star b) \star c.$
2. $\exists 1_G \in G$ such that $\forall x \in G, 1_G \star x = x \star 1_G = x.$
3. $\forall x \in G, \exists x^{-1} \in G$ such that $x \star x^{-1} = x^{-1} \star x = 1_G.$

Well, the definition of topology is quite less intuitive. However we already familiar with ‘shapes’ in $\mathbb{R}^n$ and often draw them in natural. Topology is just generalize such notions into general sets. In analysis in $\mathbb{R}^n$, the key block was *open balls*. We generalize this notion into general set also, which is called *basis*.

Definition 1.3. Basis

A *basis* $\mathcal{B}$ is collection of subsets of X having following axioms.

1. $\forall x \in X, \exists B \in \mathcal{B}$ such that $x \in B.$
2. $\forall B_1, B_2 \in \mathcal{B},$ if $x \in B_1 \cap B_2, \exists B^* \in \mathcal{B}$ such that $x \in B^* \subseteq B_1 \cap B_2.$

This is very natural property when we imagine open balls in $\mathbb{R}^n$. Another good

motivation to understand this is that, understand each $B \in \mathcal{B}$ is collection of ‘near’ elements. In metric space we’ll define later, such notion of distance is very natural but here, we need to understand abstractly. This collection is called *basis* since it generates some topology on X .

Definition 1.4. Topology generated by Basis

Let $\mathcal{B}$ is a basis of X . Then, define

$$\mathcal{T}_{\mathcal{B}} = \{U \subseteq X \mid \forall x \in U, \exists B \in \mathcal{B}, x \in B \subseteq U\}$$

Then $\mathcal{T}_{\mathcal{B}}$ is a topology on X and called topology generated by $\mathcal{B}$.

Remark 1.5

$\mathcal{T}_{\mathcal{B}}$ is well-defined topology.

As we noted $\mathcal{B}$ ‘generates’ some topology and it is called basis, there are more intuitive way to construct $\mathcal{T}_{\mathcal{B}}$.

Lemma 1.6. Topology generated by Basis

$\mathcal{T}_{\mathcal{B}}$ = collection of all unions in $\mathcal{B}$.

Then next the natural question is that, *can we compare 2 topologies with its basis?* The answer is yes, and we’ll use following theorem in many times. Following theorem allows us to compare the inclusion relation (or equivalence relation) of 2 topologies via its basis.

Theorem 1.7. Topology Comparison with Basis

Let $\mathcal{B}_1, \mathcal{B}_2$ be basis of X and $\mathcal{T}_1, \mathcal{T}_2$ be topology of X generated by $\mathcal{B}_1, \mathcal{B}_2$ respectively. Then $\mathcal{T}_1 \subseteq \mathcal{T}_2$ if and only if $\forall x \in X, \forall B_1 \in \mathcal{B}_1$ such that $x \in B_1$, there exists $B_2 \in \mathcal{B}_2$ such that $x \in B_2 \subseteq B_1$.

We observe that once we have basis $\mathcal{B}$, then we can generate some topology by unioning them. However, in 1.1 topology is closed under not only union but also finite intersection. So we can define more light basis for topology(or, basis for basis).

Definition 1.8. Subbasis

A subbasis $\mathcal{S}$ is collection of subsets of X having following axiom.

1. $\forall x \in X, \exists S \in \mathcal{S}$ such that $x \in S$.

equivalently, the entire union of $\mathcal{S}$ is X .

Then, we can observe that

$$\mathcal{S} \xrightarrow{\text{finite intersections}} \mathcal{B} \xrightarrow{\text{arbitrary unions}} \mathcal{T}$$

2 Some Basic Famous Topologies

In this section, we'll define a few useful topologies.

Definition 2.1. Standard Topology

Let $X = \mathbb{R}$ and $\mathcal{B} = \{(a, b) \mid a < b\}$. $\mathcal{T}_{\mathcal{B}}$ is called standard topology of $\mathbb{R}$.

Definition 2.2. Lower Limit Topology

Let $X = \mathbb{R}$ and $\mathcal{B} = \{[a, b) \mid a < b\}$. $\mathcal{T}_{\mathcal{B}}$ is called lower limit topology of $\mathbb{R}$.

Definition 2.3. Order Topology

Let X be a simply ordered set. i.e., $\forall x, y \in X$, $x < y$ or $x = y$ or $y < x$ holds and the relation $<$ is transitive. Then we define standard topology for X which is called order topology. Let

$$\mathcal{B} = \begin{cases} (a, b) & \forall a < b \in X \\ [m, b) & m \text{ is smallest element in } X \\ (a, M] & M \text{ is largest element in } X \end{cases}$$

Topology generated by this $\mathcal{B}$ on simply ordered set X is called *order topology*.

For order topology, we can define subbasis generates such basis. It is called open rays and define by

$$\mathcal{S} = \{(-\infty, a), (a, \infty) \mid a \in X\}$$

Remark 2.4

$\mathcal{S}$ is subbasis of $\mathcal{B}$.

One obvious step is defining 'sub' space and find its natural topology. The most natural definition is that, if Y is subspace of X , it inherits all open sets in X into Y . means that,

If Y is subspace of X , then open sets in X is also open in Y .

Definition 2.5. Subspace Topology

Let $(X, \mathcal{T}_X)$ is a topological space and $Y \subseteq X$. We define

$$\mathcal{T}_Y = \{U \cap Y \mid U \in \mathcal{T}_X\}$$

for topology of Y and call *subspace topology*. And $(Y, \mathcal{T}_Y)$ is called *subspace* of X .

Then similarly the basis of $\mathcal{T}_Y$ has same relation.

$$\mathcal{B}_Y = \{B \cap Y \mid B \in \mathcal{B}_X\} \text{ is basis for } \mathcal{T}_Y.$$

Warning 2.6

Open sets in general $Y \subseteq X$ do not need to also open in X .

Lemma 2.7. When open sets in Y is also open in X ?

If Y is subspace of X and Y is open in X , then open sets in Y is also open in X .

Proof. From definition, any open set $V \in \mathcal{T}_Y$ in Y is formed $V = U \cap Y$ and $U, Y \in \mathcal{T}_X$ so $V \in \mathcal{T}_X$. $\square$

Then also one natural question is arised. **How about subspace of simply ordered space?** Here interesting results are occur. We observe 3 subspace of $\mathbb{R}$.

$$Y_1 = [0, 1], \quad Y_2 = [0, 1) \cup \{2\}, \quad Y_3 = [0, 1] \cup [2, 3]$$

In Y_1, Y_3 , subspace topology of them are equivalent to ordered topology of themselves. However, in Y_2 , subspace topology is not equivalent to ordered topology since then singleton set $\{2\}$ is open in subspace topology but not in ordered topology. In general,

Theorem 2.8. Subspace Topology and Order Topology

Let $(X, \mathcal{T}_X)$ is ordered topology space and $Y \subseteq X$. $\mathcal{T}_1$ is subspace topology of Y and $\mathcal{T}_2$ is ordered topology of Y . (since Y also simply ordered) Then, $\mathcal{T}_2 \subseteq \mathcal{T}_1$ in general. Furthermore, $\mathcal{T}_1 = \mathcal{T}_2$ if and only if every gap in Y is either open or closed.

3 Closed Sets and Limit Points

We defined basic notions of topology via well-shaped subsets called *open sets*. Another type of sets can be useful.

Definition 3.1. Closed Sets

$C \subset X$ is closed when $X \setminus C$ is open in X

Openness and closedness is familiar notion in $\mathbb{R}^n$ space. Closed set is generalization of the closedness into general sets. Sometimes, arguing with closed set is much easier than open sets despite they are just complement of open sets.

Theorem 3.2. Topology Axioms for Closed Sets

Let $(X, \mathcal{T})$ be a topological space. Then following variation of axioms hold.

1. $\emptyset, X$ is closed set.
2. Arbitrary intersections of closed sets are closed.
3. Finite unions of closed sets are closed.

Theorem 3.3. Subspace Topology for Closed Sets

Let Y be subspace of X . Then, $D \subseteq Y$ is closed in Y if and only if there exists closed set C in X such that $D = C \cap Y$.

Proof. If $D \subseteq Y$ is closed, then $Y \setminus D$ is open in Y . So, we can find open set U in X such that $Y \setminus D = U \cap Y$. Then, $D = U^c \cap Y$ and U^c is closed in X . If $D = C \cap Y$ for closed set C in X , $Y \setminus D = Y \cap C^c$ is open in Y , so D is closed in Y . $\square$

Theorem 3.4. When closed sets in Y is also closed in X ?

If Y is subspace of X and Y is closed in X , then closed sets in Y is closed in X .

Now we define few more way to construct meaningful sets.

Definition 3.5. Interior and Closure of a Set

Let A be any subset of X . The interior of A (denoted by $\text{Int } A$) is union of all open sets contained in A . i.e., the largest open set in A . The closure of A (denoted by $\bar{A}$) is intersection of all closed sets containing A . i.e., the smallest closed set containing A .

In many times, closure of set acts quite important roles. So we state few theorems related to the closure of set.

Theorem 3.6. Closure in the Subspace

Let X be a topological space and Y is subspace of X . For $A \subseteq Y \subseteq X$, denote $\bar{A}$ by closure of A in X . Then closure of A in Y is determined by $\bar{A} \cap Y$.

Proof. Let C be closed set in X such that $A \subseteq C$. Then, $C \cap Y$ is closed in Y and $A \subseteq C \cap Y$. So, $\bar{A} \cap Y \subseteq \text{closure of } A \text{ in } Y$. And for D be closed set in Y such that $A \subseteq D$, by definition, $D = C \cap Y$ where C is closed and $A \subseteq C$. So, closure of A in Y is equivalent to $\bar{A} \cap Y$. $\square$

Theorem 3.7. Points in Closure

Let X be a topological space and $A \subseteq X$ be any subset. Then $x \in \bar{A}$ if and only if every *neighborhoods* of x *intersects* with A .

Here, *neighborhood* of x means that any open sets containing x . Also, we say A, B are intersects when $A \cap B \neq \emptyset$. This is important property of points in the closure.

Proof. If $x \in \bar{A}$ but $\exists U$ which is neighborhood of x and $U \cap A = \emptyset$, Then $A \subseteq U^c$ and U^c is closed. So by definition, $\bar{A} \subseteq U^c$ so $x \in U^c$ which is contradiction. If every neighborhoods of x intersects with A but $x \notin \bar{A}$, then it means that there exists a closed set C such that containing A but does not contain x . It implies that $x \in X \setminus C$ and $X \setminus C \cap A = \emptyset$ which is contradiction. $\square$

Definition 3.8. Limit Points

If A is a subset of topological space X , we say that x is a limit point of A when every neighborhoods of x intersects with A other than x itself. i.e., $x \in \overline{A \setminus \{x\}}$.

Then now, we can describe the closure of set A with A itself and its limit points.

Theorem 3.9. Closure Representation

Let X be a topological space and $A \subseteq X$ be any subset. Denote A' by set of limit points of A . Then,

$$\bar{A} = A \cup A'$$

In other words, $\bar{A} \setminus A = \text{limit points of } A \text{ but not in } A$.

Corollary 3.10. When A is closed?

$A \subseteq X$ is closed if and only if $A' \subseteq A$. i.e., A contains all of its limit points.

4 Basic notion of Hausdorff Space

We learned how can we define each shapes of sets and construct them into spaces. Then we can wonder, what is the good shapes for spaces? And how can we categorize them? Actually the answer for this question is dealt in further section but we define very part of them here since it will be heavily used.

Definition 4.1. Hausdorff Space

A topological space $(X, \mathcal{T})$ is called Hausdorff space when

$$\forall x \neq y \in X, \text{ there exists disjoint neighborhoods of } x, y \text{ respectively.}$$

In other words, we can *separate* two *points* via their neighborhoods always.

Actually we can understand that why this is good property for space since real line $\mathbb{R}$ have this property. Once our space is Hausdorff, we can find a few interesting facts.

Theorem 4.2. Finite sets in Hausdorff spaces

Let X be a Hausdorff space. Then, every finite set in X is closed.

Proof. Since finite union of closed sets are closed, so it is sufficient to show that any singleton set $\{x\}$ is closed in X . Now suppose that $y \in X$ is a limit point of $\{x\}$. If $x \neq y$, since X is Hausdorff we can find U_1, U_2 such that both are neighborhoods of x, y respectively and $U_1 \cap U_2 = \emptyset$. Then, according to theorem 3.7, y can't be limit point of $\{x\}$. So unless x is limit point or not, we can state $\{x\}$ contained all its limit points and so closed set. $\square$

Now we find next interesting property of Hausdorff space. Let's call the $\mathbb{N}$ -indexed sets of points in X by *sequences* and denoted by $\{x_n\}_{n \in \mathbb{N}}$. In general notion, we state that $\{x_n\}_{n \in \mathbb{N}}$ converges to $x \in X$ when

For any neighborhood U of x , we can find $N \in \mathbb{N}$ such that $x_m \in U$ for $m \geq N$.

Then following good property holds in Hausdorff space.

Theorem 4.3. Unique Convergency in Hausdorff Space

If X is a Hausdorff space, then any sequences in X converges into at most one point.

Theorem 4.4. Hausdorff spaces with Famous Topologies

Any space with Order topology, Subspace topology of Hausdorff space is Hausdorff.

5 Continuous Functions and Homeomorphisms

From highschool to undergrad analysis courses, the continuity of function is matter. By the way, the concept of continuity in $\mathbb{R}^n$ (in general, metric space) is quite obvious. Our goal is generalize them into general topological space. One beautiful definition is following.

Definition 5.1. Continuous Function

Let X, Y be any topological spaces. A function $f : X \rightarrow Y$ is *continuous* when for any open set V in Y , the preimage $f^{-1}(V) \subseteq X$ is also open set in X .

Remark 5.2

Continuity is depends on not only f itself but also topologies of X and Y .

Here, I mention why we take preimage for define continuity. First at all, this notion is called **pull-back**. This is important since it preserves many structural things in codomain space Y .

Proposition 5.3. Preimage Propositions

1. $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$
2. $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$
3. $f^{-1}(A^c) = (f^{-1}(A))^c$

These are not true in original mapping f . So, we can analyze topological structures of Y in X via pull-back. Also, the natural notion ε - δ definition of continuity also refer the preimage. Once we view this definition as just generalization of ε - δ definition, it seemed be natural.

Lemma 5.4. Checking Continuity

Let $\mathcal{S}$ be subbasis of topological space Y . Then for any $S \in \mathcal{S}$, if $f^{-1}(S)$ is open set in X , f is continuous.

Proof. Every open sets in Y can be generated by arbitrary union and finite intersection of elements in $\mathcal{S}$. Since 5.3 holds for preimages, if every $f^{-1}(S)$ is open then every preimage of open set is also open in X . $\square$

Theorem 5.5. Equivalence statements for Continuity

The followings are equivalent.

1. f is continuous.
2. For any $A \subset X$, $f(\bar{A}) \subseteq \overline{f(A)}$
3. For every closed set C of Y , $f^{-1}(C)$ is closed in X .
4. For each $x \in X$, each neighborhood V of $f(x)$, there exists neighborhood U of x such that $f(U) \subseteq V$.

(4) is just changing of forms of statement into ε - δ definition form. The remarkable one is second one. It can be written: If x is a limit point of A in X , then $f(x)$ is a limit point of $f(A)$ in Y or equivalently, continuous map preserves the limit point relation.

Definition 5.6. Homeomorphism

Let X, Y be topological spaces and $f : X \rightarrow Y$ is a bijection. If both f, f^{-1} are continuous then f is called *homeomorphism* and X, Y are *homeomorphic*.

The homeomorphism f between X and Y construct bijection between $\mathcal{T}_X$ and $\mathcal{T}_Y$ also. i.e., open sets in both spaces are one-to-one corresponds. We say that some property is topological property if it preserves by homeomorphisms. And also we say that an injective continuous map $f : X \rightarrow Y$ is *embedding*.

Theorem 5.7. Constructing Continuous Functions

Let X, Y and Z be topological spaces.

1. Any constant function is continuous.
2. If A is a subspace of X , the natural inclusion $j : A \hookrightarrow X$ is continuous.
3. If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are continuous, then $h = g \circ f$ is continuous.
4. If $f : X \rightarrow Y$ is continuous then for subspace A of X , $f|_A : A \rightarrow Y$ is continuous.
5. Let $f : X \rightarrow Y$ is continuous. If $\text{Im}(X)$ is subspace of Y , then $g : X \rightarrow \text{Im}(X)$ is continuous. If Y is subspace of Z , then $h : X \rightarrow Z$ is continuous.
6. If $X = \bigcup U_\alpha$ and every $f|_{U_\alpha}$ is continuous, then $f : X \rightarrow Y$ is continuous.

These are not hard to follow and prove. We state important lemma here.

Lemma 5.8. Pasting Lemma

Let X, Y be topological spaces and $X = A \cup B$ where A, B are closed subspaces in X . Let $f : A \rightarrow Y$ and $g : B \rightarrow Y$ be continuous. If $f(x) = g(x)$ for every $x \in A \cap B$, then we can define continuous function $h : X \rightarrow Y$ defined by

$$h(x) = \begin{cases} f(x) & x \in A \\ g(x) & x \in B \end{cases}$$

This is well-defined since we assume $f(x) = g(x)$ for $x \in A \cap B$.

Proof. As we stated, $h : X \rightarrow Y$ is well-defined. Let C be any closed set in Y . Then, $f^{-1}(C), g^{-1}(C)$ are closed in A, B respectively by 5.5. However since A, B are closed subspace of X , $f^{-1}(C), g^{-1}(C)$ is closed in X by 3.4. However,

$$h^{-1}(C) = f^{-1}(C) \cup g^{-1}(C) \implies \text{closed in } X$$

So, $h : X \rightarrow Y$ is continuous function. □

6 The Product Topology

We often consider Cartesian product of sets(or spaces). It is good time to state how Cartesian product can be generalized into topological concepts. We'll see familiar product $X \times Y$ first and then generalize into arbitrary Cartesian product $\prod_{\alpha \in J} X_\alpha$ for index set J .

Definition 6.1. Simple Cartesian Product Space $X \times Y$

Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ be topological spaces. For the Cartesian product $X \times Y$, we define *product topology* generated by following basis.

$$\mathcal{B} := \{U \times V \mid U \in \mathcal{T}_X, V \in \mathcal{T}_Y\}$$

Remark 6.2

We take $\mathcal{T}_X \times \mathcal{T}_Y$ as just a **basis** for $X \times Y$. So, the *product topology* on $X \times Y$ is arbitrary unions of $\mathcal{T}_X \times \mathcal{T}_Y$. Intuitively, imagine the $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$ space. The 'shapes' in $\mathbb{R}^2$ is more various than just $\mathcal{T}_{\mathbb{R}} \times \mathcal{T}_{\mathbb{R}}$. So, $\mathcal{T}_X \times \mathcal{T}_Y$ just basis generates product topology.

Here we can check a few properties that we stated in previous sections.

Lemma 6.3. Basis of Product Topology

Let X, Y be topological spaces. If $\mathcal{B}_X, \mathcal{B}_Y$ are basis for X, Y respectively, then $\mathcal{B} = \mathcal{B}_X \times \mathcal{B}_Y$ is a basis for product topology of $X \times Y$.

Definition 6.4. Projection

We define $\pi_1 : X \times Y \rightarrow X$ and $\pi_2 : X \times Y \rightarrow Y$ in natural manner.

Lemma 6.5. Subbasis of Product Topology

Let X, Y be topological spaces. Following is subbasis for product topology of $X \times Y$:

$$\mathcal{S} = \{\pi_1^{-1}(U) \mid U \text{ is open in } X\} \cup \{\pi_2^{-1}(V) \mid V \text{ is open in } Y\}$$

Lemma 6.6. Subspace of Product Space

Let X, Y be topological spaces. If A, B are subspaces of X, Y respectively, then $A \times B$ is subspace of $X \times Y$.

Lemma 6.7. Product of Hausdorff is Hausdorff

If X, Y are Hausdorff space, then $X \times Y$ with product topology is Hausdorff.

These are not hard to understand. We can easily generalize this into finite Cartesian product by

$$X_1 \times X_2 \times \cdots \times X_n$$

Then here 6.5 still holds. One natural following step is considering the arbitrary Cartesian products.

Definition 6.8. Indexing, J -tuple and Coordinates

Let J be an arbitrary index set. Given a set X , we define a J -tuple of elements of X to be a function

$$\mathbf{x} : J \rightarrow X$$

We often denote $\mathbf{x}(\alpha)$ by x_α and call it the α -th coordinate of $\mathbf{x}$.

With this arbitrary index set J , we can define general Cartesian product.

Definition 6.9. Cartesian Product

Let J be an arbitrary index set and $\{A_\alpha\}_{\alpha \in J}$ be an indexed family of sets. The Cartesian product of this indexed family is denoted by

$$\prod_{\alpha \in J} A_\alpha = \{\text{All } J\text{-tuples } \{x_\alpha\}_{\alpha \in J} \text{ where each } x_\alpha \in A_\alpha\}$$

Now we can define this notion into topological Cartesian product space. However, there are 2 ways to construct topology on such space. Actually, one is quite natural but less useful, and the other is related to 6.5. It means that 6.5 is not the case in general, and two topologies are distinct when J is an infinite index set.

Definition 6.10. Box Topology

Given arbitrary index set J and $\{X_\alpha\}_{\alpha \in J}$ be a family of topological spaces. Then, we can find Cartesian product $\prod_{\alpha \in J} X_\alpha$. Then we can define a basis in following way:

$$\mathcal{B} = \left\{ \prod_{\alpha \in J} U_\alpha \mid U_\alpha \text{ is open in } X_\alpha \right\}$$

Then this is topology on $\prod_{\alpha \in J} X_\alpha$ and called *box topology*.

Definition 6.11. Projection

For each $\beta \in J$, we define

$$\pi_\beta : \prod_{\alpha \in J} X_\alpha \rightarrow X_\beta$$

defined by

$$\pi_\beta : \mathbf{x} \mapsto \mathbf{x}(\beta)$$

This is called the projection mapping associated with the index β .

We generalize the notion of projection in 6.4 into general Cartesian product.

Definition 6.12. Product Topology

We can define a subbasis $\mathcal{S}$ of $\prod_{\alpha \in J} X_\alpha$ in following way:

$$\mathcal{S}_\beta = \{\pi_\beta^{-1}(U_\beta) \mid U_\beta \text{ is open in } X_\beta\} \implies \mathcal{S} = \bigcup_{\beta \in J} \mathcal{S}_\beta$$

The topology generated by this subbasis is called the *product topology*. And with this, $\prod_{\alpha \in J} X_\alpha$ is called *product space*.

Compare these two topology is not quite difficult. As we stated in 1.7, it is enough to show the differences between the basis of box topology and the basis of product topology.

Proposition 6.13. Comparison between Box topology and Product Topology

The basis of box topology contains all possible forms of $\prod_{\alpha \in J} U_\alpha$ where U_α is any open set in X_α . However, the basis of product topology is obtained by **finite intersections** of subbasis. So, the basis of product topology contains only has the following form of elements:

$$\pi_{\beta_1}^{-1}(U_{\beta_1}) \cap \cdots \cap \pi_{\beta_n}^{-1}(U_{\beta_n}) = \prod_{\alpha \in J} U_\alpha, \quad U_\alpha = \begin{cases} X_\alpha & \alpha \notin \{\beta_1, \dots, \beta_n\} \\ U_{\beta_i} & i \in [n] \end{cases}$$

So the basis of box topology contains the basis of product topology, so *box topology is finer than product topology*.

According to the textbook [1], there is an answer **why we generally use the definition of product topology**. i.e., despite the definition of box topology is more finer and stronger, we assume that the topology of $\prod_{\alpha \in J} X_\alpha$ will be product topology in general. It is because, there are many important theorems in topology such that holds for product topology, but fails for box topology. **However note that when index set J is infinite then matter. For finite Cartesian product of spaces, there are not any confusion.**

Now let's state bunch of theorems here.

Theorem 6.14. Basis generated by Basis of each Space

Suppose that $\{X_\alpha\}_{\alpha \in J}$ be family of spaces and $\mathcal{B}_\alpha$ is the basis of X_α . Then, the basis of *box topology* is given by

$$\mathcal{B}_{\text{box}} = \left\{ \prod_{\alpha \in J} B_\alpha \mid B_\alpha \in \mathcal{B}_\alpha \right\}$$

The basis of *product topology* is given by

$$\mathcal{B}_{\text{prod}} = \left\{ \prod_{\alpha \in J} B_\alpha \mid B_\alpha = \begin{cases} B_\alpha \in \mathcal{B}_\alpha & \text{Finitely many } \alpha \\ X_\alpha & \text{otherwise} \end{cases} \right\}$$

Theorem 6.15. Subspace Topology and Box and Product Topology

Let A_α be a subspace of X_α for each $\alpha \in J$. Now box topology of $\prod_{\alpha \in J} A_\alpha$ is subspace of box topology of $\prod_{\alpha \in J} X_\alpha$. And same for product topology.

Theorem 6.16. Hausdorff in Box and Product Topology

If each X_α is Hausdorff, $\prod_{\alpha \in J} X_\alpha$ is also Hausdorff in both box and product topologies.

Theorem 6.17. Closure in Box and Product Topology

Let $A_\alpha \subseteq X_\alpha$ for each $\alpha \in J$. If $\prod X_\alpha$ is given either the product or the box topology, then

$$\prod \overline{A_\alpha} = \overline{\prod A_\alpha}$$

Proof. Suppose that $\mathbf{x} = (x_\alpha)_{\alpha \in J} \in \prod \overline{A_\alpha}$. Then, let $\prod U_\alpha$ be any basis neighborhood of $\mathbf{x}$ in box or product topology of $\prod X_\alpha$. Then each U_α is neighborhood of x_α in X_α , and since $x_\alpha \in \overline{A_\alpha}$ so we can find $y_\alpha \in A_\alpha \cap U_\alpha$. So, $\mathbf{y} = (y_\alpha)_{\alpha \in J} \in \prod U_\alpha \cap \prod A_\alpha$. So $\mathbf{x} \in \overline{\prod A_\alpha}$.

Suppose that $\mathbf{x} \in \overline{\prod A_\alpha}$. Let V_β be an arbitrary open set of X_β containing x_β . Since $\pi_\beta^{-1}(V_\beta)$ is open in $\prod A_\alpha$ in both topology and containing $\mathbf{x}$, there exists $\mathbf{y} \in \pi_\beta^{-1}(V_\beta) \cap \prod A_\alpha$. This implies that, $y_\beta \in V_\beta \cap A_\beta$. So, $x_\beta \in \overline{A_\beta}$. $\square$

Following is the first situation when we can observe box topology and product topology is different.

Theorem 6.18. Continuity in Product Topology

Let $f : A \rightarrow \prod X_\alpha$ be given by the equation

$$f(a) = (f_\alpha(a))_{\alpha \in J}$$

where $f_\alpha : A \rightarrow X_\alpha$ for each α . Let $\prod X_\alpha$ have the product topology. Then the function f is continuous if and only if each function f_α is continuous.

Proof. ($\Rightarrow$) Suppose that f is continuous. Then for any open set U_β in A_β , we can define following open set in $\prod X_\alpha$,

$$U^* = \prod U_\alpha = \begin{cases} U_\beta & \alpha = \beta \\ X_\alpha & \alpha \neq \beta \end{cases}$$

Then this is a basis element of product topology of $\prod X_\alpha$, hence open. Since f is continuous mapping, $f^{-1}(U^*) = f_\beta^{-1}(U_\beta)$ is open in A . So each f_β is continuous.

($\Leftarrow$) Suppose that each f_α is continuous. According to 5.4 it is sufficient to show that preimage of subbasis is open in A . Each subbasis has the form

$$\pi_\beta^{-1}(V_\beta), \quad V_\beta \text{ is open in } X_\beta$$

So that, the preimage of each subbasis element has the form

$$f^{-1}(\pi_\beta^{-1}(V_\beta)) = f_\beta^{-1}(V_\beta) \implies \text{open in } A \text{ since } f_\beta \text{ is continuous}$$

Thus, f is continuous mapping. □

7 The Quotient Topology

The notion of quotient is always non-trivial (But, very fun if once you understand). In the Group theory, we encounter the quotient group with normal subgroups. The intuitive concept was that, grouping some subset into sort of equivalent classes. We can do same thing in topology. Whenever the space is X , we can grouping some elements in equivalent classes. Such notion is visualizable and natural, see following figure.




Figure 1: The fundamental polygon of a torus.

Now suppose that 2 sides represented as b are equivalent. Then the natural image of such rectangle can be represented as following.




Figure 2: The cylinder formed by identifying the edges b of the fundamental polygon.

Now, do same thing to the both side a also.




Figure 3: The standard generators a and b on the torus $\mathbb{T}^2$.


Our goal in this section is generalizing this notions into topological words. Let's define and understand bunch of things from now on.

Definition 7.1. Quotient Map

Let X, Y be topological spaces and $p : X \rightarrow Y$ be a surjective map. The map p is said to be a *quotient map* provided a subset U of Y is open in Y if and only if $p^{-1}(U)$ is open in X .

Remark 7.2

Following is the order of strongness of definitions.



Quotient map is continuous map with surjectivity and Embedding is continuous map with injectivity, and Homeomorphism is continuous map with bijectivity.

The if and only if condition above implies that, quotient map p is continuous($\Rightarrow$) and if preimage $p^{-1}(U)$ of something $U \subseteq Y$ is open in X , then U is open in Y . ($\Leftarrow$) **Note** that, there aren't any guarantee that image of any open set in X is also open in Y . We can obtain an alternative way to define quotient maps.

Definition 7.3. Saturated Set

Let X, Y be topological spaces and $p : X \rightarrow Y$ is a surjective map. We say that $C \subseteq X$ is saturated with respect to p if C equals the complete preimage of some subset of Y . i.e., $p^{-1}(p(C)) = C$ holds. Equivalently, C contains all $p^{-1}(\{y\})$ such that intersects with C .

Then now we can say that p is quotient map if and only if p is continuous and every open saturated set in X map to open set in Y . We also say that $f : X \rightarrow Y$ is an *open map* if image of open set in X is open in Y . Then, every surjective continuous open map is quotient map. (First one is stronger notion). Before move on, we summarize such confusing concepts in following way.

Remark 7.4. Continuous, Quotient, Homeomorphism

- If f is continuous, $U \subseteq Y$ is open $\implies$ complete preimage of U is open in X .
- If f is quotient map, there is a bijection between open sets in Y and open complete preimages in X .
- If f is Homeomorphism, there is a bijection between open sets in Y and open sets in X .

Now we can define *quotient topology* and *quotient space* which are main topics in this section.

Definition 7.5. Quotient Topology

X is a space and A is a set. For any $p : X \rightarrow A$ surjective map, there exists unique topology of A such that make p be a quotient map. Such topology is called *quotient topology* induced by p .

Such topology of A is just collection of $U \subseteq A$ such that $p^{-1}(U)$ is open in X . It automatically makes that p be a continuous map, and even quotient map. Intuitively, surjective map can be imagine as a compression of X into smaller set A . Above definition states that, there is the unique ‘shape’ for A that makes this becomes ‘good’ compression. Actually the ‘compression’ of space X is quite non-trivial in above context. We define the *quotient space* in general and see how to describe such compression in topological words.

Definition 7.6. Quotient Space

Let X be a topological space and X^* be a disjoint partition of X . We can define the projection $p : X \rightarrow X^*$ in natural manner, which is surjective map. Then 7.5 says that we can find unique *quotient topology* of X^* . With it, X^* is called quotient space of X .

Then it is not hard to understand that X^* is compressed space of space X . Each element in X^* is a sort of *representation* of its equivalent classes of elements of X . So any $U \subseteq X^*$ is collection of *equivalent classes* and its preimage $p^{-1}(U)$ is open.

As we stated in first of this section, now we can generalize the constructing torus from 2D rectangle. Let

$$X = [0, 1] \times [0, 1]$$


Then we can define X^* by:

$$X^* = \begin{cases} \{(x, y)\} & (x, y) \in (0, 1) \times (0, 1) \\ \{(0, y), (1, y)\} & y \in (0, 1) \\ \{(x, 0), (x, 1)\} & x \in (0, 1) \\ \{(0, 0), (0, 1), (1, 0), (1, 1)\} \end{cases}$$

Then X^* becomes a torus and we can pull-back open sets on torus into open sets on rectangle. We state 2 theorems related to continuity on quotient maps.

Theorem 7.7. Continuity of Quotient Map - Mapping Out

Let $p : X \rightarrow Y$ be a quotient map. Let Z be a space and let $g : X \rightarrow Z$ be a map such that constant on each $p^{-1}(y)$ for $y \in Y$. Then, g induces a map $f : Y \rightarrow Z$ such that $f \circ p = g$. f is continuous if and only if g is continuous, f is quotient map if and only if g is quotient map.



Proof. f can be well-defined as following way:

$$f : y \mapsto g(p^{-1}(y))$$

Then for any $x \in X$,

$$f \circ p(x) = g(p^{-1}p(x)) = g(x)$$

Then it is clear that if f is continuous(quotient map), then g is continuous(quotient map). Now suppose that g is continuous. Let U be any open set in Z . Then, $g^{-1}(U)$ is open in X . However, $g^{-1}(U)$ is complete preimage of some subset V in Y so $p^{-1}(V) = g^{-1}(U)$. Since p is quotient map, V is open in Y . However, $V = p \circ g^{-1}(U) = f^{-1}(U)$ is open in Y , so f is continuous. $\square$


Corollary 7.8

Let $g : X \rightarrow Z$ be a surjective continuous map. Let X^* be the following collection of subsets of X :

$$X^* = \{g^{-1}(z) \mid z \in Z\}$$

Give X^* the quotient topology.

- The map g induces a bijective continuous map $f : X^* \rightarrow Z$, which is a homeomorphism if and only if g is a quotient map.



- If Z is Hausdorff, so is X^* .

Proof. Since g is surjective, $f = g \circ p^{-1}$ clearly forms the bijection between X^* and Z . Suppose that f is a homeomorphism and $g^{-1}(U)$ is open for some subset U of Z . Since f is bijection, there exists V in X^* such that $f(V) = U$. Then by definition of

f , $p^{-1}(V) = g^{-1}(U)$. So $p^{-1}(V)$ is open in X , implies that V is open in X^* . Since f is homeomorphism, U is open in Z . Suppose that g is a quotient map. Then for any open set U in X , $p^{-1}(U)$ and $g^{-1}(U)$ both are open in X^*, Z respectively. However, they bijects via f , so f is homeomorphism. If Z is Hausdorff, X^* is clearly Hausdorff. $\square$

8 Connected Spaces

From now on, our main focus is on defining and analyzing ‘good’ shapes and spaces in general. First of all, we can imagine the good properties in $\mathbb{R}^n$ for example, *Maximum Value Theorem*, *Intermediate Value Theorem*, *Uniform Continuity Theorem* so on. However, in $\mathbb{R}$, they holds for closed interval $[a, b]$. So we can imagine that $[a, b]$ is somewhat good-shaped subspace of the good space $\mathbb{R}$. Since the condition for ‘good’ can be differ for each properties, so we want to generalize them as much as possible. The first concept is quite natural, which is called *connectedness*.

Definition 8.1. Seperation and Connected

Let X be a topological space. We say that U, V is a *seperation* of X if $U, V \neq \emptyset$, $X = U \cup V$ and U, V are open set in X . If such U, V are not exists in X , then we say that X is *connected* space.

Connectedness is obviously topological invariant. i.e., any Homeomorphism preserves the connectivity. This is clear since any Homeomorphisms preserves the seperation. We can define connected space slightly differently.

Lemma 8.2

X is connected if and only if clopen sets in X are only the trivial ones(i.e., if $A \subseteq X$ is both closed and open, then $A = \emptyset$ or X).

Proof. Existance of clopen set is equivalent to existance of seperation of X . $\square$

Lemma 8.3. Subspace Connectivity

If Y is a subspace of X , a seperation of Y is a pair of disjoint nonempty sets A and B whose union is Y , neither of which contains a limit point of the other. The space Y is connected if and only if there exists no seperation of Y .

Proof. Suppose that Y is not connected, then there exists $A, B \neq \emptyset$, $A \cup B = Y$ and A, B are open in Y . Then, A, B are also closed in Y . So, by 3.6,

$$A = \text{closure of } A \text{ in } Y = \bar{A} \cap Y$$

And similarly for B , we can obtain that A, B contains all of its limit points in themselves. So, A, B is the seperation of Y . Suppose that A, B are a seperation of Y . Then we can obtain A, B are both closed and open in Y , so Y is not connected. $\square$

Here, we can imagine that not only elements but also a limit point acts role of *connection port* of a set. First, we can state the notion of *connection port* of elements as following lemmata.

Lemma 8.4. Connected Subspaces

If the sets C and D form a separation of X , and if Y is a connected subspace of X , then Y lies entirely within either C or D .

Proof. Suppose not, then $C \cap Y, D \cap Y \neq \emptyset$ and $(C \cap Y) \cap (D \cap Y) = \emptyset$ and both are open in Y , moreover $(C \cap Y) \cup (D \cap Y) = Y$. So, contradiction to Y is connected subspace. $\square$

Lemma 8.5. Connection Port - Elements

The union of connected subspaces that **point in common** is connected.

Proof. Let $\{A_\alpha\}_{\alpha \in J}$ be a family of connected subspaces of X and there exists $x \in X$ such that $x \in A_\alpha$ for any $\alpha \in J$. Suppose that $\bigcup A_\alpha$ is not connected. Then we can find $\bigcup A_\alpha = C \cup D$ where C, D forms separation of $\bigcup A_\alpha$. W.L.O.G., $x \in C$. Then, according to ??, every $A_\alpha \subseteq C$ so $D = \emptyset$ which is contradiction. $\square$

Now we state the notion of connection port of limit points as following theorem.

Theorem 8.6. Connection Port - Limit Points

Let A be connected subspace of X . For $A \subseteq B \subseteq \bar{A}$, then B is connected subspace.

Proof. Suppose that B is not connected. Then, we can find separation U, V of B such that U contains all its limit points in B and V also. However since $A \subseteq B$, according to 8.4 A entirely in U without loss of generality. Then all elements in V is $B \setminus A \subseteq A'$. But they are limit points of $A \subseteq U$, contradiction. $\square$

Now we'll see important results for connected spaces. The big question is, **what preserves the connectedness?**

Theorem 8.7. Continuous Map Preserves Connectedness

The image of connected set under continuous map is connected.

Proof. Let $f : X \rightarrow Y$ is continuous map and X is connected space. For convenience, let $Y = \text{Im}(X)$. If Y is not connected, there are U, V forming separation of Y . Since they are open, $f^{-1}(U), f^{-1}(V)$ also open in X . And they are non-empty and disjoint partition of X , contradiction to X is connected. $\square$

Theorem 8.8. Finite Cartesian Product of connected spaces

A finite Cartesian product of connected spaces is connected.

Proof. It is enough to show that if X, Y are connected spaces, then $X \times Y$ is connected. Pick one point $a \times b$ in $X \times Y$. For any $x \in X$, since $x \times Y \cong Y$ and $X \times b \cong X$, both

of $x \times Y, X \times b$ are connected. Then 8.5 implies that following subspace is connected:

$$T_x = (X \times b) \cup (x \times Y)$$

Now, form the union $\bigcup_{x \in X} T_x$ then this is also connected since $a \times b \in T_x$ for any $x \in X$ and then 8.5. However, $\bigcup_{x \in X} T_x = X \times Y$. So $X \times Y$ is connected. $\square$

Here are some interesting results. As we referred in 6.13, when we consider the arbitrary cartesian products, the connectedness depends on what topology we take for the space.

Theorem 8.9. Box Topology and Product Topology, in connectedness

Let $\{X_\alpha\}_{\alpha \in J}$ is family of connected spaces. $\prod_{\alpha \in J} X_\alpha$ is connected under product topology, but not always true under box topology.

The example is quite tricky. We can imagine $\mathbb{R}^\omega$ as a set of sequences on $\mathbb{R}$. Then, we can categorize such sequences by:

A : Bounded sequences , B : Unbounded sequences

Then $A \cap B = \emptyset$, $A \cup B = \mathbb{R}^\omega$. Now we claim that A, B both are open. For sequence $\{x_n\}_{n \in \mathbb{N}}$, construct

$$U = (x_1 - 1, x_1 + 1) \times (x_2 - 1, x_2 + 1) \times \cdots$$

Then U is open in box topology, and $U \subseteq A$ if $\{x_n\}$ is bounded, and $U \subseteq B$ if $\{x_n\}$ is unbounded. So A, B both are open and $\mathbb{R}^\omega$ is not connected. Actually this example shows that second part of 8.9.

Proof. Choose $(x_\alpha)_{\alpha \in J}$ in $\prod X_\alpha$. We can define Y by set of $(y_\alpha)_{\alpha \in J}$ such that $y_\alpha \neq x_\alpha$ for finitely many $\alpha \in J$. Then, we can define X_K by $\prod H_\alpha$ such that $H_\alpha = X_\alpha$ if $\alpha \in K$ and $H_\alpha = \{x_\alpha\}$ if not. Then $X_K \cong \prod_{\alpha \in K} X_\alpha$ Here if K is finite, then $\prod_{\alpha \in K} X_\alpha$ is connected by 8.8 and so for X_K . Note that $Y = \bigcup_{K \subseteq J, |K| < \infty} X_K$ and each X_K has common point (x_α) . So Y is connected. Now suppose any basis element $\prod_{\alpha \in L, |L| < \infty} U_\alpha$ for $\prod X_\alpha$. In such basis element, we can always find

$$(z_\alpha) = \begin{cases} x_\alpha & \alpha \notin L \\ z_\alpha & \alpha \in L \end{cases}$$

Then it is clear that $(z_\alpha) \in \prod_{\alpha \in L, |L| < \infty} U_\alpha$ and $(z_\alpha) \in Y$. So, every basis of $\prod X_\alpha$ intersects Y . Means that every points of $\prod X_\alpha$ is limit points of Y , implies that $\bar{Y} = \prod X_\alpha$. However Y is connected, so $\bar{Y}$ also connected by 8.6. So $\prod X_\alpha$ also connected. $\square$

9 Connected Subspaces of The Real Line

In this section, we state full-of intuitive parts about connectedness. We intuitively knows that any intervals and rays are connected in $\mathbb{R}$. Such connected sets in $\mathbb{R}$ has ‘good’ properties, like *intermediate value theorem*. Our goal is find the general condition that makes our space holds *intermediate value theorem*, like $\mathbb{R}$. And also, state another fully intuitive notion called *path connected*.

Definition 9.1. Linear continuum

A simply ordered set L non-empty is called a *linear continuum* if following holds:

1. L has the least upper bound property.
2. If $x < y$, exists z such that $x < z < y$.

Where least upper bound property means that, if $A \subseteq L$ is bounded above, then $\sup A \in L$.

This is the target property that our general space behaviors like $\mathbb{R}$ (in connectedness’ perspective). First $\mathbb{R}$ -like behavior is following.

Theorem 9.2. Convex Connectedness

Let L is a linear continuum with ordered topology. Any convex set Y in L is connected. So, L itself, intervals, and rays are connected.

Proof. Suppose that our convex set $Y \subseteq L$ is not connected. Then Y has a separation A, B . Let’s choose $a \in A, b \in B$ and $a < b$ without loss of generality. Since Y is convex, $[a, b] \subseteq Y$. Let’s denote

$$A_0 = [a, b] \cap A, \quad B_0 = [a, b] \cap B$$

However, $[a, b] \subseteq Y \subseteq L$ also supspace and A_0, B_0 are non-empty open sets in $[a, b]$, which union is $[a, b]$. So A_0, B_0 forms separation of $[a, b]$. Now, let $x = \sup A_0$. This is well-defined since A_0 is bounded and L is linear continuum. And $a \leq x \leq b$, so $x \in [a, b]$. Suppose that $x \in B_0$. Since B_0 is open, we can find small open interval for x in B_0 . See it partly, we obtain $(d, x] \subseteq B_0$. Then d is smaller upperbound than x , contradiction. Vice versa, Y is connected. $\square$

Now it’s time to generalize one of the famous theorem in real analysis, which is *intermediate value theorem*. The condition is much generalized, f is continuous, X is connected, Y is order space.

Theorem 9.3. The Intermediate Value Theorem

Let $f : X \rightarrow Y$ be a continuous map where X is connected and Y is order space. For $a, b \in X$, if $f(a) < r < f(b)$ then there exists $c \in X$ such that $f(c) = r$.

Proof. Suppose that such c is not exists. Then, $r \notin \text{Im}(X)$ so we can separate $\text{Im}(X)$ by

$$A_0 = \text{Im}(X) \cap (-\infty, r), \quad B_0 = \text{Im}(X) \cap (r, \infty)$$

Since A_0, B_0 is open in $\text{Im}(X)$ and non-empty disjoint, they are separation of $\text{Im}(X)$. However, the image of connected space under continuous map is connected by 8.7. So this is contradiction, and such c is exists. $\square$

Now we define another version of connectedness, which is called *path connected*. This is drawble concepts, but not exactly equivalent to what we defined.

Definition 9.4. Path Connected

Let X be a topological space. We define a *path* between $x, y \in X$ by a continuous mapping $f : [a, b] \rightarrow X$ such that $f(a) = x, f(b) = y$. Where $[a, b] \subseteq \mathbb{R}$. X is called *path connected* if for every $x, y \in X$, there exists a path from x to y .

Theorem 9.5. Path Connected Implies Connected

If X is path connected space, then X is connected space.

Proof. Suppose that X is not connected so there is a separation A, B of X . Then since every path is image of convex(so, connected) space, *path is connected*. So by 8.4, path is entirely in A or B . Then, we can't find a path between $a \in A$ and $b \in B$, contradiction to X is path connected. So, X is connected. $\square$

Proposition 9.6. Connected Does Not Implies Path Connected

Let $S = \{x \cdot \sin(1/x) \mid 0 < x \leq 1\}$. $\bar{S}$ is connected but not path connected.

Proof. Since S is image of $(0, 1]$ under continuous map, so S is connected. This implies that $\bar{S}$ is connected by 8.6. Note that $\bar{S} = S \cup (0 \times [-1, 1])$. There is not any path between $a \in S, b \in 0 \times [-1, 1]$. If there exists a path, $f : [0, 1] \rightarrow \bar{S}$ such that $f(0) = b$ and $f(1) = a$ is continuous. However, it oscillate in $[-1, 1]$ near $x = 0$, continuity is not hold near $x = 0$. $\square$

10 Compact Space

When we say something is ‘compact’, it means that ‘Wow, that’s very good or suitable’. According to our textbook page 161, Munkres[1] refer good historical development of the word ‘compact’ in topology. First, in $\mathbb{R}$, we observe that any closed interval has very good properties like uniform continuity theorem and maximum value theorem. After that, mathematicians like Weierstrass observe that every infinite subsets of $[a, b]$ has a limit point. Later, finally we find the core generalization via open coverings.

Definition 10.1. Open Covering

References

- [1] James Munkres. Topology, 2014.