




Sowman
2015/09/06

Lab 1

Measurement, Uncertainty, and Temperature

Equipment List

- Leslie Cube
- Glass Thermometer
- IR Thermometer "gun" (2)
- Stopwatch
- Ruler
- Computer

Overview

In this weekly lab you will primarily work in pairs, with your "partner" at the same table, but feel free to discuss things with the "group" of four at your adjoining tables. We will also share data with the entire class, usually using google surveys and sheets. You will turn in your own individual lab assignment, with copies of all data and graphs, in a single pdf document uploaded to the Blackboard "assignment" for that week.

1.1 Introduction and Definitions

Today we will conduct our first measurements of physical phenomena, mostly temperature measurements, and understand the necessity of the uncertainty and units associated with those measurements.

A **physical measurement** contains three equally important parts: the numerical **value**, the **uncertainty** (also referred to as *error*) in that value, and the **units** in which these are measured.

Some examples are:

- The human body temperature is 37.0 ± 0.5 °C.
(Uncertainty represented as "plus or minus" makes a range of values. Both the value and uncertainty are in "degrees celsius".)

- A college basketball game has 40 minutes ± 0.1 seconds of playing time.
(The final minute clock measures to tenths of a second. The units can be different between the value and uncertainty.)
- The Sun is 93 million miles away
Here the uncertainty is expressed *implicitly* as “plus or minus a few of the last digit given”, so plus or minus a few million miles

Let's make a precise definition of the uncertainty/error in a measurement:

Given a particular measurement, the **uncertainty** is a statement of the range of values in which another scientist is likely¹ to make the same measurement.

There are a number of sources/components of this uncertainty:

- **Instrumental Uncertainty:** A measurement device generally has a smallest unit division shown. For an analog device — like a ruler with millimeter marks, or a thermometer with degrees — where we might hope to measure between the smallest markings, we can specify the instrumental uncertainty as such (e.g., half of a millimeter on a ruler, or 0.3°C on a thermometer). For a digital device, we can say that the instrumental uncertainty is “plus or minus a few of the last digit” (e.g., ± 0.03 seconds on a standard stopwatch).
- **(Manufacturing) Instrumental Uncertainty:** Sometimes although an instrument shows a precise demarcation, it is clear from looking at multiple “identical” instruments that they were not manufactured to that precision (e.g., multiple identical thermometers read slightly different values). The range of manufacturing uncertainty can be assessed by using multiple copies of the same device.
- **Methodological Uncertainty:** Sometimes our uncertainty is limited more by the way we make a measurement than by the instrument itself. For example, although a stopwatch has instrumental uncertainty in the hundredths of a second, our human reaction time is no better than a few *tenths* of a second, so our methodological uncertainty would be ± 0.3 s. This can be assessed by making repeated measurements.
- **Natural Variation** Almost always, the object of measurement does not have an infinitely precise value (e.g., the “temperature outside right now” depends on where exactly you measure it, and at which moment in time; the “diameter of a rock” that is not spherical). This uncertainty can sometimes be limited by a more precise specification of the measurement (e.g., “the temperature in the shade, outside Padnos, at 10:40 am”; or “the largest diameter of the rock”). And, like the methodological uncertainty, it can be assessed by repeated measurements.

1.2 Experiments

1.2.1 The time of my name

- Have your partner recite their full name to you at a steady relaxed pace. Measure the amount of time this takes using the stopwatch. You can try this multiple times. Then

¹More precisely, if we say that the uncertainty is the *standard deviation* over repeated measurements, then 68% of all measurements fall in that range

switch and recite your own name for them.

Questions

1. Write down the name, and then your time measurement with uncertainty and units.

"Benjamin Huser"

$$2.5 \text{ s} \pm 0.3 \text{ s}$$

2. Briefly explain how you arrived at the uncertainty value, making mention of all "sources" of uncertainty.

SHOULD HAVE PERFORMED
MULTIPLE MEASUREMENTS
(w/ DATA SHOWN) AND
GIVEN UNC. FROM SCATTER

- INSTRUMENTAL: $\pm 0.03 \text{ s}$
DOESN'T AFFECT OVERALL
- METHOD: MAIN ISSUE
- NAT. VARIATION: COULD BE ISSUE
LOOK @ SPREAD IN
VALUE & DISCUSS.


1.2.2 "Seeing" the uncertainty I: Room Temperature

One table has been set up with an array of thermometers. Since they are all localized in the same place, they should be making approximately the same measurement: the temperature of the air at that table right now.

- Take a careful clear photo of the array of thermometers, then return to your desk and write down each of the temperature measurements on a sheet of paper.
- Considering the range of measurements you have, put labeled tick marks on the axis below such that the values will be spread out across the line. Use a ruler to make equal spacing of tick marks. At the far right, label that axis (e.g., "Temperature (C)").
- Place a dot on/near the axis for every measurement of temperature. You can "scatter" these a bit above/below the axis for clarity.

22.5
22.0
21.5
20.5
20.0
21.0
20.5

DATA



Questions


1. Express the full measurement with value, uncertainty, units. Estimate the uncertainty using your graphical display of points.

$$21.2 \pm 0.8 \text{ } ^\circ\text{C}$$

MAKE ONLY USE THE ANALOG?

2. What is the instrumental uncertainty for this thermometer?

"THE ANALOG"



$$\pm 0.3 \text{ } ^\circ\text{C}$$

(WITHIN SMALLEST SPACING)

3. What do you think is the primary source of the uncertainty? Mention each type of uncertainty source.

PROBABLY:

- MANUFACTURING UNCERTAINTY (SLIGHT DIFF'S OF SAME MEASUREMENT)

OR:

- NATURAL VARIATION: TEMP. COULD BE DIFF @ DIFF POINTS ON DESK

1.2.3 "Seeing" the uncertainty II: Body Temperature

Here each of you will make a measurement of your body temperature using the Infrared (IR) thermometer "gun" on the palm of your hand. Note the picture on the side of the IR thermometer gun describing its "beam" of measurement (imagine a spot being projected onto the thing you are measuring, and make sure the spot is contained within the object). You will produce a two-picture plot, using a provided code (see Blackboard) and the Google CoLab implementation of python programming language.

1. What is the instrumental uncertainty associated to the IR thermometer?

$$\pm 0.1 \text{ } ^\circ\text{C} \text{ (or } \pm 0.3 \text{ } ^\circ\text{C} \text{ ?) } \text{ FROM DIG. READOUT}$$

2. Do you observe any natural variation of temperature across your hand?

$$\text{YES: } 34.1 \text{ } ^\circ\text{C} - 34.6 \text{ } ^\circ\text{C} , \pm 0.3 \text{ } ^\circ\text{C}$$

3. Write down your measurement (with value, uncertainty, and units) here, and then submit it to the google survey (see Blackboard).

$$\underline{34.3 \pm 0.3 \text{ } ^\circ\text{C}}$$

OR, BETTER, AN EXPERIMENT

MAKE FIGURE FOR THIS & DISCUSS IN CLASS?

ADD INSTRUCTIONS TO DISCUSS & MAKE QUANTITATIVE

ADD QUESTION "HOW DID YOU DETERMINE UNCERTAINTY?" (SHOW MULT MEASUREMENTS?)

ON BUENBOARD:

Lab 01 --- Measurement, Uncertainty, Temperature

- **Data:** [Human Temperature \(Google Form\)](#) and [Results](#)
- **Data:** [Other Temperature Data \(Google Form\)](#) and [Results](#)
- **Code:** Scatter and Histogram
 - [First block of code](#)
 - [Second block of code](#)

Google Form:

Lab 01 --- Human Body Temperature

hulderbg@gmail.com [Switch account](#)

Not shared

* Indicates required question

Name (or nickname/coded name if you don't want) *

Your answer

Temperature *

Your answer

Uncertainty *

Your answer

Units *

Your answer

Submit Clear form

Results:

Timestamp	Name (or nickname/coded name if you don't want)	Temperature	Uncertainty	Units
8/27/2025 9:51:08	Anna	31.5	0.2	Degrees celsius
8/27/2025 9:52:03	Josiah Van Engen	29	0.5	Degrees Celsius
8/27/2025 9:52:47	Emma T	32.4	0.1	Celsius
8/27/2025 9:55:20	Aylah Smith	32.6	0.2	Celsius
8/27/2025 9:57:39	Jenna	33.9, 31.0, 32.7, + 32.5	+ 00.5 degrees Celsius	Celsius
8/27/2025 9:59:23	Matteo Lambardi	33.5 degrees Celsius	plus/minus 0.5	degrees Celsius
8/27/2025 9:59:49	George Rousch	30.8	+ or - 0.3	c
8/27/2025 10:01:02	Charlie Baldwin	33	0.67	Degrees Celsius
8/27/2025 10:01:23	Oleksandr	34.2, 34.3, 34.4, 34.8, 35	0.46	C
8/27/2025 10:03:01	Avery Palka	33 C	+/- 0.4 C	Degrees Celsius
8/27/2025 10:03:22	Joel K	29.6	1.3	Celsius
8/27/2025 10:03:34	Mary Konnie	33.3	0.3	°C
8/27/2025 10:04:14	Kendall Noakes	33.3	0.2	Celsius
8/27/2025 10:04:41	lj	32.5,C	27.8C	4.7
8/27/2025 10:05:11	Trinity LaFlore	33.64	1.38	Celsius
8/27/2025 10:05:27	Addie	34.6°C	+/- 0.3°C	Degrees Celsius
8/27/2025 10:05:33	Clare	34.4	+/- 0.1	Degrees Celsius

SINGLE CSV w/ ONLY TEMP DATA

Untitled spreadsheet

File Edit View Insert Format D

C5

	A	B
1	31.5	
2	29	
3	32.4	
4	32.6	
5	33.9, 31.0, 32.7, = 32.5	
6	33.5 degrees Celcius	
7	30.8	
8	33	
9	34.2, 34.3, 34.4, 34.8, 35.0	
10	33 C	
11	29.6	
12	33.3	
13	33.3	
14	32.5,C	
15	33.64	
16	34.6°C	
17	34.4	
18		

CLEAN
THE
DATASET
→
(SINGLE NUM.
VALUE PER ROW)

Untitled spreadsheet


File Edit View Insert Format Data Tools

A17

	A	B	C
1	31.5		
2	29		
3	32.4		
4	32.6		
5	32.5		
6	33.5		
7	30.8		
8	33		
9	34.6		
10	33		
11	29.6		
12	33.3		
13	33.3		
14	32.5		
15	33.64		
16	34.6		
17	34.4		
18			

FILE/DOWNLOAD
⇒ UNTITLED...CSV
IN "DOWNLOADED"

Google Colab



```
import numpy as np
import pandas as pd
import matplotlib.pyplot as plt


def plot_csv_column(filename, xlabel='Value', jitter_frac=0.02):
    # Read first column (works with or without header)
    df = pd.read_csv(filename)
    col = df.iloc[:, 0].dropna().astype(float).to_numpy()
    if col.size == 0:
        raise ValueError("No numeric data found in first column.")

    # compute bins via Freedman-Diaconis; fallback to sqrt(n) bins if needed
    try:
        bins = np.histogram_bin_edges(col, bins='fd')
        if len(bins) <= 1:
            raise ValueError
    except Exception:
        nbins = max(1, int(np.sqrt(col.size)))
        bins = nbins

    # plotting
    fig, (ax_top, ax_bot) = plt.subplots(2, 1, sharex=True, figsize=(8, 4), constrained_layout=True)

    # top: horizontal scatter with small vertical jitter so points don't overlap
    rng = np.ptp(col)
    jitter = jitter_frac * (rng if rng > 0 else 1.0)
    y = np.random.uniform(-jitter, jitter, size=col.size)
    ax_top.scatter(col, y, s=18, alpha=0.7)
    ax_top.set_yticks([])
    ax_top.set_xlabel('Data (horizontal)')

    # bottom: histogram using bins computed above
    ax_bot.hist(col, bins=bins, edgecolor='black', alpha=0.75)
    ax_bot.set_xlabel(xlabel)
    ax_bot.set_ylabel('Count')
    ax_bot.set_title('Histogram')
```



RIGHT-CLICK
TO SAVE
IMAGE
(AFTER ADJUSTING
TITLES & AXIS
LABELS)

- Copy the column of "values" data for the whole class in the linked spreadsheet (see Blackboard) into your own spreadsheet and save as a csv file.
- In a web browser, go to colab.research.google.com and open a "New Notebook".
- Paste in the code provided on blackboard for *Scatter and Histogram Plot*.
- To run the code, either press the "play" button, or press shift-enter/shift-return. It will prompt you for your csv file to upload.
- This will produce a plot on the screen.
- Notice that there are generic axis labels and titles. Find the place in the code where they have been entered and edit them to make them specific to your particular data.
- To save and print your plot, you can either right-click and save to a file or screenshot to a file. Print this plot so that you can add it to your submission.

→ DONE?

→ PLOT TURNED IN?

1.2.4 Leslie Cube

- Fill up the Leslie cube with hot/warm water at the sink and then replace the rubber stopper. Fill a small beaker with cold water.
 - Place the glass thermometer first in the cold water, and then insert it into the Leslie cube such that its bulb is in the middle of the cube.
1. Record the glass thermometer measurements of the temperature in the cold water and in the Leslie cube.

REMEMBER THEM

< SHOULD HAVE: VALUE ± UNC UNITS >

(COLD & HOT MEASUREMENTS)

2. Describe how quickly the thermometer came to equilibrium with the cold water and with the hot water.

JUST SOME QUALITATIVE COMMENTS, E.G.,

"WITHIN A FEW SECONDS"

"QUICKLY THEN SLOW FOR LAST APPROACH"

3. Take and record temperature measurements of each side of the Leslie cube using the IR thermometer gun.

< VALUE ± UNC °C > FOUR TIMES

(CAN SHOW MULT MEASUREMENTS
→ UNC FOR EACH)

MAYBE
MOVE
SUMM.
QUEST
HERE?

1.2.5 Temperatures of other objects

- Using the IR thermometer gun, record measurements of temperature (with value, uncertainty, units for each) for some of the following objects:
 - The surface of the table you sitting at.
 - Something else in the room.

MAKE CLEAR
THAT EACH
STUDENT SHOULD
MEASURE & RECORD,
NOT EACH GROUP.

REMEMBER TAKEN ABOUT
UNC. &
UNITS

- Outside: Something that might be in equilibrium with the air temperature (shaded spot).
 - Outside: Something that is probably not in equilibrium with the air temperature.
 - Outside: The sky (try both clear and clouded spots).
 - Record your temperature measurements below and also report them to the google survey posted on Blackboard.
1. Record your temperature measurements here. *ADD "BRIEFLY DESCRIBE HOW YOU OBTAINED UNCERT. VALUES"*

TEMP MEASUREMENTS w/ UNC. ESTIMATES
FOR ALL OF ABOVE (5 "OBJECTS")

Timestamp	Nar	Temperatur	Uncertain	Units	Outside /	Uncertainty	Units	Sky Tem	Uncertain	Units
8/27/2025 10:24:36	Mary Kc	23.1	0.1	°C	21	0.2	°C	-15	4.3	°C
8/27/2025 10:24:56	George	22.5	+ or - 0.2	C	15.1	+ or - 0.2	C	-19.6	+ or - 0.3	C
8/27/2025 10:25:58	Oleksan	25.1	0.3	C	17.4	0.4	C	-12.5	-2	C
8/27/2025 10:26:26	Addie	23.5°C	+/-1°C	Degrees Celsius	25°C	+/-5°C	Degrees Celsius	-18.3°C	+/-0.5°C	Degrees Celsius
8/27/2025 10:27:05	Trinity	23	0.1	Celsius	19.45	2.74	celsius	-2.4	10	celsius
8/27/2025 10:27:13	Anna	23.5	0.1	degrees celcius	19.4	0.1	degrees celcius	-16.8	0.1	degrees celcius
8/27/2025 10:42:37	bj	23.9	0.65	C	15.8	0.1	C	14.1	3.8	C
8/29/2025 13:23:46	Matteo	24	plus/minus 0.	Celcius	18	plus/minus	0.5	-17	plus/minus 0.4	Celcius

ADD: Print & SHOW THE RESULTING PLOT

Summary Questions

Brief answers are fine, but please write in complete sentences.

SHOULD ALSO ASK SPECIFICALLY:
 — DIFF. BOWN CUBE & GLASS
 — DIFF BOWN SIDES

1. State the temperature measurements you made for the Leslie cube, including each type of side and the direct glass thermometer measurement. Speculate on why you found different values, if you did. → GIVE ALL 4 MEASUREMENTS w/ UNC. + GLASS (5 TOTAL)

→ SHOULD SEE REFLECTIVE SIDE GIVES LOW MEASUREMENT

→ THREE SIDES WITHIN UNC. ONE IS NOT (REFLECTIVE IS GIVING YOUR TEMP?)

"BE QUANTITATIVE"

2. Looking at either the hand-temperature data or the other plot you made with the entire class data: how did your own estimated uncertainty (for your measured value) compare to the "spread" of the data taken by all students? Should these agree? Why or why not?

• STATE YOUR OWN UNCERTAINTY

• STATE UNC. FROM SPREAD (HALF OF IT) IN OTHERS DATA.

• YES THEY SHOULD OVERLAP BASED ON THE DEFN OF UNC.

3. Two students make a measurement of the distance from the Bookstore to the Clock Tower. They are reported as follows

$$199.65 \text{ m} \pm 0.3 \text{ m}$$

$$202 \text{ m} \pm 3 \text{ m}$$

- (i) On the axis below, draw tick marks over a wide enough range to plot both measurements *and their ranges of uncertainty*. (ii) Plot the two measurements using a point for the "value" and a horizontal bar to show the uncertainty. (iii) Are these "the same" measurements to within uncertainty or not? Explain.



YES THEY OVERLAP IN UNC. RANGES ...

EACH MEASUREMENT FALLS IN RANGE OF OTHER
(SEE DEFIN OF UNC)

4. Shown below is a summary of the Global Average Temperature over most of human recorded history. The uncertainties for each data point are indicated by bars. Speculate on how these uncertainties might have been determined, how they change over time, and why.

"BY HIGHLIGHTING THREE 'BARS' AT DIFF. TIMES ..."

3 (i)-(ii)
GOOD RANGE?
GOOD SPACING?
UNITS?



- UNC. PROBS. CAME FROM SPREAD IN VALUES OF INDIVIDUAL DIFF. METHODS IN CALCULATING THAT AVERAGE

- UNC. IS DEFINITELY DECREASING OVER TIME (MORE TEMP VALUES COLLECTED)

PROBS NOT MUCH CHANGE IN, E.G. INSTRUM. UNC.