


# Guidelines for RNA-seq projects: applications and opportunities in non-model decapod crustacean species

Tuan Viet Nguyen  · Hyungtaek Jung · Guiomar Rotllant · David Hurwood · Peter Mather · Tomer Ventura

Received: 11 October 2017 / Revised: 14 May 2018 / Accepted: 11 June 2018  
© Springer International Publishing AG, part of Springer Nature 2018

**Abstract** Next-generation sequencing (NGS) has dramatically changed the way biological research is being conducted in the post-genomic era, and they have only been utilized widely over the recent decade for studies of non-model decapod crustacean species, predominantly by sequencing the transcriptome of various tissues across different life stages. Next-generation sequencing can now provide a rapid, cost-effective solution for discovery of genetic markers crucial in many applications that would previously have otherwise taken years to develop. Sequencing of

the entire transcriptome (referred to as RNA sequencing; RNA-seq) is one of the most popular NGS tools. RNA-seq studies of non-model species in crustacean taxa, however, have faced some problems, including a lack of “good” experimental study design, a relative paucity of gene annotations, combined with limited knowledge of genomic technologies and analyses. The aim of the current review is to assist crustacean biologists to develop a better appreciation for the applications and scope of RNA-seq analysis, understand the basic requirements for optimal RNA-seq studies and provide an overview of each step, from RNA-seq experimental design to bioinformatics approaches to data analysis. Insights that have resulted from RNA-seq studies across a wide range of non-model decapod species are also summarized.

---

Guest editors: Guiomar Rotllant, Ferran Palero, Peter Mather, Heather Bracken-Grissom & Begoña Santos / Crustacean Genomics

---

Tuan Viet Nguyen and Hyungtaek Jung have contributed equally to this work.

---

T. V. Nguyen (✉) · T. Ventura (✉)  
GeneCology Research Centre, Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast, 4 Locked Bag, Maroochydore, QLD 4558, Australia  
e-mail: tnguyen@usc.edu.au

T. Ventura  
e-mail: tventura@usc.edu.au

H. Jung  
Centre for Tropical Crops and Biocommodities, Institute for Future Environment, Queensland University of Technology, GPO Box 2434, Brisbane, QLD 4001, Australia

G. Rotllant  
Institut de Ciències del Mar (CSIC), Passaig Marítim de la Barceloneta 37, 08003 Barcelona, Spain

D. Hurwood · P. Mather  
School of Earth, Environment and Biological Sciences, Science and Engineering Faculty, Queensland University of Technology, GPO Box 2434, Brisbane, QLD 4001, Australia

**Keywords** RNA-sequencing · Next-generation sequencing · Differential gene expression · Shrimp · Prawn · Crab · Lobster · Crayfish · In silico

## Abbreviations

<i>P. trituberculatus</i>	<i>Portunus trituberculatus</i>
<i>S. henanense</i>	<i>Sinopotamon henanense</i>
<i>P. vannamei</i>	<i>Penaeus vannamei</i>
<i>P. monodon</i>	<i>Penaeus monodon</i>
<i>M. japonicus</i>	<i>Marsupenaeus japonicus</i>
<i>P. virginalis</i>	<i>Procambarus virginalis</i> or <i>Procambarus fallax forma virginalis</i>
<i>S. olivacea</i>	<i>Scylla olivacea</i>
<i>S. paramamosain</i>	<i>Scylla paramamosain</i>
<i>E. sinensis</i>	<i>Eriocheir sinensis</i>
<i>N. norvegicus</i>	<i>Nephrops norvegicus</i>
<i>F. merguensis</i>	<i>Fenneropenaeus merguensis</i>
<i>P. clarkii</i>	<i>Procambrarus clarkii</i>
<i>M. rosenbergii</i>	<i>Macrobrachium rosenbergii</i>
<i>S. verreauxi</i>	<i>Sagmariasus verreauxi</i>
<i>N. denticulata</i>	<i>Neocaridina denticulata</i>
<i>P. hawaiiensis</i>	<i>Parhyale hawaiiensis</i>
<i>E. carinicauda</i>	<i>Exopalaemon carinicauda</i>
<i>M. olfersi</i>	<i>Macrobrachium olfersi</i>
<i>P. elegans</i>	<i>Palaemon elegans</i>
<i>P. australiensis</i>	<i>Paratya australiensis</i>
<i>H. rubra</i>	<i>Halocaridina rubra</i>

## Introduction

Next-generation sequencing (NGS) technologies have rapidly transitioned bioscience into the post-genomic era, resulting in easier, cheaper and faster DNA sequencing. Application of advanced NGS platforms has allowed multiple techniques to be developed that address biological challenges. These include RNA-sequencing (RNA-seq), whole-exome sequencing, chromatin immunoprecipitation sequencing (ChIP-seq), microRNA sequencing (miRNA-seq), restriction-assisted DNA sequencing (RAD-seq) and small RNA sequencing. Among these, RNA-seq is a technique that has revolutionized gene-expression studies and marker discovery (single sequence repeats [SSRs]/microsatellites and single nucleotide polymorphisms [SNPs]) (Lister et al., 2009; Wang et al., 2009;

Wilhelm & Landry, 2009; Marguerat & Bähler, 2010; Ozsolak & Milos, 2011; Das et al., 2016; Mykles et al., 2016). The RNA-seq platform is based on the analysis of the transcriptome—a small portion of the whole genome that is transcribed from chromosomal DNA into RNA molecules—a dynamic set of elements that change depending on developmental stages or physiological conditions. Also, by analysing the sequenced transcriptome, genetic polymorphisms including SNPs and SSRs can be mined and analysed with ease (Jung et al., 2011, 2016; Jin et al., 2013; Lv et al., 2014; Meng et al., 2015; Jaramillo et al., 2016; Nguyen et al., 2016).

While RNA-seq techniques have had a major impact on model species (which in this review is defined as a species with a well-characterized genome, e.g. *Daphnia pulex*), the application of RNA-seq approaches in non-model decapod crustacean taxa is still limited by the small size of the research community and the subsequent bottleneck of bioinformatics analysis capabilities. Many NGS analytical tools are available and by default, are developed for model species, making it difficult for researchers investigating non-model organisms to navigate through and identify appropriate tools. Designing and evaluating a pipeline for transcriptomics projects in non-model species, therefore, can be considered a crucial step prior to project initiation. While the transcriptome can encompass many categories of different types of RNA (mi-RNA, small nuclear RNA, non-coding RNA, etc.), this review will focus mainly on mRNA sequencing using Second-Generation Sequencing (SGS) technology—we intend to use the same classification proposed by Schadt et al., (2010), which defined “Sanger sequencing” as the First Generation, “wash-and-scan” sequencing technology as the Second Generation, and single-molecule real-time sequencing as the Third Generation. Under this classification scheme, SGS includes a number of platforms, notably Illumina, Solid, Ion Torrent/Ion Proton, Roche 454, whereas PacBio and Oxford Nanopore are classified as the Third Generation Sequencing (TGS) Technology. In this study, we will focus primarily on different strategies to initiate a transcriptome study, briefly addressing several platforms that currently are available, as well as recommending a number of experimental designs, bioinformatics software for de novo assembly and specific data analyses for decapod

crustacean species. Finally, we review recent biological insights gained from application of SGS in crustacean transcriptomics and highlight opportunities as well as challenges for applied RNA-seq in the future.

## Overview of RNA-Seq technology

### Pre-sequencing

New sequencing technologies and new sequencing chemistries are being developed rapidly. The arrival of SGS, and more recently TGS, has completely changed the way researchers approach unanswered phenomena in basic, applied, and clinical research. Each sequencing platform is based on different proprietary chemistries and technologies and each has unique strengths and weaknesses. Details on sequencing chemistry have been summarized elsewhere (Metzker, 2010; Koboldt et al., 2013; Reuter et al., 2015; Goodwin et al., 2016). Currently, Illumina is the most widely utilized SGS for RNA-seq, since the platform enables deep coverage of the transcriptome and provides long, low-error reads that are suitable for mapping to reference genomes and transcriptome assemblies (Metzker, 2010; Niedringhaus et al., 2011; Goodwin et al. 2016). Performance benchmarking of many SGS platforms has been conducted for several years (Glenn, 2011; Lam et al., 2012; Liu et al., 2012; Finseth & Harrison, 2014; Goodwin et al., 2016; Lahens et al., 2017) and an online archive of sequencing platforms is available on the market and can be found at <https://allseq.com/knowledgebank/>. Given the popularity of Illumina Sequencers in general, it tends to be one of the most widely applied technologies in crustacean transcriptome projects (Havird & Santos, 2016a).


In brief, RNA-seq includes the utilization of an SGS platform to generate a huge amount of sequence data. Due to technical constraints of the approach (most SGS platforms can only generate short to medium length reads, approximately 50–300 bp), RNA transcripts must be fragmented into shorter sequences. In the absence of a reference genome, short reads are then reconstructed to make a reference transcriptome, referred to as a *de novo* assembly. Following this, raw reads can be realigned (or mapped) to the previously generated reference

sequence and counted, thus providing a digital measurement of specific transcript abundances that can facilitate biological interpretation. Where key genes are targeted (based on either high differential expression or previously identified in the literature), they can be validated by replicating samples across a range of experimental conditions (e.g. in different tissue types, at different life history stages, between sexes, etc.). A popular approach for validation includes quantitative real-time PCR (qRT-PCR) where relative transcript abundance can be assessed under more strictly controlled conditions. Most RNA-seq strategies that utilize SGS can be summarized by a basic workflow (Fig. 1).

### Experimental design

Designing an RNA-seq experiment requires a solid biological understanding of the taxa under investigation and the question(s) to be addressed. Poor or inappropriate decisions at this stage can result in a large amount of unusable data. A good experimental design for every NGS-based experiment therefore, is a basic requirement that cannot be over-emphasized.

In general, several factors must be considered prior to the initiation of any well-designed sequencing project. Essentially, an appropriate experimental design is a balance between the level of biological versus technical replication (Fig. 2) and the resulting depth of coverage for each tissue type, life stage, sex etc., within a framework of time and financial constraints. It is advisable that researchers without much prior experience should seek suggestions from professional service providers including bioinformaticians and biostatisticians, as well as the sequencing provider. This review highlights some of the pitfalls to be aware of and sets the scene for appropriate study design. Several studies provide direction on how to design a statistically valid RNA-seq experiment (Auer & Doerge, 2010; Fang & Cui, 2011; Yang & Wei, 2015; Conesa et al., 2016). In general, a comprehensive transcriptome requires multiple tissues from multiple developmental stages, while gene-expression studies require samples that represent contrasting treatments (e.g. male vs. female, control versus hormone treated, salinity vs. freshwater acclimation, or different developmental/life history stages).



**Fig. 1** A simple flowchart of an RNA-seq/transcriptomics study


### *Biological and technical replicates*

In the NGS context, technical replication refers to multiple libraries from the same biological sample (i.e. the technical steps are performed separately) (Fig. 2). While potentially increasing the depth of reads, any variation recorded among technical replicates will also help identify inconsistencies associated with sampling techniques, PCR biases or sequencing errors. In some rare cases, sample collection, storage or processing can be a source of technical variance owing to the relative instability of RNA. It is advisable to employ several randomization techniques during sequencing, for example, multiplexing (mixing of different libraries, each tagged using a different barcode), splitting technical repeats between multiple lanes, or randomization of different libraries in the same lane [an excellent review on statistical randomization for RNA-seq can be found elsewhere (Auer & Doerge, 2010)]. Ultimately, this type of replication provides

some measure of the quality and/or reliability of the analysis.

Biological replication alternatively, relates to different biological samples (e.g. same tissue type but from different individuals) that are processed separately (Fig. 2). Biological replication is desirable since it quantifies natural variation among individuals within the experimental cohort. Furthermore, increasing the sample size (number of biological replicates) not only increases sequencing depth, but also provides greater statistical power to detect differences among treatments where they may exist.

Nevertheless, with a very large sample size, accommodating both technical and biological variation can become very costly and may also result in a complex assay to analyse. When sequencing individuals from a population with large levels of genetic variation, for example when dealing with wild-caught individuals, the more biological replicates, the more likely it is to capture genuine differential expression

**Fig. 2** Technical samples versus biological samples

among groups. In general, most SGS experiments conducted on crustaceans tend to be under-replicated and while there is no gold standard for this matter, it is currently acceptable for RNA-seq experiments to consist of a minimum of three biological samples to provide adequate statistical power; a number of published studies have shown that the power to detect differential expressed genes improves from two samples to five samples per treatment (Kvam et al., 2012; Dillies et al., 2013). Similarly, other studies have proposed that sequencing fewer reads and including more biological replicates is an effective strategy to increase statistical power and accuracy in large-scale differential expression RNA-seq studies (Liu et al., 2014). More recently, results suggest that at least six biological replicates may be needed in more sophisticated RNA-seq experiments and up to 12 replicates per experimental group (Schurch et al., 2016). However, for samples that are very different from each other in terms of transcription level (for example, differential expression profiles between brain versus ovary), less replication may also be acceptable. It is also important to highlight the fact that replicates in an RNA-seq-based study are required for publication in some journals (e.g. refer to Sect. 2.6.7 at [https://www.](https://www.frontiersin.org/about/author-guidelines)

[frontiersin.org/about/author-guidelines](https://www.frontiersin.org/about/author-guidelines)). To conclude, we would recommend maximizing biological replicates to include at least three samples for each experimental condition in every non-model decapod crustacean RNA-seq study.

#### *Choice of sequencing platforms*

There are several sequencing methods that researchers can choose from, including single-end (SE)/paired-end (PE) reads, strand-specific, or non-strand-specific library preparation. The decision on which is selected will be based on the desired outcome of the study but will also depend on budget constraints. For experiments on crustacean species in general, PE sequencing is recommended to obtain a reliable de novo assembly where no reference genome is readily available. Long-read sequencing (e.g. PacBio, Nanopore sequencing), proven to be suitable for enhancing continuity of de novo transcriptome assembly, is currently relatively expensive, and its application has been described elsewhere (Cartolano et al., 2016; Chen et al., 2017; Kuo et al., 2017). Illumina short-read sequencing, however, is by far the most widely used platform for transcriptome sequencing in crustaceans due to its

cost-effectiveness (unit price per nucleotide), fast sequencing times and higher raw read accuracy.

Another consideration is to choose whether stranded sequencing will be needed. In brief, a stranded-specific RNA-seq can retain the gene orientation (sense or antisense transcript). A number of studies have attempted to compare between stranded vs non-stranded approaches, and most have shown that a stranded RNA-seq approach is more advantageous due to its better assembly of unannotated genes, ability to detect genes on the antisense strand as well as improved continuity of transcripts. New de novo assembly programs like Trinity (Grabherr et al., 2011) have a special mode for strand-specific data analysis that has proven to be more effective than non-stranded data (Parkhomchuk et al., 2009; Levin et al., 2010; Zhong et al. 2011; Sultan et al., 2012; Zhao et al., 2015). We therefore recommend strand-specific RNA-seq if possible for non-model decapod crustacean studies (Havird & Santos, 2016a).

#### *Depth of sequence (number of reads)*

The amount of sequencing needed for a given sample is determined by the aims of the experiment, the number of transcribed transcripts and the nature of the species' RNA samples (this is due to the fact that crustacean genomes can be quite complex compared to other invertebrates). To our knowledge, there has been no attempt to investigate the depth required for effective RNA-seq studies in crustaceans. A study of chicken RNA-seq data revealed that approximately 30 million reads (Illumina-75 bp PE) covered all annotated genes, while 10 million reads detected only ~ 80% (Wang et al., 2011). Whereas RNA-seq samples from six different phyla (Annelida, Arthropoda, Chordata, Cnidaria, Ctenophora and Mollusca) has suggested that approximately 20 million reads for tissue samples and 30 million for whole-animal samples were required to provide a good balance between total coverage and noise (Francis et al., 2013). Based on these data, it is acceptable that 20 million PE reads per sample for a diploid crustacean organism is a reasonable target to aim for, although there is no specific benchmark for all sequencing experiments.

It is also important to note that in order to detect transcripts with low expression, a deeper sequencing strategy may be needed. In the guidelines for the ENCODE project (<https://www.encodeproject.org/>),

an experiment to evaluate similarity between two transcriptional profiles, requires 30 million PE reads that must be mapped to the genome or known transcriptome. Guidelines to detect novel elements or quantification of known transcript isoforms requires deeper sequencing (Refer to the whole guideline at <https://www.encodeproject.org/about/experiment-guidelines/#guideline>). Another tool, Scotty, can be used to assist in the design phase of RNA-Seq experiments (Busby et al., 2013). This program can confirm if the design applied has sufficient statistical power to detect differentially expressed genes (DEGs) at the predetermined level required. The program is freely available online at <http://bioinformatics.bc.edu/marthlab/scotty/scotty.php/>.

An interim conclusion to be drawn from the above sections is that there exists a trade-off between depth of reads per sample and the number of samples (which include technical and biological repeats). The technology employed and financial limitations usually dictate a fine balance between these factors.

#### *Tissue RNA extraction and cDNA library preparation*

Library preparation is a crucial step prior to sequencing. It consists of a number of stages including RNA extraction, proper storage of RNA, quality checking of RNA, mRNA isolation and finally cDNA library generation.

In brief, extraction of total RNA from target tissue can be undertaken immediately on site or samples can be stored in RNA-later<sup>®</sup> solution for later extraction. It is important to note that RNA is extremely fragile and degrades readily if stored under inappropriate conditions. In addition, ribonucleases (RNases) which enzymatically degrade RNA pose a constant threat of contamination and degradation of purified RNA. Traditionally, RNA can be stored at -20, -80°C (most desirable) or in liquid nitrogen (-196°C) to provide protection. RNA storage solutions that include chelating agents which inhibit RNase activity can be used, although these might interfere with reverse transcription and should thus be removed prior to these steps. To our knowledge, there is no crustacean-specific RNA extraction kit available on the market; however, several commercial kits for RNA extraction are still usable for crustaceans, in addition to in-house (modified) versions of RNA extraction methods that use Beta-mercaptoethanol- or phenol-based



compounds, with the latter being more popular in recent publications. A detailed review on the effect of RNA extraction methods on RNA-seq can be found elsewhere (Sultan et al., 2014). RNA can then be assessed for quality and quantity using a Nanodrop® spectrophotometer or BioAnalyzer®. It is important to note that RNA integrity number (RIN) that has been used as a standardized metric of RNA quality for vertebrate species, is not usually valid for crustacean samples with non-typical RNA profiles. RIN is calculated based on the ratio between 18S ribosomal RNA (rRNA) and 28S rRNA band intensities, which are usually very conserved across eukaryotes. However, the 28S rRNA of arthropods tends to break down into two subunits, preventing a reliable RIN value calculation (Winnebeck et al., 2010; Macharia et al., 2015; McCarthy et al., 2015).

RNA can be stored and shipped in ambient conditions after desiccation with RNA-stable solution (Seelenfreund et al., 2014). An important consideration when it comes to RNA extraction in crustacean species is tissues with high pigment content (i.e. eyestalk). For these tissues, extra caution is suggested to avoid extracting pigment contamination that will affect the quality of library preparation. Currently, there is no threshold for deciding if a sample is too degraded for whole-transcriptome analysis. In most cases, however, sequencing facilities provide users with specific guidelines and technical notes recommended for producing the best results. Moreover, depending on each sequencing platform, different cDNA library preparation protocols may be required.

### Post-sequencing

Post-sequencing analyses include quality checking of raw sequences, trimming, de novo assembly of trimmed reads, read mapping and quantification, DEGs assessment and finally biological interpretation (Fig. 1).

#### *Quality control for SGS data*

Current SGS runs generate millions, or even hundreds of millions of read sequences. Technologies advancement reduces the error rate; however, every platform still produces read errors that require the application of a quality-control program post-sequencing. Read errors, while relatively negligible in

number compared with the massive dataset generated, still pose a hurdle for downstream analysis. For instance, errors in base-calling cause improper connection of nodes in de novo assembly (thus expanding running time and increase memory needed to store the nodes). In addition, incorrect SNP detection can result from an inability to differentiate between a true polymorphism and a sequencing error (Kelley et al., 2010). Several quality-control tools have been developed for NGS data (most popular tools are summarized in Table 1).

In general, quality control of raw reads from NGS sequencers can be completed in a few simple steps. Raw read statistics can then be checked with FASTQC software (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). A variety of parameters can be used to trim the data. The most important is the PHRED quality score [a base-calling score ranking system that allows users to judge the confidence of a nucleotide presumed to be correctly called (Ewing et al., 1998)]. Some other considerations include reads' average length, total number of base pairs and adapters' contamination. In addition, reads generated on the Illumina platform are considered to have a relatively higher error rate towards the 3'-end of the read (Schirmer et al., 2015), so if a drop in quality is detected, it is acceptable to trim off a portion of the read from that end. Some commonly used criteria for trimming reads include: minimum read length, minimum quality score, and homopolymer trimming. Read duplication is also a factor to consider during the quality-control step in NGS projects. In brief, read duplications are identical reads that map to the same genomic location (effects of PCR amplification bias, excess computational resources, and errors). Raw reads may also need to be cleaned from artificially introduced sequences—PCR primers or sequencing adapters; these are usually addressed in most quality-control packages. In a benchmarking study, it was shown that trimming applied in every sequencing project will improve not only quality of the results, but also reduce analysis duration (Del Fabbro et al., 2013). In general, normal quality trimming with a PHRED score ranging from 20 to 30 is normal for most RNA-seq experiments, while a PHRED score threshold of 30 or above is usually required for variant calling experiments (Ledergerber & Dessimoz, 2011). However, in one particular study, the authors highlighted that although strict trimming is usually applied, in

**Table 1** Commonly used quality-control software tools for NGS data

Software package	De-multiplexing	Adapter trimming	Quality Filtering/Trimming 3' and 5'	Kmer filtering	Citation
FASTX-Toolkit	+	+	+		<a href="http://hannonlab.cshl.edu/fastx_toolkit/">http://hannonlab.cshl.edu/fastx_toolkit/</a>
Trimmomatic	+	+	+		Bolger et al. (2014)
PRINSEQ		+	+		Schmieder & Edwards (2011)
FaQCs		+	+		Lo & Chain (2014)
Biopieces	+	+	+		<a href="http://maasha.github.io/biopieces/">http://maasha.github.io/biopieces/</a>
NGS QC Toolkit	+	+	+		Patel & Jain (2012)
CutAdapt		+	+		Martin (2011)
HTQC		+	+		Yang et al. (2013)
kPAL				+	Anvar et al. (2014)
khmer				+	<a href="https://github.com/dib-lab/khmer">https://github.com/dib-lab/khmer</a>
Stacks	+	+	+	+	Catchen et al. (2011)

+ Yes, – No

some cases, a more gentle trimming (PHRED score  $< 2$  or  $< 5$ ) might be more optimal (MacManes, 2014). This is due to the fact that short- and low-expressed transcripts suffer from heavy negative bias when using harsh trimming (MacManes, 2014). Therefore, lowering the PHRED score threshold in the quality-control step can result in a greater transcript discovery rate. In conclusion, we suggest gentle trimming initially as suggested in the above study. A list of some popular software packages for NGS quality control can be found in Table 1.

#### De novo assembly for non-model species and transcript clustering

For non-model decapod species, it is often difficult to align RNA-seq data to a reference genome from relatively recently diverged organisms (currently there are very few crustacean reference genomes available—see Sect. “Ongoing challenges for applied RNA-seq studies of crustaceans”). An alternative strategy therefore, is to construct a de novo assembly (new assembly) from high-quality reads. The primary aim is to extend the short reads from the sequencer into longer continuous sequences (contigs) that reflect the mRNAs transcribed in the cell without any chimeric/fusion events. A number of de novo transcriptome assemblers have been developed (initially

they were simply modified genome assemblers), including the Velvet/Oases pipeline (Zerbino & Birney, 2008; Schulz et al., 2012), SOAPdenovo (<https://soap.genomics.org.cn/soapdenovo.html>) and TransAbyss (Robertson et al., 2010). More recently, the Trinity software (Grabherr et al., 2011) has become available, developed specifically for de novo transcriptome assembly from short-read RNA-seq data. Since reads from SGS are short in length compared with pyrosequencing output (Liu et al., 2012), transcriptome de novo assemblers often employ a De Bruijn graph algorithm instead of the traditional Overlap Layout Consensus (OLC). This minimizes the amount of memory required to handle numerous parallel calculations. Further information on graph algorithms can be found elsewhere (Miller et al., 2010; Li et al., 2012).

Most de novo assemblers are freely distributed but usually required operating using command line, which deters many biologists without programming skills. To overcome this issue, bioinformatics platforms such as Galaxy (<https://www.usegalaxy.org/>) and CyVerse (<https://www.cyverse.org/>) embed command line packages into user-friendly interfaces. Yet, there is a limited flexibility in utilizing these tools. Learning how to use command line programming can be time consuming and potentially is out of reach for many non-model biology researchers and this can slow the



pace at which NGS studies are performed on these species. To address this, users can use commercial products (usually with a “point-and click” user-friendly interface) that are available on the market. A summary of some notable de novo assemblers can be found in Table 2. A performance comparison of commonly used de novo transcriptome assemblers can be found elsewhere (Surget-Groba & Montoya-Burgos, 2010; Zhao et al., 2011; Ghangal et al., 2013; Amin et al., 2014; Finseth & Harrison, 2014).

One significant challenge associated with de novo assembly is the lack of software to identify the assembly that is most accurate. To address this challenge, Sequence Comparative Analysis using Networks (SCAN) was created (Misner et al., 2013). SCAN uses a reference dataset (from a related genome) to identify the most accurate de novo assembly and to classify “good” transcripts in these assemblies (Misner et al., 2013). A similar program was generated for this purpose, named DETONATE (an abbreviation of *DE* novo TranscriptOme rNa-seq Assembly with or without the Truth Evaluation) (Li et al., 2014). This program combines multiple factors into a single evaluation score that then can be used to select the best assembler. The software is distributed freely at <https://deweylab.biostat.wisc.edu/detonate>. Another approach is to employ the CEGMA pipeline (Core Eukaryotic Genes Mapping Approach) (Parra et al., 2007) or BUSCO (Benchmarking Universal Single-Copy Orthologs) (Simão et al., 2015). These programs scan the de novo assembly against a dataset of core eukaryotic genes that are well conserved across several eukaryotic taxa, to calculate the coverage of protein-coding genes, thus estimating the degree of completeness of the reconstruction and the full-length complement of transcript sequences comprising the de novo transcriptome assembly. As a concluding remark, benchmarking assemblies are an option that can be trialled, but the practice is still in development.

### *Transcriptome mapping*

Following de novo assembly, reads can be aligned against the de novo assembly (mapping). The mapping step can serve two purposes: (i) a remapping step can be used to assess the assembly quality; and (ii) the alignment can then be quantified; gene-expression levels can be inferred from the total counts of reads

















aligned to each contig. Furthermore, mapping also enables variant calling for transcripts of interest.

Stringent parameters may result in a small subset of reads mapped, while less-stringent settings reduce mapped read specificity. To gain a balance between sensitivity and specificity, trials with different parameters can be performed. Popular aligners for RNA-seq include Bowtie 1 (Langmead et al., 2009) and Bowtie 2 (Langmead & Salzberg, 2012), BWA (Li & Durbin, 2009), GSnaps (Wu & Nacu, 2010), and commercial programs including CLC Genomics Workbench®, DNA-STAR® or Partek Genomics®. A detailed list of available aligners can be found at [https://www.ebi.ac.uk/~nf/hts\\_mappers/](https://www.ebi.ac.uk/~nf/hts_mappers/) (Fonseca et al., 2012). Comparisons of different aligners usually takes into consideration running time, accuracy, as well as the sensitivity of mapped reads (Li & Homer, 2010; Grant et al., 2011; Hatem et al., 2013; Baruzzo et al., 2017). Critically, for non-model organisms where no genome sequence is available, it is hard to define which are the best mapping parameters that can be applied. This is due to the occurrence of isoforms and splice variants that cannot be accurately determined without access to a reference genome. Reads can be mapped randomly to shared exons between splice variants, biasing the resulting count and confounding the biological interpretation.

### *Quantifying transcript level and analysis of differential gene expressions*

To quantify gene expression, RNA-seq reads need to be aligned to a reference genome from model organisms or to the transcriptome sequences reconstructed using de novo assembly strategies for organisms without reference genome sequences. The number of mapped reads is calculated based on the outcome of the alignment and can be used to estimate the relative expression level of individual genes. Following this, statistical methods are applied to test for significant differences among experimental groups. The data, however, first needs to be normalized since there are inherent differences in total reads per sample, resulting in over-represented long transcripts. With rapid development of RNA-seq technology, there are now numerous tools available to estimate gene-expression levels, which vary in their efficiency. Popular RNA-seq quantification (reads counting) tools include:

**Table 2** Summary of commonly used de novo transcriptome assemblers

Package	OS	Multiple k-mer support	Licence	Citation	Advantage	Disadvantage
Velvet/oases	  	Yes	F	Schulz et al. (2012), Zerbino & Birney (2008)	Widely used Well documented Stable Easy to run Multi kmer Fast (multithreading)	Currently inactive Need basic knowledge of Linux RAM intensive
Trans-ABYSS		Yes	F	Robertson et al. (2010)	Multi kmer Well documented Less RAM intensive	Currently inactive Slow
Trinity		Depend on version	F	Grabherr et al. (2011)	Widely used Well documented Well-established methodology Active community support Stable Relatively fast	Single kmer approach RAM extensive
SOAPdenovo- Trans		Yes	F	Xie et al. (2014)	Well documented Relatively fast	Currently inactive Relatively large amount of RAM Not widely used Not really well documented
Bridger		Yes	F	Chang et al. (2015)	Active community	Not widely used
CLC Genomics Workbench®	  	Yes	C	<a href="https://www.qiagenbioinformatics.com">https://www. qiagenbioinformatics. com</a>	Easy to use Point and click interface Genomics server option for large-scale analysis Relatively fast	Expensive licence
Geneious®	  	Yes	C	<a href="http://www.geneious.com">http://www.geneious.com</a>	Easy to use Point and click interface	Somewhat expensive
SeqMan NGen (DNASTAR® suite)	  	No	C	<a href="http://www.dnastar.com">http://www.dnastar.com</a>	Easy to use Point and click interface	Expensive licence Not widely used Not frequently updated

Licence C commercial product, F free. Platforms  Linux,  Windows,  MacOS

RSEM (Li & Dewey, 2011), eXpress (Roberts & Pachter, 2012), HTSeq (Anders et al., 2015), Salmon (Patro et al., 2017) and kallisto (Bray et al., 2016).

Several studies have also been conducted to compare the pros and cons of each tool (Li & Homer, 2010; Chandramohan et al., 2013; Teng et al., 2016).

DEG analysis programs perform statistical tests to determine if results of fold change under different experimental conditions are significant (e.g. among tissues types, life stages etc.). Many programs have been developed for DEG analysis (a brief summary of popular DEG tools can be found in Table 3) and several comparative assessments are available (Kvam et al., 2012; Robles et al., 2012; Soneson & Delorenzi, 2013; Zhang et al., 2014; Khang & Lau, 2015; Rajkumar et al., 2015). Much like assembly and mapping, there is no guarantee as to which tool is the best, or which parameters will result in the highest accuracy or robustness of the results generated (Zhang et al., 2014). Most DEG call methods are designed to address analysis of RNA-seq experiments that have biological replicates. There are a few tools, however, that can handle non-replicated experiments (e.g. GFOLD (Feng et al., 2012), EdgeR (Robinson et al., 2010), NOISeq (Tarazona et al., 2011, 2012). A recent study recommended using EdgeR (Robinson et al., 2010) or DESeq2 (Love et al., 2014) for experiments with less than 12 replicates per group, while they suggest studies with more than 12 replicates should use DESeq2 for the statistical analysis (Schurch et al., 2016). An alternative strategy is to employ several software packages and then compare the outcome of each approach, highlighting not only the similarity, but also differences among these analyses. Fold change is an important parameter to consider, but will depend on the number of reads that are assigned to a specific transcript. If the depth is low, yet with high fold change between groups, it should be considered as noise. For example,  $10 \times 100$  base reads mapped onto a 1 Kb transcript per sample in one group (giving an average depth of 1) compared to 1 read on average per sample in the other group is a tenfold change, yet the coverage is very low and should be validated using additional samples via qPCR.

### Annotation of transcripts

After all reads have been assembled de novo into contigs, the next step is to annotate all the contigs based on the most up-to-date database (i.e. identify homology to previously characterized genes). The most common way to annotate a large number of transcripts is the Basic Local Alignment Search Tool (BLAST). As the number of contigs in every de novo assembly can be thousands to a few hundred thousand sequences, usage of an automated search tool, in particular BLAST+ (Camacho et al., 2009), is essential. For non-model species, many candidate protein databases are available including the non-redundant protein database (nr), UniProt/Swiss-Prot database, and the Reference Sequence database (RefSeq). RefSeq (nucleotide and protein) and UniProt/Swiss-Prot (protein) consist of curated, well annotated sequences, whereas the nr database includes both curated and non-curated databases. For most crustacean RNA-seq experiments, the nr database is considered to be the best choice due to the fact that very few crustacean genes have been properly annotated to date, a problem that has been highlighted (Clark & Greenwood, 2016; Das & Mykles, 2016).

After transcripts have been scanned against the protein database and assigned annotations, there is a variety of downstream packages that can further analyse a contig, including Gene Ontology (GO) term analysis, functional enrichment analysis, protein domain analysis (PFAM domain search—pfam.xfam.org), and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway maps (Kanehisa & Goto, 2000). Each entry in the sequence database can be classified into a number of biologically relevant terms. In GO analysis, most genes can be assigned to one out of three basic ontologies: cellular component, biological process, or molecular function. When comparing

**Table 3** Popular read aligners for RNA-seq

Program	Algorithm	Long read	Gapped	Paired-end supported	Spliced	References
Bowtie	BWT	—	—	+	—	Ben (2010)
Bowtie2	BWT	+	+	+	—	Langmead & Salzberg (2012a)
BWA	BWT	+	+	+	—	Li & Durbin (2009a)
SOAP	HASH (reference)	—	+	+	—	Li et al. (2008)
GSNAP	HAST (read)	+	+	+	+	Wu & Nacu (2010a)

+ Yes, — No

samples from two groups, differentially represented GO terms can help define the mechanism via which the groups differ from one another. Similarly, when using KEGG, contigs can be assigned with components of specific pathways. Differential expression can allow a detailed assessment of the changes in pathways between studied groups. PFAM shows domains within the open reading frames of contigs that enables characterization of the protein function, based on the architecture of domains in a polypeptide chain. A number of software packages are capable of extracting vast numbers of GO terms from public databases including Blast2GO (Conesa et al., 2005), DAVID—<https://david.abcc.ncifcrf.gov/> (da Huang et al., 2009) or ermineJ (Lee et al., 2005). Among these programs, Blast2GO stands out as an easy-to-use, point-and-click program that has become very popular in the last few years (Table 4).

In addition to Blast2GO, databases like PFAM (Finn et al., 2016), eggNOG (Huerta-Cepas et al., 2015) and InterProScan (Jones et al., 2014) can be employed to predict the function of unknown proteins. The Trinity RNA-seq package, Trinotate (<https://trinotate.github.io/>), uses UniProt, eggNOG and GO Pathway databases for annotating novel sequences and these have been widely used in recent years (Das & Mykles, 2016; Das et al., 2016).

#### *Validation of RNA-seq results*

Validation is a very important step in every RNA-seq study. There is generally a very high correlation between RNA-seq and qPCR results with respect to relative gene expression. A significant point is that testing the same RNA samples used in the NGS platform for validation with techniques like qPCR or digital PCR only validates the sequencing accuracy

result. Therefore, additional, independent biological replicates should be included to properly validate the biological interpretation from the RNA-seq experiment. Essentially, validation post sequencing is now mandatory for publication. An approach has been proposed to set the minimum acceptable standard for qPCR validation (Fang & Cui, 2011) that takes a number of factors into consideration, including the number of genes tested and the number of isoform transcripts detected in the transcriptome. Nevertheless, up-to-date, qPCR techniques offer the easiest way to validate data in a transcriptomics study. One important note for researchers who are unfamiliar with the technology is that some RNA-seq pipelines allow RNA-seq analysis at the gene level [Trinity/RSEM for instance (Haas et al., 2013)]. However, there is a deeper level of transcripts component (in which transcripts can be isoforms resulting from alternative splicing events or a single nucleotide variation). Therefore, researchers should design primers that are not included in these regions to avoid unreliable qPCR results between biological replicates. As a concluding remark for this section, several studies have compared RNA-seq results to qPCR data, and have found excellent correlations between the approaches (Wu et al., 2014; Rajkumar et al., 2015; Everaert et al., 2017).

#### **Promises and challenges of RNA-seq-based studies in crustaceans**

##### *Studied topics*

SGS has revolutionized biological science, shifting it towards the post-genomic era. Transcriptomics studies in crustaceans include either

**Table 4** Widely used DEGs analysis tools

Software	Algorithm model	References
edgeR	Negative binomial (NB)	Robinson et al. (2010)
BBseq	NB	Zhou et al. (2011)
NOISeq	Nonparametric	Tarazona et al. (2011, 2012)
baySeq	NB	Hardcastle & Kelly (2010)
DESeq	NB	Anders & Huber (2012)
DESeq2	NB	Love et al. (2014)
DEGseq	MA-plot	Wang et al. (2010)
GFOLD	Nonparametric	Feng et al. (2012)

1. Sequencing and annotation of the transcriptome of one (or several) tissue/s, or a whole individual of a particular taxa in a specific developmental stage or under specific experimental conditions.
2. Applying RNA-seq to identify DEGs among different physiological conditions, treatments, developmental stages and/or tissues.
3. Identification of novel transcripts—enzymes, receptors, hormones, neuropeptides.
4. Screening for variant mutations—SNPs, SSRs and/or microsatellites.
5. A combination of the above.

To date, several RNA-seq projects have been initiated on a variety of crustacean species. In Table 5, we have summarized several RNA-seq-based studies on crustacean taxa that have been conducted over the last few years based on the following categories: *Aquatic toxicology*, *Reproduction & sexual differentiation*, *Disease resistance & immunology*, *Developmental biology* and *Physiology*. This is, however, by no means an exhaustive list as hundreds of applied RNA-seq studies have been undertaken in recent years, rather the list here illustrates several model comparative RNA-seq approaches.

Ongoing challenges for applied RNA-seq studies of crustaceans

#### *Experimental limitation*

A good experimental design will have a major impact on data outcomes; it can prevent wasted resources and help avoid the generation of unpublishable results. The balance between sequencing cost and experimental design constraints is a major issue that has been highlighted in many review articles. Due to budgetary limitations, there will always be an incentive to cut costs by sequencing with higher depth but with little or no biological replication. Furthermore, where depth is added, a large number of reads will be also mapped to the already well-covered regions, while if additional replication was available, greater statistical power can be achieved resulting in better biological inference. To resolve this problem, optimal guidelines for the design of RNA-seq experiments are needed and should be applied accordingly. In parallel, biological replicates (at least 3 or greater) are required for an RNA-seq study to reach a basic publishable level. Another

recommendation for best practice is undertaking a pilot-sequencing project where a high number of libraries are run on one lane initially. This can be valuable in assessing the feasibility of the larger experiment, as well as providing a good indicator for how to address trade-offs between obtaining high-quality output vs cost. Finally, although RNA-seq methods are becoming more robust and reliable and sometimes qPCR validations are proven to be unnecessary, a section for qPCR validation of selected genes/transcripts of interest may be beneficial to reveal the biological insights if the study has limited replications. Therefore, we recommend that for reliable biological interpretation and validation of RNA-seq analysis, the candidate genes themselves are tested for expression, rather than choosing random genes or genes showing high expression levels.

#### *In silico annotation and functional annotation*

Annotation of RNA-seq data is based loosely on BLAST searches. In fact, many BLAST results produce “hypothetical”, “predicted”, “uncharacterized”, or “low-quality” assignments. This highlights the fact that gene databases for non-model species currently, are very limited. To add another layer of complexity, *Daphnia pulex*, the model species currently available for crustaceans, has a large number of genes that currently remain unannotated. Furthermore, when compared with Decapoda, it is very remotely related and in many cases, shares higher similarity with insects than with other crustacean taxa. Further downstream annotation is also a constraint for crustacean RNA-seq studies, as specific GO classification and KEGG pathways are still not available for these taxa. As a result, drawing biological interpretations from predicted results can be problematic. Moreover, similarities in structure do not necessarily correlate with equivalent functionalities. It is crucial therefore, to highlight that in silico prediction is only speculative and functional annotation is very important to validate any biological interpretations (in particular for novel genes). RNAi technology (gene silencing) is now the go-to method for gene functional studies in decapod crustaceans and it has been already applied in some cases (Sagi et al., 2013). Gene editing technologies, for example CRISPR/Cas9 technology, have emerged recently and hold great potential for functional annotation in decapod crustacean species (Mykles &

**Table 5** Summary of recent RNA-seq studies in non-model decapod crustacean species

Application	Species	Key findings	References
Aquatic toxicology	<i>P. trituberculatus</i>	69 Differentially expressed genes (DEGs) were detected under ammonia stress	Ren & Pan (2014)
	<i>S. henanense</i>	Genes involved in macromolecular metabolism, oxidative phosphorylation, detoxification and anti-oxidant were upregulated significantly by Cadmium exposure	Sun et al. (2016)
	<i>P. vannamei</i>	Comparative RNA-seq analysis in hepatopancreas identified 136 significantly DEGs under acute ammonia stress	Lu et al. (2016)
	<i>P. elegans</i>	Seven genes out of 35 selected as potential targets for Teflubenzuron exposure showed significant differential expression. Markers linked to moulting and exoskeleton change (CHH, CTBS, GAP65), stress and apoptosis (HSP40, HSP70, CASP3), and detoxification (CYP6A18) were highlighted as important during the challenge	Olsvik et al. (2017)
	<i>P. australiensis</i>	Insights into stress response pathways induced in crustaceans by short-term exposure to multiple stressors present in acid drainage water (ADW). RNA-seq analysis revealed that functions associated with cuticle biosynthesis and oxidative stress were significantly enriched in the lists of transcripts after exposure to 50 or 100% ADW	Bain et al. (2016)
	<i>P. trituberculatus</i>	1572 DEGs were detected under the effect of aerial exposure, this study highlighted dramatic tissue oxidation stress and the alteration of the tissue epithelial structure	(Li et al. 2017)
Reproduction and sexual differentiation	<i>S. paramamosain</i>	RNA-seq analysis relating to sex differentiation revealed 4021 DEGs, 10,522 ovary-specific, and 19,013 testis-specific transcripts	Gao et al. (2014)
	<i>E. sinensis</i>	Insights into sexual differentiation resulted in 448 DEGs, 29 candidate genes involved in primary sex determination pathways were detected	Liu et al. (2015)
	<i>P. vannamei</i>	Transcriptome comparison between gonads of <i>P. vannamei</i> resulted in 19,279 testicular upregulated and 3529 ovarian upregulated transcripts	Peng et al. (2015)
	<i>S. olivacea</i>	Transcriptome comparison among the testes of different maturation stages revealed one gene (beta-crystallin-like gene) with the most significant differential expression between stages	Waiho et al. (2017)
	<i>P. trituberculatus</i>	RNA-seq analysis generated 5919 DEGs between ovary and testis, among which many genes related to gametogenesis and several genes previously reported to be critical in differentiation and development of gonads were confirmed	Meng et al. (2015)
	<i>M. nipponense</i>	DEGs enrichment analysis results revealed several important reproduction related terms and signalling pathways in an RNA-seq analysis between eyestalk and cerebral ganglia of female <i>M. nipponense</i> during breeding and non-breeding seasons	Qiao et al. (2017)
	<i>S. verreauxi</i>	An RNA-seq-based study to enhance understanding of key regulatory pathways of sexual development. This study highlighted several key regulatory factors that are the Dsx- and mab-3-related transcription factors (DMRUTS)	Chandler et al. (2016)
	<i>N. norvegicus</i>	Sex-specific transcripts were detected primarily in gonads followed by hepatopancreas, brain, thoracic ganglia, and eyestalk. Sex-specific markers were mainly expressed in males implying that males may experience stronger selection than females	Rotllant et al. (2017)



**Table 5** continued

Application	Species	Key findings	References
Disease resistance/ immunology	<i>E. sinensis</i>	Microbial challenged <i>Eriocheir sinensis</i> revealed numerous genes were further identified to be associated with multiple immune pathways	Cui et al. (2013)
	<i>P. vannamei</i>	1374 significant DEGs were detected under the effect of Taura syndrome virus (TSV)	Sookruksawong et al. (2013)
	<i>F. merguensis</i>	Effect of hepatopancreatic parvo-like virus (HPV) were revealed by using hepatopancreas tissues from six full-sib groups of banana shrimp. 404 DEGs were identified with 180 being over-expressed and 224 under-expressed	Powell et al. (2016)
	<i>M. rosenbergii</i>	14,569 Transcripts were differentially expressed in hepatopancreas of <i>M. rosenbergii</i> in response to <i>Vibrio parahaemolyticus</i> . Most differentially expressed immune genes are classified under various processes of the animal immune system	Rao et al. (2015)
	<i>P. clarkii</i>	Insights into crayfish antiviral immunity mechanisms against white spot syndrome virus (WSSV), over 7000 DEGs were significantly regulated and 5 key DEGs involved in the JAK-STAT signalling pathway were validated	Du et al. (2016)
	<i>E. carinicauda</i>	An RNA-seq-based study on shrimp immune system against AHPND (Acute hepatopancreatic necrosis disease) infection, which include genes and pathways associated with AHPND pathogenesis and immune defences. A total of 229 DEGs that have high homologies with the known proteins in crustacean species were identified, among which 127 genes are reported potentially related to immune function.	Ge et al. (2017)
	<i>M. rosenbergii</i>	Evaluation WSSV infection on <i>M. rosenbergii</i> . This study recorded 4951 DEGs that were identified and categorized into 244 metabolic pathways to improve understanding of the effect of this deadly disease	Cao et al. (2017)
	<i>E. sinensis</i>	Multiple immune-related genes (GNPTAB, MASP2, F7, F5, NFATC, TRAF6, MAP3K5 and TRA) were identified as DEGs and confirmed using RT-qPCR in <i>E. sinensis</i> under the effect of <i>Spiroplasma eriocheiris</i>	Wang et al. (2017)
	<i>S. paramamosain</i>	Insights of crab infected with Mud crab reovirus (MCRV). 13,856 DEGs, including 4444 significantly upregulated and 9412 downregulated transcripts, were detected in diseased crabs compared with the control	Liu et al. (2017)
Developmental biology	<i>P. vannamei</i>	Multiple genes related to exoskeleton development were identified as DEGs in the early developmental stages and adult-moulting stage	Gao et al. (2017)
	<i>P. trituberculatus</i>	1394 moult-related DEGs were identified among intermoult, premoult and postmoult	Lv et al. (2017)
	<i>M. rosenbergii</i>	Analysis of differential gene expressions among various stages of the species highlighted various candidate genes elucidating metamorphosis process between larvae and post-larvae	Ventura et al. (2013)
	<i>S. verreauxi</i>	Reveals key players during the Phyllosoma–Peurulus–Juvenile metamorphosis in the Eastern Rock Lobster	Ventura et al. (2015)
	<i>E. sinensis</i>	The study revealed several DEGs between the two stages (zoeae to megalopae), mostly involving morphological change, sensory ability, neuroendocrine system and food digestion	Li et al. (2015)

**Table 5** continued

Application	Species	Key findings	References
	<i>H. rubra</i>	A comparison RNA-seq profiles between “early” (i.e. Z1–Z2) as well as “late” (i.e. Z3–Z4) larval and adult developmental stages of the species. Given multiple differences between the stages, a large number of DEGs were identified to be upregulated, including opsins and light/stimulus detection genes; other DEGs were found to be downregulated including ion transport, digestion and reproduction	Havird & Santos (2016b)
	<i>M. olfersi</i>	35 homologues of embryonic development toolkit genes were identified. Also, developmental pathways-related genes were identified including TGF- $\beta$ , Wnt, Notch, MAPK, Hedgehog, JAK-STAT, VEGF, and ecdysteroid-inducible nuclear receptors	Jaramillo et al. (2016)
	<i>E. sinensis</i>	RNA-seq profiles, between oosperms and embryos at the 2–4 cell stage (Cs) that are separated by a cleavage event. were characterized. 432 DEGs were detected	Hui et al. (2017)
	<i>P. vannamei</i>	Multiple gene families were identified to be differentially expressed under five different development stages (embryo, nauplius, zoea, mysis and post-larvae)	Wei et al. (2014)
	<i>M. rosenbergii</i>	40 potentially candidate genes in determining adult male morphotypes (Small male, Orange claw and Blue claw) were curated.	Aziz et al. (2017)
Circadian rhythm	<i>N. norvegicus</i>	A curated list of candidate clock genes timeless, period, clock and bmal1	Sbragaglia et al. (2015)
Osmoregulation	<i>L. vannamei</i>	Multiple immune signalling pathways are highlighted, which are involved in salinity adaptation of the species	Chen et al. (2015), Zhao et al. (2015), Zhang et al. (2016),
	<i>M. rosenbergii</i>	Multiple genes of interest are differentially expressed in two stages: larval and post-larval under the effect of salinity changes. GO enrichment resulted in multiple GO that are over representative including transcriptional regulation, signal transduction, immune response and ion-binding	Chakrapani et al. (2016)
	<i>Macrobrachium</i> sp.	Group of genes that are involved in the evolutionary adaptation of <i>Macrobrachium</i> spp	Moshtaghi et al. (2016), Rahi et al. (2017), Moshtaghi et al. (2017)
	<i>E. sinensis</i>	This study recorded 1151 and 941 genes, which are significantly differentially expressed in crab gill and muscle, respectively. Also results indicated top pathways differentially expressed following osmotic challenge in gill and muscle	Li et al. (2014)
	<i>P. trituberculatus</i>	Comparative genomic analysis revealed 1705 DEGs in salinity stress compared to the controls; multiple gene families that relate to osmoregulation are highlighted	Lv et al. (2013)
	<i>C. sapidus</i>	Comparative RNA-seq between anterior (non-osmoregulatory) and posterior (osmoregulatory) gills revealed 477 DEGs related to a wide variety of physiological processes. Notably, anterior gills had more upregulated structural genes such as actin and cuticle proteins, while ion transport and energy-related genes were upregulated in posterior gills	Havird et al. (2016)

Hui, 2015). Employing RNAi and/or CRISPR-Cas9 in RNA-seq studies would be extremely helpful to highlight key genes and resolve functional roles of novel genes for crustacean species.

#### *Combining transcriptomics/RNA-seq with other OMICS techniques*

In parallel with advances in RNA-seq technologies, other OMICS technologies including genomics, proteomics, metagenomics phylogenomics and phenomics have also developed rapidly. This highlights a challenge for RNA-seq studies, to make use of other OMICS approaches and to utilize them to create a multilayer outcome. One key reason why decapod crustacean genomes are not yet available is that they are often very large and complex which makes them hard to resolve. Nevertheless, draft genomes of a few crustacean species have been made publicly available recently including draft genomes for some decapods including: *N. denticulata* (Kenny et al., 2014), *P. vannamei* (Yu et al., 2015), *E. sinensis* (Song et al., 2016), *P. hawaiiensis* (Kao et al., 2016), *P. monodon* and *M. japonicus* (Yuan et al., 2017), and *P. virginalis* (Gutekunst et al., 2018). Utilizing these new genomic resources will allow better gene annotation and functional annotation of crustacean gene pathways. There is no doubt that in the near future, when the cost barrier for sequencing is essentially overcome, coupled with improved sequencing technologies, combining RNA-seq approaches with integrated OMICS will enable researchers to answer the most complex of biological questions.

#### Conclusions

To conclude, RNA-seq offers great promise for crustacean studies. It is a very powerful tool that can lead to developing a better understanding of underlying pathways and mechanisms, which form the basis of many scientific questions. The guidelines offered here for future RNA-seq studies of crustaceans are an attempt to assist biologists who are not familiar with the complex and diverse array of bioinformatics software that are currently available. It is also important, however, to highlight the gap between in silico prediction from RNA-seq analysis and in vivo results. This may be explained in general, by

limitations on experimental designs in the past, the lack of annotation databases for crustacean species, as well as the need for question-driven research. In the future, we also suggest that RNA-seq should be integrated with other OMICS technologies to increase data output as well as improving biological insights.

**Acknowledgements** The current study was supported through a Marie Curie International Research Staff Exchange Scheme Fellowship within the 7th European Community Framework Programme (612296-DeNuGReC). Tuan Viet Nguyen was supported through the Australian Research Council Discovery Project grant awarded to Dr Tomer Ventura (No. DP160103320) and a USC International PhD scholarship. The authors would like to acknowledge the precious help of Professor Abigail Elizur (University of the Sunshine Coast, Australia) and four anonymous reviewers for numerous feedbacks that helped improve the quality of this manuscript.

#### References

- Amin, S., P. Prentis, E. Gilding & A. Pavasovic, 2014. Assembly and annotation of a non-model gastropod (*Nerita melanotragus*) transcriptome: a comparison of De novo assemblers. *BMC Research Notes* 7(1): 488.
- Anders, S. & W. Huber, 2012. Differential expression of RNA-Seq data at the gene level—the DESeq package. *European Molecular Biology Laboratory, Heidelberg*.
- Anders, S., P. T. Pyl & W. Huber, 2015. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics* 31(2): 166–169.
- Anvar, S., L. Khachatryan, M. Vermaat, M. van Galen, I. Pulyakhina, Y. Ariyurek, K. Kraaijeveld, J. den Dunnen & P. de Knijff, 2014. Determining the quality and complexity of next-generation sequencing data without a reference genome. *Genome Biology* 15(12): 555.
- Auer, P. L. & R. W. Doerge, 2010. Statistical design and analysis of RNA sequencing data. *Genetics* 185(2): 405–416.
- Aziz, D., V. T. Nguyen, M. L. Rahi, D. A. Hurwood & P. B. Mather, 2017. Identification of genes that potentially affect social dominance hierarchy in adult male giant freshwater prawns (*Macrobrachium rosenbergii*). *Aquaculture* 476: 168–184.
- Bain, P. A., A. L. Gregg & A. Kumar, 2016. *De novo* assembly and analysis of changes in the protein-coding transcriptome of the freshwater shrimp *Paratya australiensis* (Decapoda: Atyidae) in response to acid sulfate drainage water. *BMC Genomics* 17(1): 890.
- Baruzzo, G., K. E. Hayer, E. J. Kim, B. Di Camillo, G. A. FitzGerald & G. R. Grant, 2017. Simulation-based comprehensive benchmarking of RNA-seq aligners. *Nature Methods* 14(2): 135–139.
- Ben, L., 2010. Aligning Short Sequencing Reads with Bowtie. *Current Protocols in Bioinformatics*. <https://doi.org/10.1002/0471250953.bi1107s32>.

- Bolger, A. M., M. Lohse & B. Usadel, 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bray, N. L., H. Pimentel, P. Melsted & L. Pachter, 2016. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology* 34(5): 525–527.
- Busby, M. A., C. Stewart, C. A. Miller, K. R. Grzeda & G. T. Marth, 2013. Scotty: a web tool for designing RNA-Seq experiments to measure differential gene expression. *Bioinformatics* 29(5): 656–657.
- Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer & T. L. Madden, 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10: 421.
- Cao, J., L. Wu, M. Jin, T. Li, K. Hui & Q. Ren, 2017. Transcriptome profiling of the *Macrobrachium rosenbergii* lymphoid organ under the white spot syndrome virus challenge. *Fish & Shellfish Immunology* 67: 27–39.
- Cartolano, M., B. Huettel, B. Hartwig, R. Reinhardt & K. Schneeberger, 2016. cDNA library enrichment of full length transcripts for SMRT long read sequencing. *PLoS ONE* 11(6): e0157779.
- Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko & J. H. Postlethwait, 2011. Stacks: building and genotyping loci *de novo* from short-read sequences. *G3: Genes, Genomes, Genetics* 1(3): 171–182.
- Chakrapani, V., S. K. Patra, S. D. Mohapatra, K. D. Rasal, U. Deshpande, S. Nayak, J. K. Sundaray, P. Jayasankar & H. K. Barman, 2016. Comparative transcriptomic profiling of larvae and post-larvae of *Macrobrachium rosenbergii* in response to metamorphosis and salinity exposure. *Genes & Genomics* 38(11): 1061–1076.
- Chandler, J. C., J. Aizen, Q. P. Fitzgibbon, A. Elizur & T. Ventura, 2016. Applying the power of transcriptomics: understanding male sexual development in Decapod Crustacea. *Integrative and Comparative Biology* 56(6): 1144–1156.
- Chandramohan, R., P. Y. Wu, J. H. Phan & M. D. Wang, 2013. Systematic assessment of RNA-Seq quantification tools using simulated sequence data. *ACM Conference on Bioinformatics, Computational Biology and Biomedicine*. <https://doi.org/10.1145/2506583.2506648>.
- Chang, Z., G. Li, J. Liu, Y. Zhang, C. Ashby, D. Liu, C. Cramer & X. Huang, 2015. Bridger: a new framework for *de novo* transcriptome assembly using RNA-seq data. *Genome Biology* 16(1): 30.
- Chen, S.-Y., F. Deng, X. Jia, C. Li & S.-J. Lai, 2017. A transcriptome atlas of rabbit revealed by PacBio single-molecule long-read sequencing. *Scientific Reports* 7(1): 7648.
- Chen, K., E. Li, T. Li, C. Xu, X. Wang, H. Lin, J. G. Qin & L. Chen, 2015. Transcriptome and molecular pathway analysis of the hepatopancreas in the Pacific White Shrimp *Litopenaeus vannamei* under chronic low-salinity stress. *PLoS ONE* 10(7): e0131503.
- Clark, K. F. & S. J. Greenwood, 2016. Next-Generation Sequencing and the crustacean immune system: the need for alternatives in immune gene annotation. *Integrative and Comparative Biology* 56(6): 1113–1130.
- Conesa, A., S. Gotz, J. M. Garcia-Gomez, J. Terol, M. Talon & M. Robles, 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21(18): 3674–3676.
- Conesa, A., P. Madrigal, S. Tarazona, D. Gomez-Cabrero, A. Cervera, A. McPherson, M. W. Szczesniak, D. J. Gaffney, L. L. Elo, X. Zhang & A. Mortazavi, 2016. A survey of best practices for RNA-seq data analysis. *Genome Biology* 17(1): 13.
- Cui, Z., X. Li, Y. Liu, C. Song, M. Hui, G. Shi, D. Luo & Y. Li, 2013. Transcriptome profiling analysis on whole bodies of microbial challenged *Eriocheir sinensis* larvae for immune gene identification and SNP development. *PLoS ONE* 8(12): e82156.
- Das, S. & D. L. Mykles, 2016. A comparison of resources for the annotation of a *de novo* assembled transcriptome in the molting gland (Y-Organ) of the Blackback Land Crab, *Gecarcinus lateralis*. *Integrative and Comparative Biology* 56(6): 1103–1112.
- Das, S., S. Shyamal & D. S. Durica, 2016. Analysis of annotation and differential expression methods used in RNA-seq Studies in crustacean systems. *Integrative and Comparative Biology* 56(6): 1067–1079.
- Dillies, M.-A., A. Rau, J. Aubert, C. Hennequet-Antier, M. Jeanmougin, N. Servant, C. Keime, G. Marot, D. Castel, J. Estelle, G. Guernec, B. Jagla, L. Jouneau, D. Laloë, C. Le Gall, B. Schaeffer, S. Le Crom, M. Guedj & F. Jaffrézic, 2013. A comprehensive evaluation of normalization methods for Illumina high-throughput RNA sequencing data analysis. *Briefings in Bioinformatics* 14(6): 671–683.
- Du, Z., Y. Jin & D. Ren, 2016. In-depth comparative transcriptome analysis of intestines of red swamp crayfish, *Procambarus clarkii*, infected with WSSV. *Scientific reports* 6: 26780.
- Everaert, C., M. Luybaert, J. L. V. Maag, Q. X. Cheng, M. E. Dinger, J. Hellemans & P. Mestdagh, 2017. Benchmarking of RNA-sequencing analysis workflows using whole-transcriptome RT-qPCR expression data. *Scientific Reports* 7: 1559.
- Ewing, B., L. Hillier, M. C. Wendl & P. Green, 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Research* 8(3): 175–185.
- Del Fabbro, C., S. Scalabrin, M. Morgante & F. M. Giorgi, 2013. An extensive evaluation of read trimming effects on Illumina NGS data analysis. *PLoS ONE* 8(12): e85024.
- Fang, Z. & X. Cui, 2011. Design and validation issues in RNA-seq experiments. *Briefings in Bioinformatics*. <https://doi.org/10.1093/bib/bbr004>.
- Feng, J., C. A. Meyer, Q. Wang, J. S. Liu, X. Shirley Liu & Y. Zhang, 2012. GFOLD: a generalized fold change for ranking differentially expressed genes from RNA-seq data. *Bioinformatics* 28(21): 2782–2788.
- Finn, R. D., P. Coghill, R. Y. Eberhardt, S. R. Eddy, J. Mistry, A. L. Mitchell, S. C. Potter, M. Punta, M. Qureshi, A. Sangrador-Vegas, G. A. Salazar, J. Tate & A. Bateman, 2016. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Research* 44(D1): D279–D285.
- Finseth, F. R. & R. G. Harrison, 2014. A comparison of Next Generation Sequencing technologies for transcriptome assembly and utility for RNA-Seq in a non-model bird. *PLoS ONE* 9(10): e108550.

- Fonseca, N. A., J. Rung, A. Brazma & J. C. Marioni, 2012. Tools for mapping high-throughput sequencing data. *Bioinformatics*. <https://doi.org/10.1093/bioinformatics/bts605>.
- Francis, W. R., L. M. Christianson, R. Kiko, M. L. Powers, N. C. Shaner & S. H. Haddock, 2013. A comparison across non-model animals suggests an optimal sequencing depth for de novo transcriptome assembly. *BMC Genomics* 14(1): 167.
- Gao, J., X. Wang, Z. Zou, X. Jia, Y. Wang & Z. Zhang, 2014. Transcriptome analysis of the differences in gene expression between testis and ovary in green mud crab (*Scylla paramamosain*). *BMC Genomics* 15: 585.
- Gao, Y., J. Wei, J. Yuan, X. Zhang, F. Li & J. Xiang, 2017. Transcriptome analysis on the exoskeleton formation in early developmental stages and reconstruction scenario in growth-moulting in *Litopenaeus vannamei*. *Scientific reports* 7(1): 1098.
- Ge, Q., J. Li, J. Wang, J. Li, H. Ge & Q. Zhai, 2017. Transcriptome analysis of the hepatopancreas in *Exopalaemon carinicauda* infected with an AHPND-causing strain of *Vibrio parahaemolyticus*. *Fish & Shellfish Immunology* 67: 620–633.
- Ghangal, R., S. Chaudhary, M. Jain, R. S. Purty & P. Chand Sharma, 2013. Optimization of *De Novo* short read assembly of Seabuckthorn *Hippophae rhamnoides* L. transcriptome. *PLoS ONE* 8(8): e72516.
- Glenn, T. C., 2011. Field guide to next-generation DNA sequencers. *Molecular Ecology Resources* 11(5): 759–769.
- Goodwin, S., J. D. McPherson & W. R. McCombie, 2016. Coming of age: ten years of Next Generation Sequencing technologies. *Nature Reviews Genetics* 17(6): 333–351.
- Grabherr, M., B. Haas, M. Yassour, J. Levin, D. Thompson, I. Amit, X. Adiconis, L. Fan, R. Raychowdhury, Q. Zeng, Z. Chen, E. Mauceli, N. Hacohen, A. Gnirke, N. Rhind, F. di Palma, B. Birren, C. Nusbaum, K. Lindblad-Toh, N. Friedman & A. Regev, 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29(7): 644–652.
- Grant, G. R., M. H. Farkas, A. D. Pizarro, N. F. Lahens, J. Schug, B. P. Brunk, C. J. Stoeckert, J. B. Hogenesch & E. A. Pierce, 2011. Comparative analysis of RNA-Seq alignment algorithms and the RNA-Seq unified mapper (RUM). *Bioinformatics* 27(18): 2518–2528.
- Gutekunst, J., R. Andriantsoa, C. Falckenhayn, K. Hanna, W. Stein, J. Rasamy & F. Lyko, 2018. Clonal genome evolution and rapid invasive spread of the marbled crayfish. *Nature Ecology & Evolution* 2(3): 567–573.
- Haas, B. J., A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B. Couger, D. Eccles, B. Li, M. Lieber, M. D. MacManes, M. Ott, J. Orvis, N. Pochet, F. Strozzi, N. Weeks, R. Westerman, T. William, C. N. Dewey, R. Henschel, R. D. LeDuc, N. Friedman & A. Regev, 2013. De novo transcript sequence reconstruction from RNA-Seq: reference generation and analysis with Trinity. *Nature Protocols*. <https://doi.org/10.1038/nprot.2013.084>.
- Hardcastle, T. J. & K. A. Kelly, 2010. BaySeq: empirical Bayesian methods for identifying differential expressions in sequence count data. *BMC bioinformatics* 11(1): 422.
- Hatem, A., D. Bozdağ, A. E. Toland & Ü. V. Çatalyürek, 2013. Benchmarking short sequence mapping tools. *BMC Bioinformatics* 14(1): 184.
- Havird, J. C., R. T. Mitchell, R. P. Henry & S. R. Santos, 2016. Salinity-induced changes in gene expression from anterior and posterior gills of *Callinectes sapidus* (Crustacea: Portunidae) with implications for crustacean ecological genomics. *Comparative biochemistry and physiology Part D, Genomics & proteomics* 19: 34–44.
- Havird, J. C. & S. R. Santos, 2016a. Here we are, but where do we go? A systematic review of crustacean transcriptomic studies from 2014–2015. *Integrative and Comparative Biology* 56(6): 1055–1066.
- Havird, J. C. & S. R. Santos, 2016b. Developmental Transcriptomics of the Hawaiian Anchialine Shrimp *Holocaridina rubra* Holthuis, 1963 (Crustacea: Atyidae). *Integr Comp Biol* 56(6): 1170–1182.
- da Huang, W., B. T. Sherman & R. A. Lempicki, 2009. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols* 4(1): 44–57.
- Huerta-Cepas, J., D. Szklarczyk, K. Forslund, H. Cook, D. Heller, M. C. Walter, T. Rattei, D. R. Mende, S. Sunagawa & M. Kuhn, 2015. eggNOG 4.5: a hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. *Nucleic Acids Research* 44(D1): D286–D293.
- Hui, M., Z. Cui, Y. Liu & C. Song, 2017. Transcriptome profiles of embryos before and after cleavage in *Eriocheir sinensis*: identification of developmental genes at the earliest stages. *Chinese Journal of Oceanology and Limnology* 35(4): 770–781.
- Jaramillo, M. L., F. Guzman, C. L. Paese, R. Margis, E. M. Nazari, D. Ammar & Y. M. R. Müller, 2016. Exploring developmental gene toolkit and associated pathways in a potential new model crustacean using transcriptomic analysis. *Development Genes and Evolution* 226(5): 325–337.
- Jin, S., H. Fu, Q. Zhou, S. Sun, S. Jiang, Y. Xiong, Y. Gong, H. Qiao & W. Zhang, 2013. Transcriptome analysis of androgenic gland for discovery of novel genes from the oriental river prawn, *Macrobrachium nipponense*, using Illumina Hiseq 2000. *PLoS one* 8(10): e76840.
- Jones, P., D. Binns, H.-Y. Chang, M. Fraser, W. Li, C. McAnulla, H. McWilliam, J. Maslen, A. Mitchell, G. Nuka, S. Pesseat, A. F. Quinn, A. Sangrador-Vegas, M. Scheremetjew, S.-Y. Yong, R. Lopez & S. Hunter, 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* 30(9): 1236–1240.
- Jung, H., R. E. Lyons, H. Dinh, D. A. Hurwood, S. McWilliam & P. B. Mather, 2011. Transcriptomics of a Giant Freshwater Prawn (*Macrobrachium rosenbergii*): de novo assembly, annotation and marker discovery. *PLoS ONE* 6(12): e27938.
- Jung, H., B.-H. Yoon, W.-J. Kim, D.-W. Kim, D. Hurwood, R. Lyons, K. Salin, H.-S. Kim, I. Baek, V. Chand & P. Mather, 2016. Optimizing hybrid *de novo* transcriptome assembly and extending genomic resources for Giant Freshwater Prawns (*Macrobrachium rosenbergii*): the identification of genes and markers associated with reproduction. *International Journal of Molecular Sciences* 17(5): 690.



- Kanehisa, M. & S. Goto, 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research* 28(1): 27–30.
- Kao, D., A. G. Lai, E. Stamatakis, S. Rosic, N. Konstantinides, E. Jarvis, A. Di Donfrancesco, N. Pouchkina-Stancheva, M. Sémon, M. Grillo, H. Bruce, S. Kumar, I. Siwanowicz, A. Le, A. Lemire, M. B. Eisen, C. Extavour, W. E. Browne, C. Wolff, M. Averof, N. H. Patel, P. Sarkies, A. Pavlopoulos & A. Aboobaker, 2016. The genome of the crustacean *Parhyale hawaiiensis*, a model for animal development, regeneration, immunity and lignocellulose digestion. *eLife* 5: e20062.
- Kelley, D. R., M. C. Schatz & S. L. Salzberg, 2010. Quake: quality-aware detection and correction of sequencing errors. *Genome Biology* 11(11): R116.
- Kenny, N. J., Y. W. Sin, X. Shen, Q. Zhe, W. Wang, T. F. Chan, S. S. Tobe, S. M. Shimeld, K. H. Chu & J. H. Hui, 2014. Genomic sequence and experimental tractability of a new decapod shrimp model. *Neocaridina denticulata*. *Marine Drugs* 12(3): 1419–1437.
- Khang, T. F. & C. Y. Lau, 2015. Getting the most out of RNA-seq data analysis. *PeerJ* 3: e1360.
- Koboldt, Daniel C., Karyn M. Steinberg, David E. Larson, Richard K. Wilson & E. R. Mardis, 2013. The next-generation sequencing revolution and its impact on genomics. *Cell* 155(1): 27–38.
- Kuo, R. I., E. Tseng, L. Eory, I. R. Paton, A. L. Archibald & D. W. Burt, 2017. Normalized long read RNA sequencing in chicken reveals transcriptome complexity similar to human. *BMC Genomics* 18(1): 323.
- Kvam, V. M., P. Liu & Y. Si, 2012. A comparison of statistical methods for detecting differentially expressed genes from RNA-seq data. *American Journal of Botany* 99(2): 248–256.
- Lahens, N. F., E. Ricciotti, O. Smirnova, E. Toorens, E. J. Kim, G. Baruzzo, K. E. Hayer, T. Ganguly, J. Schug & G. R. Grant, 2017. A comparison of Illumina and Ion Torrent sequencing platforms in the context of differential gene expression. *BMC Genomics* 18(1): 602.
- Lam, H. Y. K., M. J. Clark, R. Chen, R. Chen, G. Natsoulis, M. O'Huallachain, F. E. Dewey & L. Habegger, 2012. Performance comparison of whole-genome sequencing platforms. *Nature Biotechnology*. <https://doi.org/10.1038/nbt.2065>.
- Langmead, B. & S. L. Salzberg, 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 9(4): 357–359.
- Langmead, B., C. Trapnell, M. Pop & S. L. Salzberg, 2009. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biology* 10(3): R25.
- Ledergerber, C. & C. Dessimoz, 2011. Base-calling for next-generation sequencing platforms. *Briefings in Bioinformatics* 12(5): 489–497.
- Lee, H. K., W. Braynen, K. Keshav & P. Pavlidis, 2005. ErmineJ: tool for functional analysis of gene expression data sets. *BMC Bioinformatics* 6(1): 269.
- Levin, J. Z., M. Yassour, X. Adiconis, C. Nusbaum, D. A. Thompson, N. Friedman, A. Gnirke & A. Regev, 2010. Comprehensive comparative analysis of strand-specific RNA sequencing methods. *Nature Methods* 7(9): 709–715.
- Li, Z., Y. Chen, D. Mu, J. Yuan, Y. Shi, H. Zhang, J. Gan, N. Li, X. Hu, B. Liu, B. Yang & W. Fan, 2012. Comparison of the two major classes of assembly algorithms: overlap-layout-consensus and de-Brujin-graph. *Briefings in Functional Genomics* 11(1): 25–37.
- Li, B. & C. N. Dewey, 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* 12: 323.
- Li, H. & R. Durbin, 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25(14): 1754–1760.
- Li, H. & N. Homer, 2010. A survey of sequence alignment algorithms for next-generation sequencing. *Briefings in Bioinformatics* 11(5): 473–483.
- Li, B., N. Fillmore, Y. Bai, M. Collins, J. A. Thomson, R. Stewart & C. N. Dewey, 2014. Evaluation of de novo transcriptome assemblies from RNA-Seq data. *Genome Biology* 15(12): 553. <https://doi.org/10.1186/s13059-014-0553-5>.
- Li, Y., M. Hui, Z. Cui, Y. Liu, C. Song & G. Shi, 2015. Comparative transcriptomic analysis provides insights into the molecular basis of the metamorphosis and nutrition metabolism change from zoeae to megalopae in *Eriocheir sinensis*. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics* 13: 1–9.
- Li, Y., S. Lai, R. Wang, Y. Zhao, H. Qin, L. Jiang, N. Li, Q. Fu & C. Li, 2017. RNA-Seq analysis of the antioxidant status and immune response of *Portunus trituberculatus* following aerial exposure. *Marine Biotechnology* 19(1): 89–101.
- Li, R., Y. Li, K. Kristiansen & J. Wang, 2008. SOAP: short oligonucleotide alignment program. *Bioinformatics* 24(5): 713–714.
- Li, E., S. Wang, C. Li, X. Wang, K. Chen & L. Chen, 2014. Transcriptome sequencing revealed the genes and pathways involved in salinity stress of Chinese mitten crab, *Eriocheir sinensis*. *Physiological Genomics*. <https://doi.org/10.1152/physiolgenomics.00191.2013>.
- Lister, R., B. D. Gregory & J. R. Ecker, 2009. Next is now: new technologies for sequencing of genomes, transcriptomes, and beyond. *Current Opinion in Plant Biology* 12(2): 107–118.
- Liu, S., G. Chen, H. Xu, W. Zou, W. Yan, Q. Wang, H. Deng, H. Zhang, G. Yu, J. He & S. Weng, 2017. Transcriptome analysis of mud crab (*Scylla paramamosain*) gills in response to Mud crab reovirus (MCRV). *Fish & Shellfish Immunology* 60: 545–553.
- Liu, Y., M. Hui, Z. Cui, D. Luo, C. Song, Y. Li & L. Liu, 2015. Comparative transcriptome analysis reveals sex-biased gene expression in juvenile Chinese Mitten Crab *Eriocheir sinensis*. *PLoS ONE* 10(7): e0133068.
- Liu, L., Y. Li, S. Li, N. Hu, Y. He, R. Pong, D. Lin, L. Lu & M. Law, 2012. Comparison of next-generation sequencing systems. *Journal of Biomedicine and Biotechnology* 2012: 11.
- Liu, Y., J. Zhou & K. P. White, 2014. RNA-seq differential expression studies: more sequence or more replication? *Bioinformatics* 30(3): 301–304.
- Lo, C. C. & P. S. Chain, 2014. Rapid evaluation and quality control of next generation sequencing data with FaQCs. *BMC bioinformatics* 15(1): 366.



- Love, M. I., W. Huber & S. Anders, 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* 15(12): 550.
- Lu, X., J. Kong, S. Luan, P. Dai, X. Meng, B. Cao & K. Luo, 2016. Transcriptome analysis of the hepatopancreas in the Pacific White Shrimp (*Litopenaeus vannamei*) under acute Ammonia stress. *PLoS ONE* 11(10): e0164396.
- Lv, J., P. Liu, B. Gao, Y. Wang, Z. Wang, P. Chen & J. Li, 2014. Transcriptome analysis of the *Portunus trituberculatus*: de novo assembly, growth-related gene identification and marker discovery. *PLoS One* 9(4): e94055.
- Lv, J., P. Liu, Y. Wang, B. Gao, P. Chen & J. Li, 2013. Transcriptome analysis of *Portunus trituberculatus* in response to salinity stress provides insights into the molecular basis of osmoregulation. *PLoS ONE* 8(12): e82155.
- Lv, J., L. Zhang, P. Liu & J. Li, 2017. Transcriptomic variation of eyestalk reveals the genes and biological processes associated with molting in *Portunus trituberculatus*. *PLoS ONE* 12(4): e0175315.
- MacManes, M. D., 2014. On the optimal trimming of high-throughput mRNA sequence data. *Frontiers in Genetics* 5: 13.
- Macharia, R. W., F. L. Ombura & E. O. Aroko, 2015. Insects RNA profiling reveals absence of hidden break in 28S Ribosomal RNA molecule of Onion Thrips, *Thrips tabaci*. *Journal of Nucleic Acids* 2015: 8.
- Marguerat, S. & J. Bähler, 2010. RNA-seq: from technology to biology. *Cellular and Molecular Life Sciences* 67(4): 569–579.
- Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet journal* 17(1): 10–12.
- McCarthy, S. D., M. M. Dugon & A. M. Power, 2015. ‘Degraded’ RNA profiles in Arthropoda and beyond. *PeerJ* 3: e1436.
- Meng, X.-L., P. Liu, F.-L. Jia, J. Li & B.-Q. Gao, 2015. De novo transcriptome analysis of *Portunus trituberculatus* ovary and testis by RNA-Seq: identification of genes involved in gonadal development. *PLoS ONE* 10(6): e0128659.
- Metzker, M. L., 2010. Sequencing technologies—the next generation. *Nature Reviews Genetics* 11(1): 31–46.
- Miller, J. R., S. Koren & G. Sutton, 2010. Assembly algorithms for next-generation sequencing data. *Genomics* 95(6): 315–327.
- Misner, I., C. Bicep, P. Lopez, S. Halary, E. Baptiste & C. E. Lane, 2013. Sequence comparative analysis using networks: software for evaluating *de novo* transcript assembly from Next-Generation Sequencing. *Molecular Biology and Evolution* 30(8): 1975–1986.
- Moshtaghi, A., M. L. Rahi, P. B. Mather & D. A. Hurwood, 2017. Understanding the Genomic Basis of Adaptive Response to Variable Osmotic Niches in Freshwater Prawns: a Comparative Intraspecific RNA-Seq Analysis of *Macrobrachium australiense*. *J Hered* 108(5): 544–552.
- Moshtaghi, A., M. L. Rahi, V. T. Nguyen, P. B. Mather & D. A. Hurwood, 2016. A transcriptomic scan for potential candidate genes involved in osmoregulation in an obligate freshwater palaemonid prawn (*Macrobrachium australiense*). *PeerJ* 4: e2520.
- Mykles, D. L., K. G. Burnett, D. S. Durica, B. L. Joyce, F. M. McCarthy, C. J. Schmidt & J. H. Stillman, 2016. Resources and recommendations for using transcriptomics to address grand challenges in comparative biology. *Integrative and Comparative Biology* 56(6): 1183–1191.
- Mykles, D. L. & J. H. Hui, 2015. *Neocaridina denticulata*: a decapod crustacean model for Functional Genomics. *Integrative and Comparative Biology* 55(5): 891–897.
- Nguyen, C., T. G. Nguyen, L. Van Nguyen, H. Q. Pham, T. H. Nguyen, H. T. Pham, H. T. Nguyen, T. T. Ha, T. H. Dau & H. T. Vu, 2016. De novo assembly and transcriptome characterization of major growth-related genes in various tissues of *Penaeus monodon*. *Aquaculture* 464: 545–553.
- Niedringhaus, T. P., D. Milanova, M. B. Kerby, M. P. Snyder & A. E. Barron, 2011. Landscape of Next-Generation Sequencing technologies. *Analytical Chemistry* 83(12): 4327–4341.
- Olsvik, P. A., B. T. Lunestad, A. L. Agnalt & O. B. Samuelsen, 2017. Impact of teflubenzuron on the rockpool shrimp (*Palaemon elegans*). *Comparative Biochemistry and Physiology Toxicology & Pharmacology* 201: 35–43.
- Ozsolak, F. & P. M. Milos, 2011. RNA sequencing: advances, challenges and opportunities. *Nature Reviews Genetics* 12(2): 87–98.
- Parkhomchuk, D., T. Borodina, V. Amstislavskiy, M. Banaru, L. Hallen, S. Krobitsch, H. Lehrach & A. Soldatov, 2009. Transcriptome analysis by strand-specific sequencing of complementary DNA. *Nucleic Acids Research* 37(18): e123.
- Parra, G., K. Bradnam & I. Korf, 2007. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics* 23(9): 1061–1067.
- Patel, R. K. & M. Jain, 2012. NGS QC Toolkit: a Toolkit for quality control of next generation sequencing data. *PLoS ONE* 7(2): e30619.
- Patro, R., G. Duggal, M. I. Love, R. A. Irizarry & C. Kingsford, 2017. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods*. <https://doi.org/10.1038/nmeth.4197>.
- Peng, J., P. Wei, B. Zhang, Y. Zhao, D. Zeng, X. Chen, M. Li & X. Chen, 2015. Gonadal transcriptomic analysis and differentially expressed genes in the testis and ovary of the Pacific white shrimp (*Litopenaeus vannamei*). *BMC Genomics* 16: 1006.
- Powell, D., W. Knibb, N. H. Nguyen & A. Elizur, 2016. Transcriptional profiling of Banana Shrimp *Fenneropenaeus merguensis* with differing levels of viral load. *Integrative and Comparative Biology*. <https://doi.org/10.1093/icb/icw029>.
- Qiao, H., H. Fu, Y. Xiong, S. Jiang, W. Zhang, S. Sun, S. Jin, Y. Gong, Y. Wang, D. Shan, F. Li & Y. Wu, 2017. Molecular insights into reproduction regulation of female Oriental River prawns *Macrobrachium nipponense* through comparative transcriptomic analysis. *Scientific reports* 7(1): 12161.
- Rahi, M. L., S. Amin, P. B. Mather & D. A. Hurwood, 2017. Candidate genes that have facilitated freshwater adaptation by palaemonid prawns in the genus *Macrobrachium*: identification and expression validation in a model species (*M. koomboolomba*). *PeerJ* 5: 2977.
- Rajkumar, A. P., P. Qvist, R. Lazarus, F. Lescai, J. Ju, M. Nyegaard, O. Mors, A. D. Børglum, Q. Li & J. H. Christensen, 2015. Experimental validation of methods for

- differential gene expression analysis and sample pooling in RNA-seq. *BMC Genomics* 16(1): 548.
- Rao, R., Y. B. Zhu, T. Alinejad, S. Tiruvayipati, K. L. Thong, J. Wang & S. Bhassu, 2015. RNA-seq analysis of *Macrobrachium rosenbergii* hepatopancreas in response to *Vibrio parahaemolyticus* infection. *Gut Pathogens* 7(1): 1.
- Ren, Q. & L. Pan, 2014. Digital gene expression analysis in the gills of the swimming crab (*Portunus trituberculatus*) exposed to elevated ambient ammonia-N. *Aquaculture* 434: 108–114.
- Reuter, Jason A., D. V. Spacek & Michael P. Snyder, 2015. High-throughput sequencing technologies. *Molecular Cell* 58(4): 586–597.
- Roberts, A. & L. Pachter, 2012. Streaming fragment assignment for real-time analysis of sequencing experiments. *Nature Methods* 10: 71.
- Robertson, G., J. Schein, R. Chiu, R. Corbett, M. Field, S. D. Jackman, K. Mungall, S. Lee, H. M. Okada, J. Q. Qian, M. Griffith, A. Raymond, N. Thiessen, T. Cezard, Y. S. Butterfield, R. Newsome, S. K. Chan, R. She, R. Varhol, B. Kamoh, A.-L. Prabhu, A. Tam, Y. Zhao, R. A. Moore, M. Hirst, M. A. Marra, S. J. M. Jones, P. A. Hoodless & I. Birol, 2010. De novo assembly and analysis of RNA-seq data. *Nature Methods* 7(11): 909–912.
- Robinson, M. D., D. J. McCarthy & G. K. Smyth, 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26(1): 139–140.
- Robles, J. A., S. E. Qureshi, S. J. Stephen, S. R. Wilson, C. J. Burden & J. M. Taylor, 2012. Efficient experimental design and analysis strategies for the detection of differential expression using RNA-Sequencing. *BMC Genomics* 13(1): 484.
- Rotllant, G., T. V. Nguyen, V. Sbragaglia, L. Rahi, K. J. Dudley, D. Hurwood, T. Ventura, J. B. Company, V. Chand, J. Aguzzi & P. B. Mather, 2017. Sex and tissue specific gene expression patterns identified following *de novo* transcriptomic analysis of the Norway lobster, *Nephrops norvegicus*. *BMC Genomics* 18: 622.
- Sagi, A., R. Manor & T. Ventura, 2013. Gene silencing in Crustaceans: from basic research to biotechnologies. *Genes* 4(4): 620.
- Sbragaglia, V., F. Lamanna, A. M. Mat, G. Rotllant, S. Joly, V. Ketmaier, H. O. de la Iglesia & J. Aguzzi, 2015. Identification, characterization, and diel pattern of expression of canonical clock genes in *Nephrops norvegicus* (Crustacea: Decapoda) eyestalk. *PLoS ONE* 10(11): e0141893.
- Schadt, E. E., S. Turner & A. Kasarskis, 2010. A window into third-generation sequencing. *Human Molecular Genetics* 19(R2): R227–R240.
- Schirmer, M., U. Z. Ijaz, R. D'Amore, N. Hall, W. T. Sloan & C. Quince, 2015. Insight into biases and sequencing errors for amplicon sequencing with the Illumina MiSeq platform. *Nucleic Acids Research* 43(6): e37.
- Schmieder, R. & R. Edwards, 2011. Quality control and pre-processing of metagenomic datasets. *Bioinformatics*. <https://doi.org/10.1093/bioinformatics/btr026>.
- Schulz, M. H., D. R. Zerbino, M. Vingron & E. Birney, 2012. Oases: robust *de novo* RNA-seq assembly across the dynamic range of expression levels. *Bioinformatics* 28(8): 1086–1092.
- Schurch, N. J., P. Schofield, M. Gierliński, C. Cole, A. Sherstnev, V. Singh, N. Wrobel, K. Gharbi, G. G. Simpson, T. Owen-Hughes, M. Blaxter & G. J. Barton, 2016. How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use? *RNA* 22(6): 839–851.
- Seelenfreund, E., W. A. Robinson, C. M. Amato, A.-C. Tan, J. Kim & S. E. Robinson, 2014. Long term storage of dry versus frozen RNA for next generation molecular studies. *PLoS ONE* 9(11): e111827.
- Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva & E. M. Zdobnov, 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31(19): 3210–3212.
- Soneson, C. & M. Delorenzi, 2013. A comparison of methods for differential expression analysis of RNA-seq data. *BMC Bioinformatics* 14(1): 91.
- Song, L., C. Bian, Y. Luo, L. Wang, X. You, J. Li, Y. Qiu, X. Ma, Z. Zhu, L. Ma, Z. Wang, Y. Lei, J. Qiang, H. Li, J. Yu, A. Wong, J. Xu, Q. Shi & P. Xu, 2016. Draft genome of the Chinese mitten crab *Eriocheir sinensis*. *GigaScience* 5: 5.
- Sookruksawong, S., F. Sun & Z. Liu, 2013. RNA-Seq analysis reveals genes associated with resistance to Taura syndrome virus (TSV) in the Pacific white shrimp *Litopenaeus vannamei*. *Developmental & Comparative Immunology* 41(4): 523–533.
- Sultan, M., V. Amstislavskiy, T. Risch, M. Schuette, S. Dökel, M. Ralser, D. Balzereit, H. Lehrach & M.-L. Yaspo, 2014. Influence of RNA extraction methods and library selection schemes on RNA-seq data. *BMC Genomics* 15(1): 675.
- Sultan, M., S. Dökel, V. Amstislavskiy, D. Wuttig, H. Sülmann, H. Lehrach & M.-L. Yaspo, 2012. A simple strand-specific RNA-Seq library preparation protocol combining the Illumina TruSeq RNA and the dUTP methods. *Biochemical and Biophysical Research Communications* 422(4): 643–646.
- Sun, M., Y. Ting Li, Y. Liu, S. Chin Lee & L. Wang, 2016. Transcriptome assembly and expression profiling of molecular responses to cadmium toxicity in hepatopancreas of the freshwater crab *Sinopotamon henanense*. *Scientific Reports* 6: 19405.
- Surget-Groba, Y. & J. I. Montoya-Burgos, 2010. Optimization of *de novo* transcriptome assembly from next-generation sequencing data. *Genome Research* 20(10): 1432–1440.
- Tarazona, S., F. García, A. Ferrer, J. Dopazo & A. Conesa, 2012. NOIseq: a RNA-seq differential expression method robust for sequencing depth biases. *EMBnet Journal* 17: 18–19.
- Tarazona, S., F. García-Alcalde, J. Dopazo, A. Ferrer & A. Conesa, 2011. Differential expression in RNA-seq: a matter of depth. *Genome Research* 21(12): 2213–2223.
- Teng, M., M. I. Love, C. A. Davis, S. Djebali, A. Dobin, B. R. Graveley, S. Li, C. E. Mason, S. Olson, D. Pervouchine, C. A. Sloan, X. Wei, L. Zhan & R. A. Irizarry, 2016. A benchmark for RNA-seq quantification pipelines. *Genome Biology* 17(1): 74.
- Ventura, T., Q. P. Fitzgibbon, S. C. Battaglene & A. Elizur, 2015. Redefining metamorphosis in spiny lobsters: molecular analysis of the phyllosoma to puerulus transition in *Sagmariasus verreauxi*. *Scientific Reports* 5: 13537.
- Ventura, T., R. Manor, E. D. Aflalo, V. Chalifa-Caspi, S. Weil, O. Sharabi & A. Sagi, 2013. Post-embryonic

- transcriptomes of the prawn *Macrobrachium rosenbergii*: multigenic succession through metamorphosis. *PLoS ONE* 8(1): e55322.
- Waiho, K., H. Fazhan, M. S. Shahreza, J. H. Z. Moh, S. Noor-baiduri, L. L. Wong, S. Sinnasamy & M. Ikhwanuddin, 2017. Transcriptome analysis and differential gene expression on the testis of orange mud crab, *Scylla olivacea*, during sexual maturation. *PLoS ONE* 12(1): e0171095.
- Wang, L., Z. Feng, X. Wang, X. Wang & X. Zhang, 2010. DEGseq: an R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26(1): 136–138.
- Wang, Z., M. Gerstein & M. Snyder, 2009. RNA-Seq: a revolutionary tool for transcriptomics. *Nature Reviews Genetics* 10(1): 57–63.
- Wang, Y., N. Ghaffari, C. Johnson, U. Braga-Neto, H. Wang, R. Chen & H. Zhou, 2011. Evaluation of the coverage and depth of transcriptome by RNA-Seq in chickens. *BMC Bioinformatics*. <https://doi.org/10.1186/1471-2105-12-S10-S5>.
- Wang, Y., Y. Xiu, K. Bi, J. Ou, W. Gu, W. Wang & Q. Meng, 2017. Integrated analysis of mRNA-seq in the haemocytes of *Eriocheir sinensis* in response to *Spiroplasma eriocheiris* infection. *Fish Shellfish Immunol* 68: 289–298.
- Wei, J., X. Zhang, Y. Yu, H. Huang, F. Li & J. Xiang, 2014. Comparative transcriptomic characterization of the early development in Pacific white shrimp *Litopenaeus vannamei*. *PLoS ONE* 9(9): e106201.
- Wilhelm, B. T. & J.-R. Landry, 2009. RNA-Seq—quantitative measurement of expression through massively parallel RNA-sequencing. *Methods* 48(3): 249–257.
- Winnebeck, E. C., C. D. Millar & G. R. Warman, 2010. Why Does Insect RNA Look Degraded? *Journal of Insect Science* 10: 159.
- Wu, T. D. & S. Nacu, 2010. Fast and SNP-tolerant detection of complex variants and splicing in short reads. *Bioinformatics* 26(7): 873–881.
- Wu, A. R., N. F. Neff, T. Kalisky, P. Dalerba, B. Treutlein, M. E. Rothenberg, F. M. Mburu, G. L. Mantalas, S. Sim, M. F. Clarke & S. R. Quake, 2014. Quantitative assessment of single-cell RNA-sequencing methods. *Nature Methods* 11(1): 41–46.
- Xie, Y., G. Wu, J. Tang, R. Luo, J. Patterson, S. Liu, W. Huang, G. He, S. Gu, S. Li, X. Zhou, T.-W. Lam, Y. Li, X. Xu, G. K.-S. Wong & J. Wang, 2014. SOAPdenovo-Trans: de novo transcriptome assembly with short RNA-Seq reads. *Bioinformatics*. <https://doi.org/10.1093/bioinformatics/btu077>.
- Yang, X., D. Liu, F. Liu, J. Wu, J. Zou, X. Xiao, F. Zhao & B. Zhu, 2013. HTQC: a fast quality control toolkit for Illumina sequencing data. *BMC bioinformatics* 14(1): 33.
- Yang, C. & H. Wei, 2015. Designing Microarray and RNA-Seq experiments for greater systems biology discovery in modern plant genomics. *Molecular Plant* 8(2): 196–206.
- Yu, Y., X. Zhang, J. Yuan, F. Li, X. Chen, Y. Zhao, L. Huang, H. Zheng & J. Xiang, 2015. Genome survey and high-density genetic map construction provide genomic and genetic resources for the Pacific White Shrimp *Litopenaeus vannamei*. *Scientific Reports* 5: 15612.
- Yuan, J., X. Zhang, C. Liu, Y. Yu, J. Wei, F. Li & J. Xiang, 2017. Genomic resources and comparative analyses of two economical penaeid shrimp species, *Marsupenaeus japonicus* and *Penaeus monodon*. *Marine Genomics*. <https://doi.org/10.1016/j.margen.2017.12.006>.
- Zerbino, D. R. & E. Birney, 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Research* 18(5): 821–829.
- Zhang, Z. H., D. J. Jhaveri, V. M. Marshall, D. C. Bauer, J. Edson, R. K. Narayanan, G. J. Robinson, A. E. Lundberg, P. F. Bartlett, N. R. Wray & Q.-Y. Zhao, 2014. A comparative study of techniques for differential expression analysis on RNA-Seq data. *PLoS ONE* 9(8): e103207.
- Zhang, D., F. Wang, S. Dong & Y. Lu, 2016. *De novo* assembly and transcriptome analysis of osmoregulation in *Litopenaeus vannamei* under three cultivated conditions with different salinities. *Gene* 578(2): 185–193.
- Zhao, Q.-Y., Y. Wang, Y.-M. Kong, D. Luo, X. Li & P. Hao, 2011. Optimizing *de novo* transcriptome assembly from short-read RNA-Seq data: a comparative study. *BMC Bioinformatics*. <https://doi.org/10.1186/1471-2105-12-S14-S2>.
- Zhao, S., Y. Zhang, W. Gordon, J. Quan, H. Xi, S. Du, D. von Schack & B. Zhang, 2015a. Comparison of stranded and non-stranded RNA-seq transcriptome profiling and investigation of gene overlap. *BMC Genomics* 16(1): 675.
- Zhao, Q., L. Pan, Q. Ren & D. Hu, 2015b. Digital gene expression analysis in hemocytes of the white shrimp *Litopenaeus vannamei* in response to low salinity stress. *Fish & Shellfish Immunology* 42(2): 400–407.
- Zhong, S., J. G. Joung, Y. Zheng, Y. R. Chen, B. Liu, Y. Shao, J. Z. Xiang, Z. Fei & J. J. Giovannoni, 2011. High-throughput illumina strand-specific RNA sequencing library preparation. *Cold Spring Harbor Protocols* 2011(8): 940–949.
- Zhou, Y. H., K. Xia & F. A. Wright, 2011. A powerful and flexible approach to the analysis of RNA sequence count data. *Bioinformatics* 27(19): 2672–2678.