

AAS-PROVIDED PDF • OPEN ACCESS

## SPHEREx Discovery of Strong Water Ice Absorption and an Extended Carbon Dioxide Coma in 3I/ATLAS

To cite this article: C. M. Lisse *et al* 2025 *Res. Notes AAS* **9** 242

Manuscript version: AAS-Provided PDF

This AAS-Provided PDF is © 2025 **The Author(s)**. Published by the American Astronomical Society.



Original content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence

<https://creativecommons.org/licenses/by/4.0>

Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required.

View the [article online](#) for updates and enhancements.

## **SPHEREx Discovery of Strong Water Ice Absorption and an Extended Carbon Dioxide Coma in 3I/ATLAS**

- C.M. Lisse**, Johns Hopkins University Applied Physics Laboratory, Planetary Exploration Group, Space Department, 11100 Johns Hopkins Road, Laurel, MD 20723, USA
- Y.P. Bach**, Korea Astronomy and Space Science Institute, 776 Daedeok-daero, Lee Won Chul Hall 301, Yuseong-gu, Daejeon 34055, Republic of Korea
- S. Bryan**, Arizona State University, 650 E Tyler Mall, Tempe, AZ 85281, USA
- B.P. Crill**, California Institute of Technology/Jet Propulsion Lab, 4800 Oak Grove Drive, Pasadena, CA 91009, USA
- A. Cukierman**, Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA
- O. Doré**, California Institute of Technology/Jet Propulsion Lab, 4800 Oak Grove Drive, Pasadena, CA 91009, USA
- B. Fabinsky**, California Institute of Technology/Jet Propulsion Lab, 4800 Oak Grove Drive, Pasadena, CA 91009, USA
- A. Faisst**, Infrared Processing and Analysis Center, Caltech, 770 South Wilson Street, Pasadena, CA 91125, USA
- P. M. Korngut**, Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA
- G. Melnick**, Center for Astrophysics, Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA
- Z. Rustamkulov**, Infrared Processing and Analysis Center, Caltech, 770 South Wilson Street, Pasadena, CA 91125, USA
- V. Tolls**, Center for Astrophysics, Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA
- M. Werner**, California Institute of Technology/Jet Propulsion Lab, 4800 Oak Grove Drive, Pasadena, CA 91009, USA
- M.L. Sitko**, Space Science Institute, 4765 Walnut Street, Suite B, Boulder, CO 80301, USA
- C. Champagne**, Northern Arizona University, Department of Astronomy and Planetary Science, Flagstaff, AZ, USA
- M. Connelley**, University of Hawaii, 640 N. Aohoku Place, Hilo, HI 96720, USA
- J.P. Emery**, Northern Arizona University, Department of Astronomy and Planetary Science, Flagstaff, AZ, USA
- Y.R. Fernandez**, University of Central Florida, Department of Physics, 4000 Central Florida Boulevard, Orlando, FL 32816-2385, USA
- B. Yang**, Planetary Science Institute, 1700 East Fort Lowell, Suite 106, Tucson, AZ 85719
- and the SPHEREx Science Team**

**Abstract.** In mid-August 2025, 0.75-5.0 $\mu$ m SPHEREx imaging spectrophotometric and ancillary NASA-IRTF/SpeX 0.7-2.5 $\mu$ m low-resolution spectral observations of 3I/ATLAS were obtained. The combined spectrophotometry is dominated by features due to water ice absorption and CO<sub>2</sub> gas emission. A bright,  $\sim 3'$  radius CO<sub>2</sub> gas coma was clearly resolved, corresponding to  $Q_{\text{gas,CO}_2} \sim 9.4 \times 10^{26}$  molecules/sec. From the SPHEREx photometry, we put conservative, preliminary  $3\sigma$  upper limits on the gas production rates for H<sub>2</sub>O and CO of  $1.5 \times 10^{26}$  and  $2.8 \times 10^{26}$  molecules/sec, respectively. No obvious jet, tail, or trail structures were found in SPHEREx images. If we assume all observed 1 $\mu$ m flux is scattered light from a  $p_v = 0.04$  albedo spherical nucleus, then the radius would be  $R_{\text{nuc}} \sim 23$  km. Given the nucleus size limit  $R_{\text{nuc}} < 2.8$  km from Jewett+ 2025, we conclude  $>99\%$  of the measured SPHEREx continuum flux is from coma dust.

**1. SPHEREx Observations.** From 08-Aug- to 12-Aug-2025, the NASA SPHEREx mission (Doré+2018, Crill+2020) observed interstellar object 3I/ATLAS (hereafter 3I) from 0.75-5.0 $\mu$ m at the heliocentric distances 3.1-3.3 au. The observations consisted of 72 special pointings of 115 seconds each, with 3I centered on unique positions across SPHEREx's 6 LVF-patterned detectors. Approximately  $2''$  of object motion occurred during each integration, smaller than a SPHEREx  $6.15'' \times 6.15''$  pixel. The image position of 3I was determined to  $< 1''$  using the JPL Horizons Orbit #22, the mission software packages *skyloc* and *kete*, plus the SPHEREx Sky Simulator (Crill+2025). Photometry was obtained by locating 3I, placing a 2-pixel radius aperture at 3I's location, and subtracting the background counts from a surrounding annulus. While the SPHEREx platescale was optimal for detecting 3I's extended morphology and gas emission, a significant fraction of the pointings were contaminated by flux from background objects (galactic latitude<sub>3I</sub> was  $\sim 23^\circ$ ) and removed from further analysis until the background sky, without 3I, is re-observed in Nov-Dec 2025. For the reflectance spectra, the photometry was corrected for 3I's motion versus the Sun and Earth assuming a  $1/r_h^2 * 1/\Delta^2$  dependence. No correction for 3I's phase function, rotation, or production rate variability was applied.

**2. Ancillary Measurements.** On 11-Aug-2025, 3I was observed from the NASA/IRTF 3.2m using the SpeX imaging spectrometer (Rayner+2003) with 0.7-2.5 $\mu$ m,  $R \sim 200$  spectroscopy. Two G2V stars (HD148744 and SA110-361) were used as local and absolute solar analogue references, respectively.

**3. Spectral Results.** The combined SPHEREx and IRTF photometry/reflectance spectrophotometry is shown in Figure 1a. There is good consistency between the different

measurements and published results, showing a generally rising slope from 0.7-1.0 $\mu$ m and a flattened slope from 1-1.5 $\mu$ m. Clear IRTF/SpeX absorptions at 1.5, 2.1, and 2.4 $\mu$ m are new, although the latter is in a region of low SNR. SPHEREx confirms the downward trend at the end of the IRTF/SpeX spectrum, and continues its very strong drop through 2.5-3.1 $\mu$ m not readily observable from the ground for this  $V \sim 16.6$  magnitude object. The SPHEREx reflectance drops to  $\sim 20\%$  of its 1 $\mu$ m flux level by 3 $\mu$ m, and stays low out to 5 $\mu$ m except for a strong peak located at  $\sim 4.3\mu$ m. The combined reflectance spectral shape is consistent with the CO<sub>2</sub>/H<sub>2</sub>O ice+organics KBO “Cliff” category of Pinilla-Alonso+2025 (Figure 1a).

**4. Imaging Results.** A SPHEREx continuum dust image of 3I created by stacking 1.0-1.5 $\mu$ m images is shown in Figure 1b. 3I looks like a point source. SPHEREx images taken near the H<sub>2</sub>O (2.8-3.1 $\mu$ m), CO<sub>2</sub> (4.25-4.2 $\mu$ m), and CO (4.7 $\mu$ m) gas emission lines are shown in Figures 1c-e. While no obvious coma was found in H<sub>2</sub>O or CO, the CO<sub>2</sub> image shows a central condensation and detectable structure above the background with radius  $\gtrsim 3''$ , far beyond the central PSF (Figure 1f). The observed CO<sub>2</sub> coma is roughly symmetric with respect to the Sun and orbital velocity directions.

**5. Temporal Variability.** Continuum SPHEREx geometry-corrected flux measurements of 3I at 1.0-1.5 $\mu$ m were corrected for the Solar spectrum and plotted against time. We see no evidence for variability of more than 20% over the timespan 08–12 Aug 2025, giving us confidence in the construction of our preliminary 0.75-5.0 $\mu$ m spectrum.

**6. Discussion/Interpretation.** Adopting  $N_{\text{molec}} = 4\pi r_h^2 \Delta^2 F_{\text{vd}}(\lambda) / (g * h * \lambda)$  (Lisse+2009), and  $Q_{\text{gas}} = N_{\text{gas}} * (2\pi v_{\text{gas}} \rho)$ , where  $r_h = 3.2\text{au}$ ,  $\Delta = 2.6\text{au}$ , the fluorescence efficiency  $g = 2.9\text{e-}4$ ,  $2.6\text{e-}3$ , and  $2.6\text{e-}4$  for H<sub>2</sub>O 3.05 $\mu$ m, CO<sub>2</sub> 4.25 $\mu$ m, and CO 4.7 $\mu$ m emission, respectively (Crovisier+1997),  $N_{\text{gas}}$  is the column number density of gas,  $v_{\text{gas}}$  is the gas emission velocity  $0.85/\text{sqrt}(3.2\text{au}) = 0.48\text{ km/sec}$  (Cochran+2012), and  $\rho$  is the projected distance on the sky ( $\sim 11,600\text{ km/pixel}$ ), we find the estimated rate of CO<sub>2</sub> gas loss,  $Q_{\text{gas,CO}_2} \sim 9.4 \times 10^{26} \pm 30\%$  molecules/sec. Conservative, preliminary 3- $\sigma$  upper limits for  $Q_{\text{gas,H}_2\text{O}} = 1.5 \times 10^{26}$  and for  $Q_{\text{gas,CO}} = 2.8 \times 10^{26}$  molecules/sec were also found. This rate of CO<sub>2</sub> emission and upper limit for CO production is consistent with the activity of thermally processed short-period Solar System comets at 3.2au (Harrington-Pinto & Womack 2022).

From extant surveys, comets are known to be mixtures of ice and refractory dust, with H<sub>2</sub>O, CO<sub>2</sub>, and CO ices dominating the icy component (Bockelee-Morvan & Biver 2017). 3I appears to be an active object with morphology created mainly by CO<sub>2</sub> gas emission

effects. This is physically reasonable, since 3I at  $r_h \sim 3.2 \text{ au}$  was well within the Solar system's  $\text{CO}_2$  “ice line” at  $\sim 14 \text{ au}$  during the observations (Lisse+2021, 2022). The strong reflectance signature of water ice can be explained by the presence of abundant water ice in 3I's nucleus and coma dust particles. The lack of a bright water gas coma is puzzling as 3I was not far outside the Solar system's “water ice line” at  $2.5 \text{ au}$  during the observations. Most likely, 3I is emitting large chunks of mixed  $\text{CO}_2 + \text{H}_2\text{O}$  ice into its coma, where evaporative cooling of  $\text{CO}_2$  is pinning the chunks' temperature at  $\sim 120 \text{ K}$  and greatly suppressing the  $\text{H}_2\text{O}$  ice's vapor pressure (Lisse+2021).


The lack of any obvious dust coma extension at the  $6''$ -scale is consistent with the  $\sim 3''$ -wide by  $5''$ -long dust continuum morphology reported by Jewitt+2025. Assuming all observed  $1\text{-}\mu\text{m}$  flux is scattered light from the nucleus and an  $p_v = 0.04$  albedo,  $\phi(a) = 0.035 \text{ mag/deg}$  surface, the radius  $R_{\text{nuc}}$  of a spherical nucleus emitting  $F_{\text{scat}} = 1.5 \pm 0.05 \text{ mJy} = (p_v * 10^{-0.4 * \phi(a)} * \pi R_{\text{nuc}}^2) (F_{\text{sun}}/\pi)/(r_h^2 \Delta^2)$  (Lisse+1999) at  $\lambda = 1 \mu\text{m}$  is  $23 \pm 0.7 \text{ km}$ . This is  $\sim 10$  times the nucleus size limit found by Jewitt+2025 using HST imaging, arguing that  $>99\%$  of the observed SPHEREx flux is due to scattering from icy coma dust.

**7. Future Work.** The photometric measurements reported here are from the first  $\frac{1}{2}$  of all the August 2025 SPHEREx 3I pointings. Treatment of stellar contamination will greatly improve after the spacecraft re-observes the same field in 6 months. 3I/ATLAS will also pass through SPHEREx's planned survey pattern again in November-December 2025.

## 8. References

- Bockelée-Morvan,D & Biver,N. 2017, *Phil. Trans. R.Soc.A* **375**:20160252
- Cochran,A.+ 2012, *Icarus* **218**:144
- Crill,B.+ 2025, <https://arxiv.org/abs/2505.24856>
- Crill,B.+ 2020, *Proceed. SPIE*, 11443:114430I
- Crovisier,J.+ 1997, *Science* **275**:1904
- Doré,O.+ 2018, arXiv:1805.05489
- Harrington-Pinto,O.+ 2022, *PSJ* **3**:247
- Huebner,W.+ 1992, *Astrophysics & Space Science* **195**:1
- Jewitt,D.+ 2025, <https://arxiv.org/pdf/2508.02934>
- Lisse,C.+ 1999, *Icarus* **140**:189
- Lisse,C.+ 2009, *ApJ* **701**:2019
- Lisse,C.+ 2021, *Icarus* 356:114072
- Lisse,C+ 2022, *PSJ* **3**:112
- Pinilla-Alonso, N.+, 2025, *Nature Astron* **9**:230
- Rayner,J.+ 2003, *PASP* **115**:3

## 9. Figure



**Figure 1a : SPHEREx 0.75-5.0 $\mu$ m 2-pixel radius aperture spectrophotometry (left) and solar spectrum corrected reflectance (right) for 3I.** Red arrows denote possible significant background contamination due to known background sources. The arrows start at the measured flux+1 $\sigma$ . For the reflectance, the NASA-IRTF/SpeX 0.7-2.5 $\mu$ m spectrum (**green**) is overlaid. The two sets of measurements are consistent, and characterized by a slightly rising slope from 0.7-1.0 $\mu$ m, a flat regime from 1.0-1.5 $\mu$ m, a shallow absorption at 1.5 and 2.1 $\mu$ m, and a strong falloff starting at 2.4 $\mu$ m. The SPHEREx spectrophotometry stays low from 2.5-5.0 $\mu$ m, with a huge uptick at 4.3 $\mu$ m due to CO<sub>2</sub> gas emission. Also overlaid for comparison is the average “cliff” KBO spectrum of Pinilla-Alonso+2025 (**pink**). **Figure 1b: Stacked 1.0-1.5 $\mu$ m SPHEREx geometry- and Solar flux-corrected image of 3I.** Assuming the reflectance is flat across these wavelengths, this median stacking produces a deep image of the nucleus + dust scattered light continuum flux from 3I. There is no significant extension found in the stacked image beyond a stellar point source in a SPHEREx 6.15” pixel. **Figures 1c, d, and e: SPHEREx images centered at the H<sub>2</sub>O/CO<sub>2</sub>/CO coma gas wavelengths of 3.0/4.26/4.7 $\mu$ m, respectively.** 3I is undetected in H<sub>2</sub>O and CO, with 2-pixel radius preliminary conservative upper limits of 0.45 and 0.99mJy. By contrast, a bright CO<sub>2</sub> coma with central PSF flux = 33mJy that extends out to at least 30-pixel (348,000 km) was found. This observed minimum size for the CO<sub>2</sub> coma corresponds to  $\sim 700,000$ -second mean lifetime at  $v_{\text{gas}} = 0.48$  km/sec. The reported photodissociation lifetime for CO<sub>2</sub> at 1au is 135hrs = 486,000 seconds (Huebner+1992), which scales by  $1/r_h^2$  to  $\sim 4,670,000$  seconds at 3.2au. **Figure 1f: Azimuthally-averaged radial profile for the SPHEREx 3I CO<sub>2</sub> coma.** The observed falloff with projected distance is steeper than  $1/\rho$ , closer to  $1/\rho^{3/2}$ , which is consistent with a near-nucleus source of CO<sub>2</sub> coupled with photodissociation + charge exchange ionization processing of free-flying molecules in the coma.